

Catalogue of the Cerambycidae (Coleoptera) of Canada and United States of America. Part I. Subfamilies Parandrinae, Prioninae and Spondylidinae

June 2022

MIGUEL A. MONNÉ

Museu Nacional, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista, 20940-040, Rio de Janeiro, RJ, Brazil.

monne@uol.com.br

EUGENIO H. NEARNS

National Museum of Natural History, Smithsonian Institution Washington, DC 20560 USA

Recommended format for citation: Available from:

https://cerambycids.com/catalog/Monne&Nearns_Jun2022_NearcticCat_part_I.pdf

This catalogue is an attempt to offer the names, the geographical distribution and bibliographic references for extant taxa of Cerambycidae of Canada and United States of America.

All informations of the present catalogue were previously published, there is not unpublished data. The geographical distribution of the extant families and subfamilies in the North America is based on previously published records, supplemented by data extracted from several collections. The geographical distribution of the species from Canada were adapted from Bousquet, Laplante, Hammond and Langor, 2017. Tribes are listed in alphabetical order, and in each tribe genera and species follow the same order. Each species-group name is followed by author (s), publication year, page, and figure (if any), acronym of the institution depository of the type, host plants (when available) and when not strictly taxonomical, an abbreviated indication of the matter is given as:

ab. - aberration
 biol. - biological data
 cat. - catalogue
 distr. - distribution
 ecol. - ecological data
 emend. -emendation
et al., - more than four authors
 lect. - lectotype (designation).
 mim. - mimetism
 paras. - parasites
 pred. - predators
 refs - bibliographic references
 rev. - revision
 reval. - revalidation

syn. - synonymy

terat - teratology

var. - variety

Acronyms of the institutions or private collections mentioned in the Part III of the catalogue:

AMNH - American Museum of Natural History, New York, New York, United States.

ANSP - Academy of Natural Sciences, Philadelphia, Pennsylvania, United States.

BMNH - The Natural History Museum, London, United Kingdom.

CASC - California Academy of Sciences, San Francisco, California, United States.

CNCI - Canadian National Collection of Insects, Ontario, Ottawa, Canada.

EMEC - Essig Museum of Entomology, University of California, Berkeley, California, United States.

FMNH - Field Museum of Natural History, Chicago, Illinois, United States

IUBC - Department of Zoology collection, Indiana University, Bloomington, United States

LACM - Los Angeles County Museum of Natural History, Los Angeles, California, United States.

LSUK - Linnean Society, London, United Kingdom.

MCZN - Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, United States.

MNHN - Muséum National d'Histoire Naturelle, Paris, France

NHMW - Naturhistorisches Museum, Wien, Austria.

NHRS - Naturhistoriska Riksmuseet, Stockholm, Sweden

USNM - National Museum of Natural History, Washington, D. C., United States.

UZIU - Uppsala University, Uppsala, Sweden.

ZMHB - Museum für Naturkunde der Humboldt-Universität, Berlin, Germany.

ZMUM - Zoological Museum, Moscow State University, Moscow, Russia.

PARANDRINAE Blanchard, 1845

Parandrides Blanchard, C.E., 1845: 134.

Type-genus: *Parandra* Latreille, 1802.

Type-species: *Attelabus glaber* De Geer, 1774 by monotypy. Availability (under Article 11.7.2).

Parandraeidae; Gistel 1848a: (8).

PARANDRINI Blanchard, 1845

Parandrides Blanchard, C. E., 1845: 134.

Parandrae; Lameere, 1912b: 180; 1913: 4 (cat.); 1919: 4.

Parandrini; Gilmour, 1956: 2; Monné, M.A., 1994: 1 (cat.); Santos-Silva, 2002b: 30 (key gen.); Monné, M.A., 2006: 8 (cat.); Santos-Silva & Martins, 2010: 15; Cardona-Duque, Santos-Silva & Wölf, 2010: 138 (key gen.); Bousquet, Heffern, Bouchard & Nearn., 2009: 14; Santos-Silva, Heffern & Matsuda, 2010: 6 (key genera); Bouchard *et al.*, 2011: 454.

Type-genus: *Parandra* Latreille, 1802.

Type-species: *Attelabus glaber* De Geer, 1774 by monotypy. Availability (under Article 11.7.2): Parandrae Blanchard, 1845 (Lameere 1913: 4).

***Neandra* Lameere, 1912**

Parandra (*Neandra*) Lameere, 1912b: 114; 1919: 15; Linsley, 1962: 5; Hatch, 1971: 91; Monné, M.A., 1994: 3 (cat.); Chemsak, 1996: 7; Santos-Silva, 2001: 239; 2002: 30; Bousquet, 2008: 62

Neandra; Santos-Silva, 2002: 30; Monné, M.A. & Hovore, 2006: 5 (checklist); Bousquet, 2008: 62; Santos-Silva, Heffern & Matsuda, 2010: 6; Svacha & Lawrence, 2014: 138; Bousquet, Laplante, Hammond & Langor, 2017: 28

Type species - *Tenebrio brunneus* Fabricius (monotypy)

1. *Neandra brunnea* (Fabricius, 1798)

Syntypes locality - Syntypes: Indiis. (ZMUC). **Distribution** - Eastern half of the United States and southeast Canada. **Host plants** - *Acer rubrum* Linnaeus (Aceraceae), *Castanea dentata* (Marshall) Borkhausen, *C. pumila* (Linnaeus) Miller, *Fagus* sp., *Quercus velutina* Lamarck (Fagaceae), *Carya* sp., *Juglans* sp. (Juglandaceae), *Liriodendron tulipifera* Linnaeus (Magnoliaceae), *Pinus rigida* Miller (Pinaceae), *Prunus cerasus* Linnaeus (Rosaceae), *Populus nigra* Linnaeus, *Salix alba* Linnaeus (Salicaceae), *Ailanthus glandulosa* Desfontaines (Simarubaceae) *Tilia americana* Linnaeus, *T. cordata* Miller, *T. platyphyllos* Scopoli (Tiliaceae)

Tenebrio brunneus Fabricius, 1798: 49; 1801: 148; Zimsen, 1964: 46

Parandra brunnea; Schoenherr, 1817: 334; Lepeletier & Audinet-Serville, 1825: 3; Harris, 1838: 88; White, 1853: 2; Horn, 1861: 43; Bland, 1861: 93; Thomson, 1861b: 84; Osten-Sacken, 1862: 118, pl. 2, fig. 6 (larva); Murray, 1862: 452, pl. 47, fig. 7; Thomson, 1867b: 113; Lacordaire, 1868: 23; Pettit, 1871: 105; Packard, 1872: 494; Horn, 1875: 150; Popenoe, 1877: 33; Provancher, 1877: 582; Schaupp, 1879: 24 (hosts); Riley, 1880: 23; Packard, 1881: 127; Townsend, 1886: 66; Packard, 1890: 481; Townsend, 1893: 202; Hopkins, 1893: 191; Chittenden, 1894: 96; Hamilton, 1895: 337; Smith, 1900: 284; Lameere, 1902a: 80; Dury, 1902: 158; Ulke, 1903: 25; Chagnon, 1905: 36; Tucker, 1907: 162; Fall & Cockerell, 1907: 190; Wickham, 1909: 28; Champlain, 1909: 181; Blatchley, 1910: 1007; Snyder, 1910: 1; Smith, 1910: 323; Gahan, 1911: 288; Casey, 1912: 216; Snyder, 1913: 131, fig. 12; Johnson, 1915: 314; Brooks, 1915: 1; Craighead, 1915: 12, pl. 3, fig. 6; pl. 4, figs 5.6; Chagnon, 1917: 229; Morris, 1918: 42; Nicolay, 1919: 20; Britton, 1920: 266, pl. 16; Craighead, 1923, 29, pl. 6, fig. 4; Hatch, 1925: 579; Fletcher, 1926: 143; Kirk & Knull, 1926: 21; Leonard, 1928: 432; Schaeffer, 1929: 39; Craighead & Middleton, 1930: 13; Barrett, 1932: 291; Beaulne, 1932: 197; Loding, 1933: 148; Herrick, 1935: 42, fig. 18; Sheppard, 1935: 97; Doane *et al.*, 1936: 161; Sheppard, 1936: 75; Chagnon, 1936: 239; Brimley, 1938: 209; Loding, 1945: 112; Knull, 1946: 144; Craighead, 1950: 258; Beal, Haliburton & Knight, 1952: Duffy, 1953: 102, figs 24-29; Gilmour, 1954: 3; Anderson, 1960: 282, fig. 11.4; Nüssler, 1961: 130 (introduction); Chagnon & Robert, 1962: 239; Swan & Papp, 1972: 441, fig. 924; Baker, 1972: 197; Furniss & Carolin, 1977: 287; Waters & Hyche, 1984: 284; White, 1985: 281; Gosling, 1986: 157; Bense, 1995: 96, fig. 288

Isocerus brunneus; Illiger, 1802: 295:

Parandra (Neandra) brunnea; Lameere, 1912b: 114; 1919: 16 (syn.); Hatch, 1971: 92; Chemsak, Linsley & Noguera, 1992: 12 (cat.); Chemsak, 1996: 8, pl. 1, figs 3, 4; Schiefer, 1998: 115; Heffern, 1998: 6; Sama, 2003: 8, 2 figs

Parandra brunnea brunnea; Linsley, 1957: 1; Gosling, 1973: 65; Kirk & Balsbaugh, 1975: 96; Stein & Tagestad, 1976: 27; Laliberté, Chantal & LaRochelle, 1977: 96; Turnbow & Wappes, 1978: 367; Turnbow & Franklin, 1980: 338; Gosling, 1984: 72; MacRae, 1993: 26; Yanega, 1996: 25, pl. 1, figs 1a, 1b; Linsley & Chemsak, 1997: 414 (hosts); Vlasák & Vlasakova, 2002: 204

Parandra (Neandra) brunnea brunnea; Linsley, 1962: 8; Monné, M.A., 1994: 3 (cat.); Peck & Thomas, 1998: 116

Neandra brunnea brunnea; Santos-Silva, 2002: 31

Neandra brunnea; Ferrer, Barclay & Hancock, 2004: 579, figs 6-10 (syntypes); Monné, M.A. & Hovore, 2006: 5 (checklist); Webster, McCorquada & Majka, 2009: 291; Bouyer, Drumont & Santos-Silva, 2012: 97; Biffi & Fuhrmann, 2013: 13; Holt, 2013: 242; Bouchard, 2014: 543; Klingeman *et al.*, 2017: 294; Bousquet, Laplante, Hammond & Langor, 2017: 18, pl. 1; Haack, Keena & Eyre, 2017: 76; Webster, 2016: 484; Haack, 2017: 118; Rice, MacRae & Merickel, 2017: 669

Tenebrio purpureus Herbst, 1799: 21, pl. 119, fig. 2

Type locality - Holotype: without locality (ZMHB)

Parandra Sayi Thomson, 1867b: 114; Lacordaire, 1868: 23; Thomson, 1878: 4

Syntypes locality - Syntypes: Amer. bor. (MNHN)

Parandra quadricollis Thomson, 1867: 114; Lacordaire, 1868: 23; Thomson, 1878: 4

Type locality – Holotype : Amer. Borealis. (MNHN)

Parandra conformis Thomson, 1867b: 115; Lacordaire, 1868: 23; Thomson, 1878: 4

Type locality - Holotype: Amer. Borealis. (MNHN)

Parandra dentata Thomson, 1867b: 115; Lacordaire, 1868: 23; Thomson, 1878: 4

Type locality 8 Holotype female; Amer. Borealis. (MNHN)

Parandra minuta Thomson, 1867b: 116; Lacordaire, 1868: 23; Thomson, 1878: 4

Type locality - Holotype female: Amer. Borealis. (MNHN)

Parandra ampliceps Casey, 1912: 216 ; Lingafelter *et al.*, 2014: 39, 369, fig. 36u (lect.)

Type locality - Lectotype male: United States, Indiana. (USNM)

Parandra gravidula Casey, 1912: 216 ; Lingafelter *et al.*, 2014: 71, 369, fig. 36w (lect.)

Type locality - Lectotype male: United States : Pennsylvania (USNM)

Parandra brunnea coloradensis Linsley, 1957: 2

Type locality - Holotype male: United States, Colorado. (AMNH)

Parandra brunnea quebecensis Abdullah, 1968: 75

Type locality - Holotype: Canada, Quebec, Morgans Arboretum. (depository unknown)

2. *Neandra marginicollis* (Schaeffer, 1929)

Type locality - United States, California: Los Angeles County. (MCZN). **Distribution** – United States (California and southern Arizona). **Host plants** - *Alnus oblongifolia* Torrey, *A. rhombifolia* Nuttall (Betulaceae), *Quercus agrifolia* Née (Fagaceae), *Juglans regia* Linnaeus (Juglandaceae), *Platanus racemosa* Nuttall ex Audubon (Platanaceae), *Malus* sp. (Rosaceae).

Parandra marginicollis Schaeffer, 1929: 40; Doane *et al.*, 1936: 162; Moore, 1937: 86; Swan & Papp, 1972: 441, fig. 926

Parandra (Neandra) marginicollis; Linsley, 1962: 9, fig. 3; Chemsak, Linsley & Noguera, 1992: 13 (cat.); Chemsak, 1996: 10

Parandra marginicollis marginicollis; Hovore & Giesbert, 1976: 349; Linsley & Chemsak, 1997: 414

Neandra marginicollis; Santos-Silva, 2002: 3; Monné, M.A., & Hovore, 2006: 5 (checklist)

Parandra punctillata Schaeffer, 1929: 40; Doane *et al.*, 1936: 162; Linsley, Knull & Statham, 1961: 3; Lingafelter *et al.*, 2014: 396, fig. 149c (holotype)

Parandra (Neandra) marginicollis punctillata; Chemsak, Linsley & Noguera, 1992: 13

Parandra marginicollis punctillata; Furniss & Carolin, 1977: 288, fig. 170

Type locality – United States, Arizona, Santa Catalina Mountains and Prescott. (USNM)

PRIONINAE Latreille, 1802

Prionii Latreille, 1802: 212.

Type-genus: *Prionus* Geoffroy, 1762

Type-species: *Cerambyx coriarius* Linnaeus, 1758 (see ICZN 1994: 60).

CALLIPOGONINI Thomson, 1861

Callipogonitae Thomson, 1861a: 293, 323.

Callipogonides; Lacordaire, 1868: 91.

Callipogonines; Lameere, 1904a: 7.

Callipogonini; Lameere, 1912b: 181; 1919: 63; Melzer, 1919: 79; Heyrovsky, 1955: 68; Gilmour, 1956: 111; Monné, M.A., 2006: 28 (cat.). Bousquet *et al.*, 2009: 15; Bouchard *et al.*, 2011: 456.

Type-genus: *Callipogon* Audinet-Serville, 1832

Type-species: *Prionus barbatus* Fabricius, 1781 (monotypy).

Anacanthitae Thomson, 1864: 285.

Type-genus: *Anacanthus* Audinet-Serville, 1832 (junior homonym of *Anacanthus* Gray, 1831 [Pisces])

Type-species: *Anacanthus costatus* Audinet-Serville, 1832 (monotypy). Comment. This name is permanently invalid because it is based on a preoccupied type genus (Article 39).

Enoploceritae Thomson, 1864: 290.

Type-genus: *Enoplocerus* Audinet-Serville, 1832

Type-species: *Prionus armillatus* Fabricius, 1775 (monotypy).

Orthomegitae Thomson, 1864: 294.

Type-genus: *Orthomegas* Audinet-Serville, 1832

Type-species: *Cerambyx corticinus* Olivier, 1790 designated by Desmarest (1860: 307).

Ctenoscelitae Thomson, 1864: 295.

Ctenoscelinae Pascoe, 1869: 661, 663.

Type-genus: *Ctenoscelis* Audinet-Serville, 1832

Type-species: *Prionus ater* Olivier, 1795 designated by Thomson (1864: 297).

Ergatites Fairmaire in Jacquelin DuVal, 1864: 117, 191.

Ergatides; Lacordaire, 1868: 93.

Ergatini; LeConte, 1873b: 286; LeConte & Horn, 1883: 271; Leng, 1884: 8; Linsley, 1962: 24.

Type-genus: *Ergates* Audinet-Serville, 1832.

Type-species: *Prionus serrarius* Panzer, 1793 (monotypy). Availability (under Article 11.7.2).

***Trichocnemis* LeConte, 1851**

Trichocnemis LeConte, 1851: 110; Thomson, 1864: 298; Casey, 1912: 221; Swift, Santos-Silva & Nearn, 2010: 39; Monné, M. A., 2012: 128.

Ergates (*Trichocnemis*); Casey, 1890: 490; 1891: 20; Lameere, 1904a: 46; 1919: 81.

Type-species - *Trichocnemis spiculatus* LeConte, 1851 (monotypy).

Ergates; LeConte & Horn, 1883: 271; Leng, 1884: 8; Horn, 1891: 41; Linsley, 1962: 24; Arnett, 1962: 855; Hatch, 1971: 92; Monné, M.A., 2006: 37 (cat.).

1. *Trichocnemis pauper* (Linsley, 1957)

Type locality - Holotype female: United States, California: Tulare County. (CASC).

Distribution - Sierra Nevada and Coast Range mountains of California. **Host plants** – *Quercus agrifolia* Née, *Q. chrysolepis* Liebman, *Q. kelloggii* Newberry, *Q. wislizenii* A. de Candolle (Fagaceae)

Ergates pauper Linsley, 1957: 5; 1962: 29; Tyson, 1966: 203 (hosts); 1967: 125 (larva, pupa); Leech, 1968: 86(distr.); Hovore & Giesbert, 1976: 350 (hosts); Chemsak, Linsley & Noguera, 1992: 17 (cat.); Monné, M.A., & Giesbert, 1994: 9 (checklist); Monné, M.A., 1995: 11 (cat.); Chemsak, 1996: 89, fig. 11; Linsley & Chemsak, 1997: 373 (hosts); Monné, M.A., & Hovore, 2006: 5 (checklist)

Trichocnemis pauper; Swift, Santos-Silva & Nearn, 2010: 42, 2 figs

2. *Trichocnemis spiculatus spiculatus* LeConte, 1851

Type locality - Holotype male: United States, California. (MCZN). **Distribution** - Pacific Coast from British Columbia to southern California, northern Rocky Mountains and Mexico (Baja California, Durango). **Host plants** - *Abies concolor* Gordon & Glen, *A. magnifica* A. Murray, *Pinus contorta* Douglas ex Loudon, *P. insignis* Douglas ex Loudon, *P. jeffreyi* Balfour, *P. ponderosa* Douglas ex Lawson & P. Lawson, *P. tuberculata* Gordon, *Pseudotsuga menziesii* (Mirbel) Franco, *P. mucronata* Rafinesque (Pinaceae), *Sequoia sempervirens* (D. Don) Endlicher (Taxodiaceae).

Trichocnemis spiculatus LeConte, 1851: 110; Thomson, 1864: 298; Swift, Santos-Silva & Nearn, 2010: 43, figs 3-8.

Macrotoma spiculata; White, 1853: 40.

Ergates spiculatus; LeConte, 1854c: 218; 1857: 59, pl. 2, fig. 9a; Lacordaire, 1868: 96; LeConte, 1873b: 286; Snow, 1877: 19 (distr.); Riley, 1880: 238 (hosts); Packard, 1881: 162 (biol.); Snow, 1883: 42 (distr.); LeConte & Horn, 1883: 271; Leng, 1884: 8, pl. 2, fig. 9; Ricksecker, 1885: 97 (biol.); Rivers, 1886b: 6 (hosts); Horn, 1890: 161; 1891: 41 (syn.); Packard, 1890: 704 (hosts); Blaisdell, 1892: 34 (hosts); Beutenmuller, 1896: 73 (hosts); Wickham, 1897a: 167 (hosts); Harrington, 1899: 107 (distr.); Fall, 1901: 142 (distr.); Hopkins, 1904: 21, 35 (biol.); Wright & Coolidge, 1908: 68 (distr.); Craighead, 1915: 17, pl. 3, fig. 7, pl. 4, figs 1, 2, pl. 6, fig. 2 (larva, pupa); Garnett, 1918: 172 (distr.); Hardy & Preece, 1926: 34 (hosts);

- Essig, 1926: 448, fig. 348 (larva); Beaulne, 1932: 197 (hosts); Ingles, 1933: 59 (biol.); Doane *et al.*, 1936: 162, figs 75, 77 (biol.); Moore, 1937: 87 (distr.); Kimmey & Furniss, 1943: 23, figs 6a-b (biol.); Spencer, 1949: 20 (biol.); DeLeon, 1952: 79 (hosts); Keen, 1952: 192, fig. 89 (biol.); Duffy, 1953: 116, fig. 49 (larva, pupa); Spencer & Buckell, 1957: 30 (paras.); Hardy, 1957: D64 (distr.); Linsley, 1957: 6; Schoening & Tilden, 1959: 167 (biol.); Duffy, 1960: 64 (larva, pupa); Linsley, 1962: 25, fig. 8; Tyson, 1967: 125, figs 5, 7 (larva, pupa); Hatch, 1971: 93, pl. 9, fig. 1; Swan & Papp, 1972: 443, fig. 932; Chemsak, Linsley & Noguera, 1992: 17 (cat.); Chemsak, 1996: 86; Haack, Keena & Eyre, 2017: 76 (eggs)
- Ergates (Trichocnemis) spiculatus*; Casey, 1890: 491, pl. 4, fig. 17; 1891: 21; Lameere, 1904a: 46; 1919: 81, pl. 5, fig. 1; Nishio, 1956b: 68, pl. 5, figs 3, 4; Monné, M.A., & Giesbert, 1994: 9 (checklist);
- Ergates spiculatus spiculatus*; Linsley, 1962: 27, figs 7, 8; Tyson, 1966: 203 (hosts); Arnaud, 1968: 81 (biol.); Hovore & Giesbert, 1976: 350 (hosts); Hovore, 1988: 3 (distr.); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 87, pl. 5, figs 7, 8; Linsley & Chemsak, 1997: 374 (hosts); Monné, M.A., 2002: 7 (cat. hosts); Monné, M.A., & Hovore, 2006: 11 (checklist); Monné, M.A., 2006: 37 (cat.).
- Trichocnemis spiculatus spiculatus*; Rice, Merickel & MacRae, 2017: 669 (distr.); Bousquet. Laplante, Hammond & Langor, 2017: 28, pl 1
- Macrotoma californica* White, 1853: 37.
- Syntypes locality** - Syntypes male: United States, California. (BMNH).
- Macrotoma spiculigera* White, 1853:39.
- Type locality** - Holotype female: United States, California. (BMNH).

2a. *Trichocnemis spiculatus neomexicanus* (Casey, 1890)

- Type locality** - Holotype male: United States, New Mexico. (USNM). **Distribution** - Rocky Mountains to New Mexico, southern Arizona and northern Mexico. **Host plants** - *Quercus sideroxyla* Kunth (Fagaceae), *Pinus arizonica* Engelm ex Rothrock, *P. engelmanni* Carr. var. *blancoi* Martínez, *P. latifolia* H. Mayr, *P. leiophylla chihuahuana* (Engelm) A. E. Murray, *P. ponderosa* Douglas ex Lawson & P. Lawson (Pinaceae).
- Ergates (Trichocnemis) neomexicanus* Casey, 1890: 491, fig. 16; 1891: 20; Lameere, 1915: 54; Lingafelter *et al.*, 2014: 104, figs 115m, n (lect. designation).
- Ergates spiculatus* var. *neomexicanus*; Casey, 1893: 597.
- Trichocnemis neomexicana*; Casey, 1912: 221.
- Ergates neomexicanus*; Spieth, 1950: fig. 40.
- Ergates spiculatus neomexicanus*; Linsley, Knull & Statham, 1961: 29, fig. 8; Swan & Papp, 1972: 443; Kirk & Balsbaugh, 1975: 96 (distr.); Lewis, 1979: 22 (biol., distr.); Hovore, Penrose & Neck, 1987: 322 (hosts); Terrón, 1992: 299 (biol., distr.); Chemsak, Linsley & Noguera, 1992: 17 (cat.); Monné, M.A., & Giesbert, 1994: 9 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 88, pl. 5, figs 9, 10; Linsley & Chemsak, 1997: 374 (hosts); Heffern, 1998: 6 (distr.); Monné, M.A., 2002: 8 (cat. hosts); Monné, M.A., & Hovore, 2006: 11 (checklist) Monné, M.A., 2006: 36 (cat.).
- Ergates spiculatus*; Hamilton in Leng & Hamilton, 1896: 164; Knaus, 1904: 156 (distr.); Snow, 1906b: 170 (distr.); Fall & Cockerell, 1907: 191 (distr.).
- Ergates spiculatus* var. *marmoratus* Cockerell, 1890: 161; Horn, 1890: 161.
- Type locality** - Holotype female: United States, Colorado: Mountain Valley. (BMNH).

MACROTOMINI Thomson, 1861

- Macrotominae Thomson, 1861a: 290, 312; 1864: 298; 1865: 477.
- Macrotomides; Lacordaire, 1868: 96.
- Macrotominae; Pascoe, 1869: 661, 666.
- Macrotomiens; Lameere, 1903a: 89.
- Macrotomini; Lameere, 1912b: 180; 1913: 7 (cat.); Melzer, 1919: 27; Bradley, 1930: 227; Gilmour, 1956: 9; Linsley, 1962: 12; Villiers, 1980b: 140; Monné, M.A., 1995b: 1 (cat.); Chemsak, 1996: 70; Monné, M.A., 2006: 45 (cat.); Santos-Silva & Martins, 2006b: 398 (key

genera); Bousquet, Heffern, Bouchard & Nearn, 2009: 17; Bouchard *et al.*, 2011: 457; Santos-Silva & Wappes, 2012: 4 (key genera North and Central America).

Type-genus: *Macrotoma* Audinet-Serville, 1832.

Type-species: *Prionus palmatus* Fabricius, 1793 designated by Desmarest (1860: 307). Comment. *Macrotoma* Audinet-Serville, 1832 [July] is a junior homonym of *Macrotoma* Laporte, 1832 [April], a junior synonym of *Longina* Wiedemann, 1830 (Diptera). Heffern *et al.* (2006) applied the reversal of precedence (Article 23.9) to qualify *Macrotoma* Audinet-Serville of *nomen protectum*.

ARCHETYPINA Lameere, 1912

Archetypina Lameere, 1912b: 180.

Archetypina; Bousquet, Heffern, Bouchard & Nearn, 2009: 17; Bouchard *et al.*, 2011: 457.

Type genus: *Archetypus* Thomson, 1861.

Type-species: *Archetypus parandroides* Thomson, 1861 (subsequent designation, Thomson, 1864: 307).

Strongylaspis Thomson, 1861

Strongylaspis Thomson, 1861a: 313; 1864: 299; 1865: 477; Lacordaire, 1868: 100; Bates, 1879: 6; Lameere, 1903: 24; 1919: 23; Melzer, 1919: 28; Bradley, 1930: 227; Zayas, 1957: 160; Linsley, 1962: 12; Arnett, 1962: 855; Zajciw, 1970: 1; Cerda, 1974: 41; Zayas, 1975: 19; Villiers, 1980b: 143; Chemsak, 1996: 70; Monné, M.L. & Santos-Silva, 2003: 32 (rev.); Monné, M.A., 2006: 60 (cat.); Monné, M.A. & Hovore, 2006: 14 (checklist); Santos-Silva & Durán, 2009: 353 (key spp.); Monné, M.A., 2012: 130.

Type-species - *Strongylaspis scobinatus* Thomson, 1861, monotypy [= *Ergates corticarius* Erichson, 1849].

1. *Strongylaspis corticarius* (Erichson, 1848)

Syntypes locality - Syntypes male and female: Guiana. (ZMHB). **Distribution** - United States (Florida), Mexico (Veracruz), Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Guyana, French Guiana, Venezuela, Brazil (Roraima), Cuba, Jamaica, Dominica, Martinique, St. Lucia, Dominican Republic.

Ergates corticarius Erichson in Schomburgk, 1849: 571; White, 1853: 11.

Strongylaspis corticarius; Bates, 1884: 234; Leng & Mutchler, 1914: 443 (distr.); Schwarz in Holland, 1917: 341 (distr.); Gowdey, 1926: 20 (distr.); Fisher, 1944: 3 (distr.); Franz, 1954: 215 (distr.); Zayas, 1957: 160, pl. 8; Linsley, 1962: 13; Zayas, 1975: 20, pl. 1, fig. b; Chemsak, Linsley & Mankins, 1980: 28 (distr.); Villiers, 1980a: 130; 1980b: 143, fig. 8; Wendt, 1984: 331 (syntypes); Chemsak, Linsley & Noguera, 1992: 16 (cat.); Maes *et al.*, 1994: 6 (distr.); Monné, M.A., & Giesbert, 1994: 7 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 70, pl. 4, figs 1, 2; Toledo *et al.*, 2002: 525 (distr.); Monné, M.L. & Santos-Silva, 2003: 45, figs 14, 18–28; Turnbow, Cave & Thomas, 2003: 4 (distr.); Lozada Piña, Fernández García & Trujillo Anaya, 2004: 105 (distr.); Monné, M.A., 2006: 61 (cat.); Peck, 2005: 165 (distr.); 2006: 191 (distr.); Hovore, 2006: 371 (distr.); Monné, M.A. & Hovore, 2006: 14 (checklist); Noguera *et al.*, 2007: 310 (distr.); Touroult, 2007: 6 (distr.); Audureau, 2008: 4 (distr.); Swift *et al.*, 2010: 8 (distr.); Maes *et al.*, 2010a: 71, 7 figs (distr.); Touroult, 2012: 76, fig. 10 (distr.); Lingafelter, 2015: 355, 1d, 13a. (distr.); Devesa, Fonseca & Barro, 2015: 29, 50, figs 1–10; Peck, 2016: 171 (distr.).

Strongylaspis (Strongylaspis) corticaria; Lameere, 1903a: 31; Rosales, 1966: 56, pl. 2, figs 3, 4; Vitali & Rezbanyai-Reser, 2003: 5, figs 1 a, b.

Strongylaspis corticaria; Galileo & Martins, in Rafael, 1991: 328 (distr.); Maes, 1998: 884 (distr.); Bouyer & Santos-Silva, 2016: 87, figs 1–4; Audureau & Roguet, 2018: 58 (distr.).

Strongylaspis scobinatus Thomson, 1861a: 313; Chevrolat, 1862: 272; Thomson, 1864: 299; Lacordaire, 1868: 101; Bates, 1872: 167; 1879: 6, pl. 1, fig. 7, pl. 2, fig. 9; 1884: 232; Thomson, 1878a: 5 (type); Gahan, 1895a: 82; Schaeffer, 1909: 149; Gowdey, 1926: 20 (distr.).

Type locality - Holotype female: Mexico. (MNHN).

Strongylaspis scobinatus var. *belti* Bates, 1884: 233.

Type locality - Holotype: Nicaragua, Chontales. (BMNH).

Strongylaspis lobulifer Bates, 1892b: 145.

Type locality - Holotype male: Mexico, Veracruz: Atoyac. (BMNH).

Hovatoma gramreta Gilmour, 1956: 30.

Hovatoma (Hovatoma) gramreta; Ferreira & Veiga Ferreira, 1959: 12; Gilmour, 1965: 3 (key).

Parahovatoma gramreta; Santos Ferreira, 1980: 63, 67.

Type locality - Holotype male: South Africa. (MAGD).

BASITOXINA Lameere, 1912

Basitoxi Lameere, 1912b: 180.

Basitoxina; Bousquet, Heffern, Bouchard & Nears., 2009: 17; Bouchard *et al.*, 2011: 457.

Type-genus: *Basitoxus* Audinet-Serville, 1832

Type-species: *Basitoxus armatus* Audinet-Serville, 1832 designated by Desmarest (1860: 307).

Mécosarthrines Lameere, 1903b: 307 (based on *Mecosarthron* Buquet, 1840). *Nomen nudum*.

Comment. This name is unavailable under Article 11.7.2 (vernacular name proposed after 1899)

Mecosarthrini Melzer, 1919: 35.

Type-genus: *Mecosarthron* Buquet, 1840

Type-species: *Mecosarthron buphagus* Buquet, 1840 by monotypy.

Archodontes Lameere, 1903

Basitoxus (Archodontes) Lameere, 1903b: 216.

Archodontes; Lameere, 1919: 27; Casey, 1924: 224; Bradley, 1930: 227; Linsley, 1962: 13;

Arnett, 1962: 855; Chemsak, 1996: 71; Monné, M.A., 2006: 46 (cat.); Monné, M.A. &

Hovore, 2006: 12 (checklist); Monné, M.A., 2012: 129.

Type-species - *Cerambyx melanopus* Linnaeus, 1767 (original designation).

Aplagiognathus; Thomson, 1864: 307 (*partim*).

Mallodon; Horn in Leng, 1884: 9 (*partim*).

Paramallus Casey, 1912: 222, 227.

Type-species - *Cerambyx melanopus* Linnaeus, 1767 (original designation).

1. *Archodontes melanopus melanopus* (Linnaeus, 1767)

Type locality - Type: United States, Carolina. (depository unknown). **Distribution** - Southern

United States from North Carolina to Louisiana. **Host plants** – *Acer negundo* Linnaeus

(Aceraceae), *Diospyros virginiana* Linnaeus (Ebenaceae), *Carya illinoensis* (Wagenheim)

K.Koch, *Quercus geminate* (Fagaceae), *Prunus persica* (Linnaeus) Batsch (Rosaceae).

Cerambyx melanopus Linnaeus, 1767: 623

Prionus melanopus; Fabricius, 1781: 208; 1787: 130; 1793: 250; Olivier, 1795: 18, pl. 112, fig.

46; Fabricius, 1801: 264; Schoenherr, 1817: 344; White, 1853:

Cerambyx (Prionus) melanopus; Gmelin, 1790: 1816

Mallodon melanopus; White, 1853: 46; Riley, 1880: 238; Packard, 1881: 28; Horn in Leng, 1884:

10; Riley, 1884: 410; Hubbard, 1886: 13; Packard, 1890: 50, pl. 35, fig. 1; Beutenmuller,

1896: 73 (hosts); Fall, 1901: 142; Knaus, 1903: 189; Hopkins, 1904: 37; Herrick, 1904: 21,

figs 13,14; Craighead, 1915: 15; Garnett, 1918: 172; Dozier, 1920: 339, fig. 8b

Aplagiognathus melanopus; Thomson, 1864: 307; 1867c: 90

Basitoxus (Archodontes) melanopus; Lameere, 1903b: 217

Paramallus melanopus; Casey, 1912: 227

Archodontes melanopus; Lameere, 1913: 10; Casey, 1924: 224; Doane *et al.*, 1936: 263; Brimley,

1938: 209; Loding, 1945: 112; Sherman, 1946: 126; Alexander, 1958: 45; Linsley, 1962: 15;

Baker, 1972: 206; Chemsak, Linsley & Noguera, 1992: 14; Brown & Peck, 1996: 2158;

Chemsak, 1996: 72; Korotyaev *et al.*, 2005: 253 (hosts); Vlasak & Vlasakova, 2021: 5

(hosts)

Stenodontes melanopus; Herrick, 1935: 219;

Archodontes (Mallodon) melanopus; Craighead, 1950: 252

Archodontes melanopus melanopus; Linsley, 1962: 16; Turnbow & Franklin, 1980: 338; Monné, M.A., & Giesbert, 1994: 5 (checklist); Monné, M.A., 1995: 5 (cat.); Chemsak, 1996: 73, pl. 4, figs 3, 4; Yanega, 1996: 26, pl. 1, fig. 5; Linsley & Chemsak, 1997: 345 (hosts); Schiefer, 1998: 115; Peck & Thomas, 1998: 116; Monné, M.A. & Hovore, 2006: 12 (checklist); Holt, 2013: 243; Haack, 2017: 113. Klingeman *et al.*, 2017: 294

Prionus cilipes Say, 1824: 328;

Orthosoma cilipes; Haldeman, 1847a: 31

Mallodon cilipes Haldeman, 1847b: 371; LeConte, 1851: 111; White, 1853: 46; LeConte, 1859a: 48

Aplagiognathus cilipes; Thomson, 1867c: 90

Paramallus cilipes; Casey, 1912: 228

Archodontes cilipes; Casey, 1924: 224*bi*-

Syntypes locality - Syntypes female: United States, Carolina. (depository unknown).

Mallodon simplicicollis Haldeman, 1847a: 30

Type locality - Syntypes female: United States. (MCZH)

Mallodon simplicicollis var. *bi-impressum* Haldeman, 1847a: 30

Type locality - Holotype male: United States (MCZH)

1a. *Archodontes melanopus aridus* Casey, 1924

Type locality - Holotype male: United States, Arizona "probably southern". (USNM).

Distribution - United States (Colorado River in California and Arizona), Mexico (Baja California). **Host plants** - *Salix* sp. (Salicaceae).

Archodontes aridus Casey, 1924: 224; Lingafelter *et al.*, 2014: 19, figs 19q, r (holotype).

Archodontes melanopus aridus; Linsley, 1962: 17, fig. 4; Chemsak, Linsley & Noguera, 1992: 14 (cat.); Monné, M.A., & Giesbert, 1994: 5 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 75, pl. 4, figs 7, 8; Linsley & Chemsak, 1997: 345 (hosts); Monné, M.A., 2002: 12 (cat.); Monné, M.A., 2006: 46 (cat.); Monné, M.A. & Hovore, 2006: 12 (checklist);

1b. *Archodontes melanopus serrulatus* (LeConte, 1854)

Syntypes locality - Syntypes: United States, Texas: Laredo to Ringgold Barracks. (MCZN).

Distribution - Southwestern United States from Arkansas to Arizona, northern Mexico. **Host plants** - *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae), *Citrus paradisi* Macfadyen (Rutaceae), *Populus* sp. (Salicaceae).

Mallodon serrulatus LeConte, 1854b: 82; 1858a: 40; 1859e: 127, pl. 3, fig. 5; Riley, 1880: 238 (hosts); Horn *in* Leng, 1884: 9, 10; Wickham, 1893: 143; Snow, 1906a: 146 (distr.); Schaeffer, 1908: 329 (distr.).

Aplagiognathus serrulatus; Thomson, 1867c: 91 (error).

Paramallus serrulatus; Casey, 1912: 228.

Archodontes melanopus serrulatus; Dean, 1953: 174; Linsley, Knull & Statham, 1961: 3; Linsley, 1962: 16, fig. 4; Manley & French, 1976: 46; MacKay, Zak & Hovore, 1987: 363; Hovore, Penrose & Neck, 1987: 293; Chemsak, Linsley & Noguera, 1992: 14 (cat.); Lingafelter & Horner, 1993: 163; Monné, M.A., & Giesbert, 1994: 5 (checklist); Chemsak, 1996: 74, pl. 4, figs 5, 6; Linsley & Chemsak, 1997: 345 (hosts); Monné, M.A., 2006: 46 (cat.); Monné, M.A. & Hovore, 2006: 12 (checklist);

Archodontes serrulatus; Alexander, 1958: 45

MALLODONINI Thomson, 1861

Mallodonitae Thomson, 1861a: 292 (key), 318; 1864: 305.

Mallodonites; Thomson, 1867c: 85.

Mallodontides; Lacordaire, 1868: 122.

Mallodontinae; Pascoe, 1869: 661, 671.

Mallodontini; LeConte, 1873b: 286; LeConte & Horn, 1883: 271; Leng, 1884: 8; Casey, 1912: 221; Villiers, 1980b: 140.

Mallodonini Drumont & Komiya, 2010: 91; Bouchard *et al.*, 2011: 459.

Type-genus: *Mallodon* Lacordaire, 1830

Type-species: *Cerambyx spinibarbis* Linnaeus, 1758 (monotypy).

Stenodontines Lameere, 1902c: 63, 104 (based on *Stenodontes* Audinet-Serville, 1832). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (vernacular name proposed after 1899).

Stenodontini Lameere, 1903a: 4 (key), 54.

Type-genus: *Stenodontes* Audinet-Serville, 1832.

Type-species: *Prionus exsertus* Olivier, 1795 designated by Desmarest (1860: 307).

***Mallodon* Lacordaire, 1830**

Mallodon Lacordaire, 1830: 171; Chevrolat in D'Orbigny, 1846a: 609; Desmarest, 1860: 307; Strauch, 1861: 124; Schjodte, 1865: 194; Thomson, 1867c: 92; Lacordaire, 1868: 125; Desmarest in Chenu, 1870: 307; Bates, 1879: 8; Leng, 1884: 9; Lameere, 1902c: 91; Casey, 1912: 221, 222; Lameere, 1913: 12; 1919: 31; Fragoso & Monné, M.A., 1995a: 217 (syn.); Chemsak, 1996: 78; Monné, M.A., 2006: 48 (cat.); Monné, M.A. & Hovore, 2006: 12 (checklist); Bousquet, 2007: 619; Monné, M.A., 2012: 129.

Type-species - *Cerambyx spinibarbis* Linnaeus, 1758 (monotypy).

Stenodontes (*Orthomallodon*) Linsley, 1957: 3; 1962: 19.

Type-species - *Cerambyx spinibarbis* Linnaeus, 1758 (original designation).

1. *Mallodon dasystemus dasystemus* (Say, 1824)

Type locality - Type: Lower Missouri River. (Depository unknown). **Distribution** - Southern United States, Mexico (Jalisco, Nayarit, Sinaloa, Tamaulipas, Veracruz), Honduras, Tobago, Colombia. **Host plants** - *Acer negundo* Linnaeus, *A. pseudoplatanus* Linnaeus (Aceraceae), *Alnus* sp. (Betulaceae), *Bursera simaruba* (Linnaeus) Sargent (Burseraceae), *Celtis laevigata* Willdenow (Cannabaceae), *Casuarina equisetifolia* Linnaeus (Casuarinaceae), *Quercus* sp. (Fagaceae), *Liquidambar styraciflua* Linnaeus (Hamamelidaceae), *Carya* sp. (Juglandaceae), *Inga* sp. (Mimosaceae), *Ficus* sp., *Morus* sp. (Moraceae), *Platanus occidentalis* Linnaeus, *P. wrightii* S. Watson (Platanaceae), *Salix* sp. (Salicaceae), *Bumelia lanuginosa* (Michaux) (Sapotaceae), *Ulmus crassifolia* Nuttall (Ulmaceae).

Prionus dasystemus Say, 1824: 326; 1835: 193; LeConte, 1859b: 184, 663.

Mallodon dasystemus; LeConte, 1851: 112; White, 1853: 46; LeConte, 1858a: 40 (distr.); Riley, 1880: 238 (hosts); Packard, 1881: 28 (biol.); Horn in Leng, 1884: 9, 10, pl. 2, fig. 5; Packard, 1890: 79, fig. 27 (biol.); Beutenmuller, 1896: 73 (hosts); Hebard, 1903: 261; Townsend, 1903: 75 (distr.); Hopkins, 1904: 37 (biol.); Tucker, 1906: 87 (distr.); Snow, 1906a: 147 (distr.); Schaeffer, 1908: 327; Wickham, 1909a: 40 (distr.); Mason, 1910: 24 (distr.); Casey, 1912: 223; Craighead, 1915: 14, pl. 3, fig. 2, pl. 5, fig. 2, pl. 7 (larva, pupa); Fragoso & Monné, M.A., 1995a: 219; Chemsak, 1996: 78, pl. 4, figs 11, 12; Monné, M.A., 2002: 13 (cat. hosts); Monné, M.A. & Hovore, 2006: 12 (checklist); Mankin *et al.*, 2018: 321 (biol.).

Aplagiognathus dasystemus; Thomson, 1861a: 320; 1867c: 91.

Stenodontes (*Mallodon*) *dasystemus dasystemus*; Lameere, 1902c: 77.

Stenodontes (*Mallodon*) *dasystemus*; Hatch & Ortenburger, 1930: 14 (distr.); Craighead, 1950: 252 (biol.); Gilmour, 1954: 7 (distr.); Franz, 1954: 216 (distr.); Duffy, 1960: 62 (larva, pupa).

Stenodontes dasystemus; Brimley, 1938: 209 (distr.); Loding, 1945: 112 (distr.); Vogt, 1949: 138 (distr., hosts); Dean, 1953: 174 (hosts); Dillon & Dillon, 1961: 576, pl. 57, No. 8; Solomon, Newsome & Darwin, 1972: 78.

Stenodontes (*Orthomallodon*) *dasytomus*; Linsley, 1957: 4; 1962: 19; Skiles, 1978a: 414; Chemsak, Linsley & Noguera, 1992: 15 (cat.).

Stenodontes (*Orthomallodon*) *dasytomus dasytomus*; Linsley, 1957: 4 (syn.); 1962: 20, fig. 5; Hovore, Penrose & Neck, 1987: 283, fig. 5 (biol., distr.); Lingafelter & Horner, 1993: 164 (distr.); MacRae, 1993: 226 (distr.); Monné, M.A., & Giesbert, 1994: 6 (checklist); Noguera & Chemsak, 1996: 396 (cat.).

Stenodontes (*Mallodon*) *dasytomus*; Duffy, 1960: 62 (larva).

- Stenodontes dasytomus*; Baker, 1972: 205 (biol.); Manley & French, 1976: 46, fig. 1; Browne & Peck, 1996: 2158, 2160; Toledo *et al.*, 2002: 525 (distr.); Peck, Cook & Hardy, 2002: 17 (distr.).
- Stenodontes dasytomus dasytomus*; Perry, 1974: 215 (distr.); Thomas, 1977: 83 (hosts); Turnbow & Franklin, 1980: 338 (distr.); Linsley & Chemsak, 1997: 434 (hosts).
- Mallodon dasystemus dasystemus*; Monné, M.A., 1995b: 9 (cat.); Turnbow, Cave & Thomas, 2003: 3 (distr.); Hovore & Santos-Silva, 2004: 51 (syn.); Monné, M.A., 2006: 48 (cat.); Santos-Silva & Martins, 2006: 397 (syn.); Hovore, 2006: 371 (distr.); Noguera *et al.*, 2007: 310 (distr.); Swift *et al.*, 2010: 8 (distr.); Maes *et al.*, 2010a: 47, 12 figs (distr.); Gutiérrez & Noguera, 2015: 136 (distr.); Audureau & Roguet, 2018: 58 (distr.); Vlasak & Vlasakova, 2021: 5 (hosts)
- Mallodon dasystemus*; García Morales *et al.*, 2015:112 (distr.).
- Mallodon costulata* LeConte, 1851: 111; White, 1853: 46.
- Aplagiognathus costulatus*; Thomson, 1867c: 91.
- Syntypes locality** - Syntypes female: United States, Georgia. (MCZN).
- Mallodon degeneratum* Thomson, 1867c: 95.
- Syntypes locality** - Syntypes male: America boreali. (MNHN).
- Mallodon masticator* Thomson, 1867c: 99; 1878a: 5 (type).
- Stenodontes (Mallodon) dasystemus masticator*; Lameere, 1902c: 78.
- Stenodontes (Mallodon) masticator*; Linsley, 1934b: 161; 1942: 85.
- Stenodontes (Stenodontes) masticator*; Linsley, Knull & Statham, 1961: 3, fig. 2.
- Stenodontes (Orthomallodon) dasytomus masticator*; Linsley, 1962: 21; Chemsak, Linsley & Noguera, 1992: 15 (cat.); Maes *et al.*, 1994: 5 (distr.); Noguera & Chemsak, 1996: 396 (cat.); Maes, 1998: 883 (distr.).
- Stenodontes dasytomus masticator*; Linsley & Chemsak, 1997: 435 (hosts).
- Mallodon dasystemus masticator*; Martínez, 2000: 84 (distr.).
- Syntypes locality** - Syntypes male: Colombia. (MNHN).
- Mallodon angustatum* Thomson, 1867c: 100; 1878: 5 (type); Bates, 1872: 168; 1879: 9 (distr.); 1884: 236; Dugés, 1884: 13, pl. 2 (larva, pupa).
- Syntypes locality** - Syntypes male: Mexico. (MNHN).
- Aplagiognathus guatemalensis* Casey, 1912: 227; Lingafelter *et al.*, 2014: 73, figs 78m, n (holotype).
- Type locality** - Guatemala, Escuintla. (USNM).
- Mallodon debile* Casey, 1912: 222; Lingafelter *et al.*, 2014: 50, figs 53c, d (holotype).
- Stenodontes debile*; Loding, 1945: 113 (distr.).
- Type locality** - Holotype female: United States, Alabama. (USNM).
- Mallodon baroni* Casey, 1912: 223; Lingafelter *et al.*, 2014: 25, figs 25u, v (holotype).
- Stenodontes (Orthomallodon) baroni*; Chemsak, Linsley & Noguera, 1992: 15 (cat.); Noguera & Chemsak, 1996: 396 (cat.).
- Nothopleurus baroni*; Fragoso & Monné, M.A., 1995a: 219; Monné, M.A., 2006: 54 (cat.).
- Type locality** - Holotype female: Mexico, Guerrero. (USNM). Distribution - Mexico (Guerrero).
- Stenodontes dasytomus socorrensis* Linsley & Chemsak, 1966c: 251; Chemsak & Linsley, 1978: 139 (distr., host).
- Stenodontes (Orthomallodon) dasytomus socorrensis*; Chemsak, Linsley & Noguera, 1992: 15 (cat.); Monné, M.A., & Giesbert, 1994: 6 (checklist); Noguera & Chemsak, 1996: 396 (cat.).
- Mallodon dasystemus socorroensis*; Monné, M.A., 1995b: 10 (cat.).
- Type locality** - Holotype male: Mexico, Islas Revilla Gigedo: Isla Socorro. (CASC).
- Stenodontes (Mallodon) molarius*; Linsley, 1942: 83 (not Bates, 1879).
- Mallodon melanopus*; Haldeman, 1847a: 30 (not Linnaeus, 1767).
- Mallodon spinibarbe*; Haldeman, 1847a: 31 (not Linnaeus, 1758).
- Mallodon mandibularis* Lackerbeck, 1998: 517, figs 1, 2.
- Type locality** - Holotype male: Costa Rica, Alajuela: Cariblanco. (KLPC).

***Neomallodon* Linsley, 1957**

Stenodontes Neomallodon) Linsley, 1957: 2; 1962: 18.

Neomallodon; Skiles, 1978a: 409; Chemsak, 1996: 83; Santos-Silva & Martins, 2006: 399; Monné, M.A. & Hovore, 2006: 13 (checklist); Santos-Silva & Wappes, 2013: 5.

Type-species - *Paramallus arizonicus* Casey, 1912: 228. (original designation).

1. *Neomallodon arizonicus* (Casey, 1912)

Type locality - Lectotype female: United States (USNM). **Distribution** - United States (Arizona), Mexico (Sonora, Jalisco). **Host plants** - *Quercus* sp. (Fagaceae).

Paramallus arizonicus Casey, 1912: 228.

Stenodontes (Neomallodon) arizonicus; Linsley, 1957: 3 (syn.); Linsley, Knull & Statham, 1961: 3 (distr.); Linsley, 1962: 19.

Neomallodon arizonicus; Skiles, 1978a: 409, fig. 1; Chemsak, Linsley & Noguera, 1992: 14 (cat.); Monné, M.A., & Giesbert, 1994: 5; Fragoso & Monné, M.A., 1995: 219

(checklist); Chemsak, 1996: 83, pl. 5, figs 5, 6; Linsley & Chemsak, 1997: 434 (hosts); Monné, M.A. & Hovore, 2006: 13 (checklist); Lingafelter *et al.*, 2014: 20, 369, fig. 20a (lect.); Heffern & Santos-Silva, 2016: 571.

Aplagiognathus remotus Linsley, 1934c: 161.

Type locality - Holotype male: United States, Arizona: Mt. Washington, near Nogales, Santa Cruz County. (CASC).

Nothopleurus Lacordaire, 1868

Nothopleurus Lacordaire, 1868: 125; Bates, 1879: 8; Lameere, 1912b: 129; Casey, 1912: 222, 224; Linsley, 1934b: 162; Quentin & Villiers, 1975: 29; Villiers, 1980b: 140; Fragoso & Monné, M. A., 1995a: 219; Chemsak, 1996: 79; Monné, M.A., 2006: 54 (cat.); Monné, M.A. & Hovore, 2006: 13 (checklist); Santos-Silva, Swift & Nearn, 2010: 8; Monné, M.A., 2012: 129.

Stenodontes (Nothopleurus); Lameere, 1902d c: 94; 1919: 32; Gilmour, 1956: 13.

Type-species - *Nothopleurus ebeninus* Lacordaire, 1868 (monotypy).

Mallodon Audinet-Serville, 1832: 176; Thomson, 1864: 307 (not Lepeletier & Audinet-Serville in Lacordaire, 1830).

Type-species - *Prionus maxillosus* Fabricius, 1775 (Thomson designation, 1864: 308).

1. *Nothopleurus lobigenis* Bates, 1884

Type locality - Holotype male: Mexico, Bahia de Tehuantepec. (MNHN). **Distribution** - United States (Texas to southern California), Mexico (Baja California, Durango, Jalisco, Sonora, Oaxaca, Nayarit, Chiapas, Puebla, Morelos, Sinaloa, Guerrero). **Host plants** - *Schinus molle* Linnaeus (Anacardiaceae), *Baccharis sergiloides* Gray, *B.sarothroides* A. (Asteraceae), *Bursera* sp. (Burseraceae), *Parkinsonia torreyana* S. Watson (Caesalpinaceae), *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae), *Ficus* sp. (Moraceae).

Nothopleurus lobigenis Bates, 1884: 235; Fragoso & Monné, M.A., 1995a: 219; Chemsak, 1996: 81, pl. 5, figs 3, 4; Monné, M.A., 2002: 16 (cat. hosts); Noguera *et al.*, 2002: 623 (distr.); Monné, M.A., 2006: 55 (cat.); Monné, M.A. & Hovore, 2006: 13 (checklist); Noguera *et al.*, 2007: 310 (distr.); Noguera *et al.*, 2009: 87 (distr.); Santos-Silva, Swift & Nearn, 2010: 12, figs 3, 4, 23, 24, 38, 39, 58, 67, 71, 76; Noguera *et al.*, 2012: 620 (distr.); Amith & Lingafelter, 2017: 308 (distr.); Noguera *et al.*, 2017: 3 (distr.); Vlasak & Vlasakova, 2021: 5 (hosts)

Stenodontes (Nothopleurus) lobigenis; Lameere, 1902c: 101; Linsley, 1934b: 59 (distr.); 1934c: 162; 1935a: 69 (distr.); Linsley & Ross, 1940: 76 (biol.); Linsley, 1942: 24; Gilmour, 1954: 7, pl. 6, fig. 9.

Stenodontes lobigenis; Brimley, 1938: 209 (distr.); Chemsak & Noguera, 1995: 57 (distr.); Linsley & Chemsak, 1997: 435 (hosts).

Stenodontes (Mallodon) lobigenis; Linsley, 1962: 23, fig. 6; Skiles, 1978a: 414; Monné, M.A., & Giesbert, 1994: 6 (checklist); Hovore, 1988: 2 (hosts, distr.); Chemsak, Linsley & Noguera, 1992: 15 (cat.); Noguera & Chemsak, 1996: 396 (cat.).

Stenodontes lobigenus; Turnbow & Franklin, 1980: 347 (distr.).

Mallodon gnatho LeConte, 1858b: 81 (not *Mallodon gnatho* White, 1853); 1858a: 40; LeConte & Horn, 1883: 272.

Type locality - Holotype male: Sonora. (MCZN).

Mallodon mandibulare Gemminger, 1872: 254 (not *Prionus mandibularis* Fabricius, 1801); Bates 1879: 10; 1884: 234; Horn *in* Leng, 1884: 9; Horn, 1894: 337 (distr.); Garnett, 1918: 172 (distr.).

Nothopleurus komiyai Santos-Silva & Martins, 2006: 194, figs 7-9; Monné, M.A. *et al.*, 2017a: 85 (holotype).

Type locality - Holotype female: Brazil, Bahia: Santo Antonio. (MZSP).

2. *Nothopleurus madericus* (Skiles, 1978)

Type locality - Holotype male: United States, Arizona: Santa Cruz Co., Madera Canyon (Roundup Picnic Ground, 5400 ft). (USNM). **Distribution** - United States (Arizona), Mexico (Chihuahua). **Host plants** - *Quercus hypoleuca* Engelmann (Fagaceae).

Stenodontes (Mallodon) madericus Skiles, 1978a: 414, figs 2-6; Chemsak, Linsley & Noguera, 1992: 15 (cat.); Monné, M.A., & Giesbert, 1994: 6 (checklist); Lingafelter *et al.*, 2014: 94, figs 104c, d (holotype).

Nothopleurus madericus; Fragoso & Monné, M.A., 1995a: 225, fig. 7B; Chemsak, 1996: 80, pl. 5, figs 1, 2; Monné, M.A., 2002: 16 (cat. hosts); Monné, M.A., 2006: 55 (cat.); Monné, M.A. & Hovore, 2006: 13 (checklist); Santos-Silva, Swift & Nearn, 2010: 19, figs 5, 6, 26, 27, 41, 42, 60, 76.

Stenodontes madericus; Linsley & Chemsak, 1997: 435 (hosts).

Stenodontes arizonicus; Hovore & Giesbert, 1976: 350 (not Casey, 1912); Hovore, Penrose & Giesbert, 1978: 99 (hosts).

Stenodontes Audinet-Serville, 1832

Stenodontes Audinet-Serville, 1832: 173; Laporte, 1840: 403; Blanchard, C.E., 1845: 142; Drapiez, 1845: 123; Chevrolat *in* D'Orbigny, 1849: 10; Gistel & Bromme, 1850: 425; Jacquelin DuVal *in* Sagra, 1857: 257; Strauch, 1861: 124; Thomson, 1861a: 323; 1864: 308; 1865: 480; 1867b: 105; Lacordaire, 1868: 123; Desmarest *in* Chenu, 1870: 307; Leng, 1884: 9; Lameere, 1902d: 67; Casey, 1912: 222, 224; Melzer, 1919: 43; Lameere, 1919: 30; Bradley, 1930: 227; Gilmour, 1956: 10; Zayas, 1957: 160; Dillon & Dillon, 1961: 576; Linsley, 1962: 17; Arnett, 1962: 855; Zayas, 1975: 21; Fragoso & Monné, M.A., 1995a: 219; Chemsak, 1996: 76; Monné, M.A., 2006: 58 (cat.); Monné, M.A. & Hovore, 2006: 14 (checklist); Monné, M.A., 2012: 130.

Stenodontes (Stenodontes); Laporte, 1840: 403; Lameere, 1902d: 91; 1919: 33; Linsley, 1962: 17.

Type-species - *Prionus exsertus* Olivier, 1795 (subsequent designation, Desmarest *in* Chenu, 1860: 307).

1. *Stenodontes chevrolati* Gahan, 1890

Syntypes locality - Syntypes male and female: Cuba. (BMNH). **Distribution** - United States (Florida), Cuba, Bahamas, Cayman Islands. **Host plants** - *Spondias purpurea* Linnaeus (Anacardiaceae), *Cordia gerascanthus* Linnaeus (Boraginaceae), *Bursera simaruba* (Linnaeus) Sargent (Burseraceae), *Cecropia peltata* Linnaeus (Cecropiaceae), *Tectona grandis* Linnaeus fil. (Lamiaceae), *Ceiba pentandra* (Linnaeus) Gaertner (Malvaceae), *Samanea saman* (Jacquin) Merrill (Mimosaceae), *Pinus caribaea* Morelet (Pinaceae).

Stenodontes chevrolati Gahan, 1890: 23; 1895a: 82; Casey, 1912: 224; Leng & Mutchler, 1914: 443 (distr.); Schwarz *in* Holland, 1917: 340; Zayas, 1957: 160, pl. 10; Chemsak, 1969a: 185 (distr.); Zayas, 1975: 21, pl. 1, fig. c; Chemsak, Linsley & Noguera, 1992: 14 (cat.); Browne, Peck & Ivie, 1993: 40 (distr.); Chemsak, 1996: 77, pl. 4, figs 9, 10; Linsley & Chemsak, 1997: 434 (hosts); Monné, M.A., 2002: 17 (cat. hosts); Aragón, 2006: 54 (distr.); Monné, M.A., 2006: 59 (cat.); Monné, M.A. & Hovore, 2006: 14 (checklist); Thomas & Turnbow, 2007: 582 (distr.); Fernández García, Favila Castillo & López Iborra 2009: 322 (distr.);

- Thomas, Turnbow & Steiner, 2013: 20 (distr.); Devesa, Fonseca & Barro, 2015b: 29, 46, figs 1-10.
- Stenodontes (Stenodontes) chevrolati*; Lameere, 1902d: 92; Cazier & Lacey, 1952: 5; Linsley, 1957: 4 (syn.); Duffy, 1960: 60 (larva); Linsley, 1962: 18; Skiles, 1978a: 414; Monné, M.A., & Giesbert, 1994: 6 (checklist); Peck, 2005: 164 (distr.).
- Stenodontes damicornis*; Chevrolat, 1862: 273 (not Linnaeus, 1771); LeConte *in* Schwarz, 1878: 470.
- Stenodontes exsertus*; Horn, 1880b: xvii (not Olivier, 1795).
- Stenodontes mandibularis*; Leng, 1884: 9 (not pl. 2, fig. 10; not Fabricius, 1801).
- Stenodontes cubensis* Casey, 1912: 224 ; Leng & Mutchler, 1914: 443 (distr.); Peck, 2005: 165 (distr.); Lingafelter *et al.*, 2014: 48, figs 51m, n (lect. designation).
- Type locality** - Lectotype male: Cuba, Habana. (USNM).
- Stenodontes parallelus* Casey, 1912: 225; Lingafelter *et al.*, 2014: 295, figs 127q, r (holotype).
- Type locality** - Holotype male: Southern Florida ?. (USNM).

MEROSCELISINI Thomson, 1861

- Meroscelisidae Thomson, 1861a: 285, 299; 1864: 279.
- Meroscelisides; Lacordaire, 1868: 46.
- Meroscelisini; Galileo, 1987a: 148 (rev.); Monné, M. A., 1995b: 63 (cat.); Chemsak, 1996: 125; Monné, M. A., 2006: 71 (cat.); Bousquet, Heffern, Bouchard & Nearn., 2009: 19; Bouchard *et al.*, 2011: 459.
- Type-genus:** *Meroscelisus* Audinet-Serville, 1832
- Type-species:** *Meroscelisus violaceus* Audinet-Serville, 1832 (monotypy).
- Aegosomeitae Thomson, 1861a: 308 (*partim*).
- Orthosomeitae Thomson, 1864: 284 (*partim*).
- Tragosomeitae Thomson, 1864: 286.
- Tragosomides; Lacordaire, 1868: 163.
- Tragosomini; LeConte, 1873b: 289; LeConte & Horn, 1883: 274; Leng, 1884: 8; Linsley, 1961: 629; 1962: 53
- Tragosomiens; Lameere, 1912b: 59
- Tragosomae; Lameere, 1919: 157.
- Type-genus:** *Tragosoma* Audinet-Serville, 1832
- Type-species:** *Cerambyx deorsarius* Linnaeus, 1767 (monotypy).
- Tragosomites; Fairmaire *in* Jacquelin DuVal, 1864: 119, 191. (based on *Tragosoma* Audinet-Serville, 1832). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Fairmaire 1864).
- Clostérides Lacordaire, 1868: 149 (*partim*).
- Closterinae; Pascoe, 1869: 661, 676 (*partim*).
- Closteriens; Lameere, 1912b: 5 (*partim*).
- Closteri; Lameere, 1913: 81 (cat.); 1919: 145 (*partim*).
- Closterina; Gilmour, 1956: 220.
- Closterini; Quentin & Villiers, 1974: 250; Fragoso & Monné, M. A., 1982: 519.
- Type-genus:** *Closterus* Audinet-Serville, 1832.
- Type-species:** *Closterus flabellicornis* Audinet-Serville, 1832 (monotypy). Availability (under Article 11.7.2): *Closteri* Lacord[aire], 1869 (Lameere 1913: 81).
- Monodesmides Lacordaire, 1868: 140, 157 (based on *Monodesmus* Audinet-Serville, 1832). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Lacordaire 1868 [1869]).
- Monodesminae Gahan, 1890: 299.
- Monodesmiens; Lameere, 1912b: 80.
- Type-genus:** *Monodesmus* Audinet-Serville, 1832
- Type-species:** *Monodesmus callidioides* Audinet-Serville, 1832 (monotypy).
- Luluina Gilmour, 1956: 181 (key), 222.
- Type-genus:** *Lulua* Burgeon, 1931
- Type-species:** *Lulua squamosa* Burgeon, 1931 (monotypy).

***Tragosoma* Audinet-Serville, 1832**

Tragosoma Audinet-Serville, 1832: 159; Mulsant, 1839: 23; Laporte, 1840: 398; Drapiez, 1845: 479; Chevrolat *in* D'Orbigny, 1849: 633; Gistel, 1848b: 130; LeConte, 1851: 107; Thomson, C. G., 1859: 148; Desmarest, 1860: 306; Thomson, 1861a: 290; 1865: 472; Mulsant, 1863: 42, 348; Fairmaire *in* Jacquelin DuVal, 1864a: 120; Schjodte, 1865: 195; Lacordaire, 1868: 167; Desmarest *in* Chenu, 1870: 306; LeConte, 1873b: 289; Redtenbacher, 1874: 396; Provancher, 1877: 583; LeConte & Horn, 1883: 274; Leng, 1884: 59; Casey, 1890: 491; 1899: 98 (key spp.); Reitter, 1912: 4; Lameere, 1912b: 59; 1919: 158; Schaufuss, 1916: 825; Planet, 1924: 22; Casey, 1924: 225; Portevin, 1927: 8; Picard, 1929: 38; Chagnon, 1936: 239; Plavilstshikov, 1936: 99; Knull, 1946: 147; Heyrovsky, 1955: 72; Linsley, 1962: 54; Arnett, 1962: 855; Chagnon & Robert, 1962: 239; Hatch, 1971: 92; Galileo, 1987a: 165 (rev.); Chemsak, 1996: 126; Monné, M.A., 2006: 76 (cat.); Monné, M.A. & Hovore, 2006: 17 (checklist); Monné, M.A., 2012: 131; Laplante, 2017: 4 (rev.); Bousquet. Laplante, Hammond & Langor, 2017: 30 (key spp)

Type-species - *Cerambyx depsarius* Linnaeus, 1767 (monotypy).

1. *Tragosoma depsarium* (Linnaeus, 1767)

Type locality - Type: Suecia. **Distribution** - Coniferous forests of the Northern Hemisphere.

Host plants – *Picea abietis* Linnaeus, *P. excelsa* Link, *Pinus contorta* Douglas ex Loudon, *P. monophylla* Torrey ex Frémont, *P. peuce* Grisebach, *P. pumilio* Haenkel, *P. rigida* Miller, *P. sibirica* Mayr, *P. strobus* Linnaeus, *P. sylvestris* Linnaeus, *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae).

Cerambyx depsarius Linnaeus, 1767: 624;

Tragosoma depsarium; Hamilton, 1889: 145; 1892: 296; 1894: 395; 1895: 337; Leng & Hamilton, 1896: 164 (syn.); Casey, 1899: 98; Mutchler, 1912: 213; Lameere, 1912b: 61; Garnett, 1918: 173; Casey, 1924: 225; Essig, 1926: 450; Mank, 1934: 79; Knowlton & Thatcher, 1936: 278; Chagnon, 1936: 239, pl. 15, fig. 2; Moore, 1937: 87; Linsley, Knull & Statham, 1961; 7 (distr.); Linsley, 1962: 54, fig. 19; 1953: 166; Chemsak, 1965: 93; Tyson, 1967: 124, figs 1-3 (larva, pupa); Hatch, 1971: 84, pl. 9, fig. 3; Baker. 1972: 198; Gosling, 1973: 68; Kirk & Balsbaugh, 1975: 97; Hovore & Giesbert, 1976: 350 (hosts); Furniss & Carolin, 1977: 390 (hosts); Galileo, 1987: 109, figs; Chemsak, Linsley & Noguera, 1992: 22; Monné, M.A., 1995: 64 (cat.); Yanega, 1996: 27, pl. 1, fig. 7; Chemsak, 1996: 127, pl., 9, figs 6.7; Linsley & Chemsak, 1997: 446 (hosts); Heffern, 1998: 7; Vlasák & Vlasakova, 2002: 204; Monné, M.A. & Hovore, 2006: 17 (checklist); Laplante, 2017: 10, fig. 8; Rice, MacRae & Merickel, 2017: 669 (distr.)

2. *Tragosoma harrisii* LeConte, 1851

Syntypes locality - Syntypes: Newfoundland, Connecticut. (MCZN). **Distribution** - Canada, Mexico, USA: Northeastern USA north to Labrador, west to southern Northwest Territories and British Columbia, south along the Rocky Mountains to northern Mexico (Chihuahua), along the Cascade range and the Sierra Nevada to southern California. **Host plants** - *Pinus ponderosa* Douglas ex Lawson & P. Lawson (Pinaceae).

Tragosoma Harrisii LeConte, 1851: 107; White, 1853: 28; Fitch, 1858: 715; LeConte, 1861b: 354; Lacordaire, 1868: 168; LeConte, 1867: 371; Pettit, 1871: 105 (distr.); LeConte, 1873b: 289; Horn, 1873: 717; Provancher, 1877: 583; Schaupp, 1878: 20 (distr.); Riley, 1880: 238 (hosts); LeConte & Horn, 1883: 274; Wickham, 1897b: 83

Tragosoma harrisii; Harrington, 1881: 33 (hosts); Packard, 1881: 162 (hosts); Snow, 1883: 42 (distr.); Leng, 1884: 60, pl. 2, fig. 7; Hamilton, 1889: 144 (hosts); Packard, 1890: 704 (hosts); Casey, 1890: 491; Beutenmuller, 1896: 74 (hosts); Wickham, 1897c: 156 (distr.); 1897a: 167 (hosts). Chagnon, 1897c: 14; Harrington, 1899: 107; Casey, 1899: 98; Smith, 1900: 284 (distr.); Fall, 1901: 143 (distr.). Ouellet, 1902: 122 (distr.); Knaus, 1903: 176 (distr.); Chagnon, 1905: 36 (distr.); Felt, 1906: 659, fig. 188 (distr.); Fall & Cockerell, 1907: 191 (distr.); Smith, 1910: 323 (distr.); Fisher & Kirk, 1912: 309 (distr.); Craighead, 1915: 17, pl. 3, fig. 1, pl. 4, fig. 3 (larva); Chagnon, 1917: 230 (distr.); Craighead, 1923: 29; Beaulne,

- 1932: 197 (hosts); Ingles, 1933: 59 (biol.); Craighead, 1950: 269; Keen, 1952: 183 (hosts); Clark, 1956: 41; Monné, M.A., & Giesbert, 1994: 18 (checklist); Webster, 2016: 484; Laplante, 2017: 10, figs 5, 7 (rev.); Bousquet, Laplante, Hammond & Langor, 2017: 30, pl 2
- Tragosoma depsarium* var. *harrisii*; Hamilton, 1890: 44; Champlain, Kirk & Knull, 1925: 105 (hosts); 1925: 139 (distr.); Hatch, 1925: 579 (distr.); Kirk & Knull, 1926: 21 (distr.); Procter, 1927: 110 (distr.); Leonard, 1928: 433 (distr.); Knowlton & Thatcher, 1936: 278 (distr.); Doane *et al.*, 1936: 165; Procter, 1946: 176; Knull, 1946: 147, pl. 3, fig. 8; Hardy, 1948: 31 (distr.); Knowton & Wood, 1950: 10 (distr.).
- Tragosoma depsarium* race *harrisii*; Schaeffer & Leng, 1912: 213.
- Tragosoma sodalis* Casey, 1899: 98; Fall & Cockerell, 1907: 191 (distr.); Lingafelter *et al.*, 2014: 361 (holotype); Laplante, 2017: 10 (syn. lectotype).
- Tragosoma sodale*; Galileo, 1987: 183 (rev.).
- Syntypes locality** - Syntypes male and female: United States, Colorado. (USNM).
- Tragosoma parvicollis* Casey, 1899: 99; Fall & Cockerell, 1907: 191 (distr.); Lingafelter *et al.*, 2014: 361 (holotype); Laplante, 2017: 10 (syn.).
- Type locality** - Holotype: United States, Utah: Southwestern Utah. (USNM).
- Tragosoma repens* Casey, 1924: 226; Knowlton & Thatcher, 1936: 278 (distr.); Laplante, 2017: 10 (syn.).
- Type locality** - Holotype male: United States, Utah. (USNM).

3. *Tragosoma pilosicorne* Casey 1890

- Type locality** - Holotype female: United States, California: Mount Diablo. (USNM).
- Distribution** - Northern, coastal and traverse mountain ranges of California and southern Oregon. **Host plants** - *Pinus ponderosa* Douglas ex Lawson & P. Lawson, *P. sabiniana* Douglas ex D, Don (Pinaceae)
- Tragosoma pilosicornis* Casey, 1890: 492m pl. 4, figs 14, 14^a; Leng & Hamilton, 1896: 164; Casey, 1899: 99; Fall, 1901: 143 (distr.); Klages, 1901: 289; Casey, 1924: 226; Leech, 1957: 51 (distr.); Linsley, 1962: 58; Tyson, 1967: 123, figs 2, 4 (larva, nympe). Galileo, 1987: 177, figs; Chemsak, Linsley & Noguera, 1992: 22 (cat.); Monné, M.A., & Giesbert, 1994: 18 (checklist); Monné, M.A., 1995: 65 (cat.); Chemsak, 1996: 130, pl. 9, figs 10, 11; Monné, M.A. & Hovore, 2006: 17 (checklist); Wescott *et al.*, 2006: 10 (distr.); Lingafelter *et al.*, 2014: 299, fig. 132e (holotype); Laplante, 2017: 10, fig; 6 (distr.)
- Tragosoma depsarium* form *pilosicornis*; Garnett, 1918: 173

4. *Tragosoma soror* Laplante 2017

- Type locality** - Holotype male: United States: Idaho, Ada County. (CNCI); **Distribution** - From southern British Columbia southeastward in the Rocky Mountains to central western Wyoming and southward along the Cascade range to the Sierra Nevada and the San Bernardino Mountains in central California. **Host plants** - *Pinus coniota* Douglas ex Loudon (Pinaceae)
- Tragosoma soror* Laplante, 2017: 4, figs 1-3; Bousquet, Laplante, Hammond & Langor, 2017: 31, fig. 14; Vlasak & Vlasakova, 2021: 6 (hosts)

5. *Tragosoma spiculum* Casey 1890

- Type locality** - Holotype male: United States, New Mexico: Las Vegas. (USNM).
- Distribution** - United States, Southern Rocky Mountains (southern Colorado, New Mexico) and isolated mountains in southeastern Arizona. **Host plants** - *Pinus leiophylla chihuahuana* (Engelmann) A.E. Murray (Pinaceae)
- Tragosoma spiculum* Casey, 1890: 492, pl. 4, fig. 15; Leng & Hamilton, 1896: 164; Casey, 1899: 99; Fall & Cockerell, 1907: 191 (distr.); Casey, 1912: 231; 1924: 226; Lingafelter *et al.*, 2014: 322, fig. 158c (holotype); Laplante, 2017: 8, 11 (syn.)
- Tragosoma depsarium* race *spiculum*; Schaeffer & Leng, 1912: 213
- Tragosoma chiricahuae* Linsley, 1959: 127; Linsley, Knull & Statham, 1961: 7, fig. 3 (distr.); Linsley, 1962: 57; Ruetter, 1970: 22 (paratype); Lewis, 1979: 22 (distr.); Galileo, 1987: 180, figs; Chemsak, Linsley & Noguera, 1992: 22 (cat.); Monné, M.A., & Giesbert, 1994: 18

(checklist); Monné, M.A., 1995: 64 (cat.); Chemsak, 1996: 128, pl. 9, figs 8, 9; Linsley & Chemsak, 1997: 446 (hosts); Heffern 1998: 7 (distr.); Monné, M.A. & Hovore, 2006: 17 (checklist)

PRIONINI Latreille, 1802

Prionii Latreille, 1802: 212.

Type-genus: *Prionus* Geoffroy, 1762

Type-species: *Cerambyx coriarius* Linnaeus, 1758 (see ICZN 1994: 60).

Prioniens; Chevrolat in D'Orbigny, 1847: 468.

Prionitae Verae Thomson, 1861a: 327.

Prionites; Fairmaire in Jacquelin DuVal, 1864a: 116, 191.

Prionites vrais; Lacordaire, 1868: 59.

Prionini; LeConte, 1873b: 286, 288; LeConte & Horn, 1883: 273; Leng, 1884: 8; Lameere, 1912b: 182; 1919: 104; Melzer, 1919: 128; Bradley, 1930: 227; Heyrovsky, 1955: 70; Gilmour, 1956: 125; Linsley, 1962: 30; Monné, M.A., 1995b: 48 (cat.); Chemsak, 1996: 91; Monné, M.A., 2006: 77 (cat.); Bousquet *et al.*, 2009: 19; Bouchard *et al.*, 2011: 460.

Prioceria Rafinesque, 1815: 116 (based on *Prioceras* Rafinesque, 1815). *Nomen nudum*. Comment. This name is unavailable under Article 11.7.1.1 (not based on an available generic name).

Cyrtognathites Blanchard, C.E. 1845: 138, 164 (based on *Cyrtognathus* Faldermann, 1835). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Blanchard 1845).

Psalidognathites Blanchard, C.E. 1845: 138, 165 (based on *Psalidognathus* Gray, 1832). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Blanchard 1845).

Prionommitae Thomson, 1861a: 295 (key), 327

Type-genus: *Prionomma* White, 1853

Type-species: *Prionus orientalis* Olivier, 1795 (monotypy).

Cyrtognathitae Thomson, 1861a: 328

Type-genus: *Cyrtognathus* Faldermann, 1835

Type-species: *Prionus paradoxus* Falderman, 1833 (monotypy).

Psalidognathitae Thomson, 1861a: 331.

Type-genus: *Psalidognathus* Gray, 1832

Type-species: *Psalidognathus friendii* Gray, 1832 (monotypy).

Orthosomitae Thomson, 1864: 284.

Type-genus: *Orthosoma* Audinet-Serville, 1832

Type-species: *Prionus cylindricus* Fabricius, 1775 (monotypy).

Pithoclitae Thomson, 1864: 291.

Type-genus: *Pithocles* Thomson, 1864

Type-species: *Derobrachus procerus* Thomson, 1860 (original designation).

Derobrachitae Thomson, 1864: 291.

Type-genus: *Derobrachus* Audinet-Serville, 1832

Type-species: *Derobrachus brevicollis* Audinet-Serville, 1832 (monotypy).

Derobrachinae Pascoe, 1869: 661.

Derobrachini; LeConte, 1873b: 286, 287; LeConte & Horn, 1883: 272; Leng, 1884: 8.

Titanitae Thomson, 1864: 292.

Type-genus: *Titanus* Audinet-Serville, 1832

Type-species: *Cerambyx giganteus* Linnaeus, 1771 (monotypy).

Aulacoceritae Thomson, 1864: 292.

Type-genus: *Aulacocerus* White, 1853

Type-species: *Aulacocerus mundus* White, 1853 (monotypy).

Psalidocoptides Lacordaire, 1868: 37, 38 (based on *Psalidocoptus* White, 1856). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Lacordaire 1868 [1869]).

Polyarthrides Lacordaire, 1868: 38, 44 (based on *Polyarthron* Audinet-Serville, 1832). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Lacordaire 1868 [1869]).

Micropsalides Lacordaire, 1868: 42 (based on *Micropsalis* Burmeister, 1865). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Lacordaire 1868 [1869]).

Polyarthrini Gounelle, 1911: 326.

Type-genus: *Polyarthron* Audinet-Serville, 1832

Type-species: *Prionus pectinicornis* Fabricius, 1793 (monotypy).

Micropsalini Gounelle, 1911: 326.

Type-genus: *Micropsalis* Burmeister, 1861 (junior homonym of *Micropsalis* Meyer, 1859 [Crustacea])

Type-species: *Micropsalis heterogama* Burmeister, 1861 by monotypy. Comment. This name is permanently invalid because it is based on a preoccupied type genus (Article 39).

***Derobrachus* Audinet-Serville, 1832**

Derobrachus Audinet-Serville, 1832: 154; Laporte, 1840: 402; Blanchard, C. E., 1845: 141; LeConte, 1851: 109; Desmarest, 1860: 307; Strauch, 1861: 125; Thomson, 1861a: 294; 1864: 291; 1865: 474; Lacordaire, 1868: 73; Desmarest *in* Chenu, 1870: 307; LeConte, 1873b: 288; Bates, 1879: 4; LeConte & Horn, 1883: 273; Leng, 1884: 10; Lameere, 1911: 260; 1919: 122; Bradley, 1930: 227; Linsley, 1940: 253; 1962: 30; Arnett, 1962: 855; Chemsak & Linsley, 1977: 508; Chemsak, 1996: 91; Monné, M.A., 2006: 79 (cat.); Monné, M.A. & Hovore, 2006: 17 (checklist); Santos-Silva, 2007a: 3 (rev.); Monné, M.A., 2012: 131.

Derobrachus (*Derobrachus*); Lameere, 1911: 263; 1919: 123.

Type-species - *Derobrachus brevicollis* Audinet-Serville, 1832 (monotypy).

Pithocles Thomson, 1864: 291; Lacordaire, 1868: 72; Bates, 1879: 4.

Type-species - *Pithocles procerus* Thomson, 1864 (original designation).

Braderochus Lacordaire, 1868: 74 (*partim*).

1. *Derobrachus brevicollis* Audinet-Serville, 1832

Syntypes locality - Syntypes male and female: United States, Georgia. (BMNH).

Distribution - Southern United States from North Carolina and Florida to Texas, **Host plants** - *Quercus* sp. (Fagaceae)

Derobrachus brevicollis Audinet-Serville, 1832: 155; Laporte, 1840: 402; Chevrolat, 1844: 62; Haldeman, 1847a: 31; LeConte, 1851: 109; White, 1853: 12; Strauch, 1861: 125; Thomson, 1864: 291; Lacordaire, 1868: 73; Chenu, 1870: 307; LeConte, 1873: 287; LeConte & Horn, 1883: 273; Leng, 1884: 11; Schaeffer, 1902: 236; Wickham, 1909b: 402; Dozier, 1920: 366; Doane *et al.*, 1936: 163; Loding, 1945: 113; Sherman, 1946: 126; Fattig, 1947: 4; Linsley, 1962: 33; Swan & Papp, 1972: 442; Chemsak & Linsley, 1977: 514; Morris, 1987: 140; Chemsak, Linsley & Noguera, 1992: 19; Monné, M.A., & Giesbert, 1994: 13 (checklist); Monné, M.A. 1995: 46; Browne & Peck, 1996: 2158; Chemsak, 1996: 99, pl. 6, figs 7, 8; Linsley & Chemsak, 1997: 364; Peck & Thomas, 1998: 116; Monné, M.A. & Hovore, 2006: 17 (checklist); Santos-Silva, 2007: 8, figs; Holt, 2013: 243; Klingeman *et al.*, 2017: 294

Derobrachus (*Derobrachus*) *brevicollis*; Lameere, 1911: 263

2. *Derobrachus geminatus* LeConte, 1853

Type locality - Holotype female: United States, New Mexico: Albuquerque. (MCZN).

Distribution - United States (Texas to California), Mexico (Baja California, Durango, Chihuahua, Sonora, Sinaloa). **Host plants** - *Cercidium microphyllum* Rose & I.M. Johnston (Caesalpinaceae), *Olea* sp. (Oleaceae).

Derobrachus geminatus LeConte, 1853: 233; 1858a: 40 (distr.); 1859a: 16, pl. 2, fig. 12; 1873b: 287, 288; LeConte & Horn, 1883: 273; Snow, 1883: 42 (distr.); Leng, 1884: 11, pl. 2, fig. 11; Bates, 1884: 231; 1892: 145; Horn, 1894: 337 (distr.); Townsend, 1895: 46 (distr.); Griffith,

1900: 569 (distr.); Fall, 1901: 142; Schaeffer, 1902: 236; Fall & Cockerell, 1907: 191 (distr.); Garnett, 1918: 173 (distr.); Smyth, 1934: 116 (biol.); Linsley, 1934b: 59 (distr.); Doane *et al.*, 1936: 163; Vogt, 1949: 138 (biol.); Thomas, 1951: 35 (hosts); Linsley, 1962: 31, fig. 10; Chemsak & Linsley, 1977: 510; Furniss & Carolin, 1977: 288 (hosts); MacKay, Zak & Hovore, 1987: 363 (hosts); Hovore, Penrose & Neck, 1987: 294; Hovore, 1988: 3 (distr.); Chemsak, Linsley & Noguera, 1992: 19 (cat.); Lingafelter & Horner, 1993: 164 (distr.); Monné, M.A., & Giesbert, 1994: 13 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 95, figs 1, 3; Linsley & Chemsak, 1997: 364 (hosts); Monné, M.A., 2002: 21 (cat. hosts); Monné, M.A., 2006: 80 (cat.); Monné, M.A. & Hovore, 2006: 18 (checklist); Santos-Silva, 2007a: 51, figs (syn.); Noguera *et al.*, 2009: 87 (distr.); Santos-Silva & Galileo, 2011: 43 (distr.).

Braderochus geminatus; Lacordaire, 1868: 74.

Derobrachus (Derobrachus) geminatus geminatus; Lameere, 1911: 264.

Derobrachus (Derobrachus) geminatus; Duffy, 1960: 68 (hosts).

Derobrachus forreri Bates, 1884: 230; Schaeffer, 1902: 236; Grossbeck, 1912: 325 (distr.); Chemsak & Linsley, 1970: 405 (lect.); 1977: 511; Wendt, 1984: 333 (syntype); Chemsak, Linsley & Noguera, 1992: 19 (cat.); Noguera & Chemsak, 1996: 396 (cat.)

Derobrachus (Derobrachus) geminatus forreri; Lameere, 1911: 265.

Derobrachus geminatus var. *forreri*; Van Dyke, 1934: 58 (hosts).

Derobrachus geminatus forreri; Linsley, 1942: 25; 1962: 33, fig. 10; Terrón, 1992: 300 (distr.); Linsley & Chemsak, 1997: 364 (hosts).

Derobrachus geminatus; LeConte, 1861a: 335 (not LeConte, 1853).

Type locality - Lectotype male: Mexico, Sinaloa: Presidio de Mazatlán. (BMNH).

3. *Derobrachus hovorei* Santos-Silva, 2007

Type locality - Holotype male: United States, California: Palm Springs (Riverside Co.).

(EMEC). **Distribution** - United States (California, Nevada, Arizona, New Mexico, Texas), Mexico (Chihuahua, Sonora, Nuevo León, Durango, Tamaulipas, Coahuila). **Host plants** - *Cercidium microphyllum* Rose & I.M. Johnston (Caesalpiniaceae), *Quercus* sp. (Fagaceae), *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae), *Morus rubra* Linnaeus (Moraceae), *Citrus* sp. (Rutaceae), *Populus* sp. (Salicaceae), *Ulmus* sp. (Ulmaceae), *Vitis* sp. (Vitaceae).

Derobrachus hovorei Santos-Silva, 2007a: 19, figs (host); García Morales *et al.*, 2015:112 (distr.); Moura & Von Groll, 2017: 457 (paratype).

Derobrachus geminatus; Bates, 1884: 231 (distr.); Schaeffer, 1902: 236 (distr.); Heyne & Taschenberg, 1908: 237; Linsley, 1962: 31; Chemsak, 1996: 95, figs 1, 3; Linsley & Chemsak, 1997: 364 (hosts) (not LeConte, 1853)

Derobrachus (Derobrachus) geminatus; Lameere, 1911: 264 (part); 1913: 67 (cat.); 1919: 123 (part).

Derobrachus geminatus geminatus; Linsley, Knull & Statham, 1961: 5; Linsley, 1962: 32, figs 9, 10; Linsley & Chemsak, 1997: 364 (hosts); Kingsley, 1998: 52; Monné, M.A., 2002: 21 (hosts) (not LeConte, 1853)

4. *Derobrachus leechi* Chemsak & Linsley, 1977

Type locality - Holotype male: United States, California: Imperial Co., Holtville. (CASC).

Distribution - United States (California, Arizona and Nevada), Mexico (Baja California, Sonora). **Host plants** - *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae).

Derobrachus leechi Chemsak & Linsley, 1977: 512; Chemsak, Linsley & Noguera, 1992: 19 (cat.); Monné, M.A., & Giesbert, 1994: 13 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 97, pl. 6, figs 5, 6; Monné, M.A., 2002: 22 (cat. hosts); Monné, M.A., 2006: 81 (cat.); Monné, M.A. & Hovore, 2006: 18 (checklist); Santos-Silva, 2007a: 17, figs; Noguera *et al.*, 2009: 87 (distr.).

Orthosoma Audinet-Serville, 1832

Orthosoma Audinet-Serville, 1832: 155; Laporte, 1840: 401; Drapiez, 1842: 195; LeConte, 1851: 110; Thomson, 1861a: 289; 1864: 285; Lacordaire, 1868: 145; Chenu, 1870: 307; LeConte, 1873b: 288; Provancher, 1877: 582; LeConte & Horn, 1883: 273; Leng, 1884: 11; Wickham, 1897: 82; Casey, 1912: 230; Linsley, 1940: 253; Knull, 1946: 145; Dillon & Dillon, 1961: 577; Linsley, 1962: 35; Arnett, 1962: 855; Monné, M.A., 1995: 48; Chemsak, 1996: 100; Monné, M.A. & Hovore, 2006: 18 (checklist); Bousquet, Laplante, Hammond & Langor, 2017: 29;

Type species - *Prionus cylindricis* Fabricius, 1781 (*Cerambyx brunneus* Forster, 1771) (monotypy)

1. *Orthosoma brunneum* (Forster 1771)

Type locality - Type: America Septentrionali (LSUK). **Distribution.** Southeastern Canada, eastern Canada, eastern and Central United States. **Host plants** - *Ostrya virginiana* (Miller) K. Koch (Betulaceae). *Cercis canadensis* Linnaeus (Caesalpiniaceae). *Juniperus virginiana* Linnaeus (*Cupressaceae*), *Robinia pseudoacacia* Linnaeus (Fabaceae). *Juglans nigra* Linnaeus (Juglandaceae), *Pinus strobus* Linnaeus, *P. rigida* Mill., *P. virginiana* Mill., *Tsuga canadensis* (Linnaeus) Carrière (Pinaceae).

Cerambyx brunneus Forster 1771: 37;

Cerambyx (Lamia) brunneus Gmelin, 1790: 1828

Lamia brunnea; Olivier, 1797: 453

Orthosoma brunneum; LeConte, 1873b: 288; Provancher, 1877: 583; Popenoe, 1877: 33; Riley, 1880: 238; Packard, 1881: 30, fig. 72; LeConte & Horn, 1883: 273; Leng, 1884: 11, pl. 2, fig. 8; Saunders, 1884: 54, fig. 21; Horn, 1886: 138; Harrington, 1887: 30, fig. 14; 1890: 185; Packard, 1890: 82, fig. 238, pl. 20, fig. 1; Riley & Howard, 1892: 239; Gahan, 1895: 83; Hamilton, 1895: 337; Fyles, 1896: 24, figs 9, 10; Beutenmuller, 1896: 74; Wickham, 1897b: 83, fig. 12; Ehrmann, 1897: 170; Smith, 1900: 284; Ehrman, 1900: 620; Dury, 1902: 158; Ulke, 1903: 25; Hopkins, 1904: 37; Melander, 1904: 20; Morris, 1908: 446, fig. 15; Wickham, 1909a: 28; Smith, 1910: 323; Blatchley, 1910: 1011, fig. 423; Lameere, 1911: 262; Leng, 1911: 215; Casey, 1912: 230; Fisher & Kirk, 1912: 309; Frost, 1912: 306; Leng & Mutchler, 1914: 444; Craighead, 1915: 15, pl. III, fig. 4, pl. IV, figs 6 & 7, pl. V, fig. 1, pl. VIII, figs 2 & 3; Johnson, 1915: 314; Chagnon, 1817: 230; Dozier, 1918: 334; Nicolay, 1919: 63; Britton, 1920: 266; Dozier, 1920: 340; Kempers, 1923: 102, fig. 571; Casey, 1924: 226; Kirk & Knull, 1926: 21; Fletcher, 1926: 143; Leonard, 1928: 433; Ware, 1929: 368; Beaulne, 1932: 197; Barrett, 1932: 190; Wolcott & Montgomery, 1933: 154; Goldman, 1933: 96, pl. 5, figs 54-57; Moennich, 1934: 98; Sheppard, 1935: 97; Chagnon, 1936: 239, pl. 15, fig. 1; Doane *et al.*, 1936: 163, fig. 77; Brimley, 1938: 210; Savely, 1939: 339; Loding, 1945: 113; Knull, 1946: 145, pl. 2, fig. 1; Craighead, 1950: 258; Beal *et al.*, 1952: 133; Zayas, 1957: 157; Linsley, 1957: 4; Alexander, 1958: 46; Gibson & Carrillo, 1959: 117; Anderson, 1960: 279; Dillon & Dillon 1961: 577, pl. 42, fig. 5; Chagnon & Robert, 1962: 239, pl. 15, fig. 1; Linsley, 1962: 35; Paim & Beckel, 1963: 1149, figs 1-10; 1964: 99; 1964: 295; 1964: 327; Baker, 1972: 205; Solomon, [Newsome] & Darwin, 1972: 78; Swan & Papp, 1972: 442, fig. 930; Gosling, 1973: 66; Meeking, Seabrook & Paim, 1974: 257; White, Paim & Seabrook, 1974: 235; Zayas, 1975: 44; Kirk & Balsbaugh, 1975: 96; Fox, 1975: 237; Solomon, Doolittle & Spilman, 1976: 290; Laliberté, Chantal & LaRochelle, 1977: 96; Turnbow & Franklin, 1980: 338; White, 1985: 281, fig. 122; Rice, Turnbow & Hovore, 1985: 18; Arnett, 1985: 360; Gosling, 1986: 157; Chemsak, Linsley & Noguera, 1992: 19; MacRae, 1993: 226; Monné, M.A., & Giesbert, 1994: 14 (checklist); Monné, M.A., 1995: 48; Chemsak, 1996: 100, pl. 6, figs 8-10; Yanega, 1996: 26, figs 8a, 8b; Linsley & Chemsak, 1997: 413; Schiefer, 1998: 115; Peck & Thomas, 1998: 116; Vlasák & Vlasakova, 2002: 204; Monné, M.A. & Hovore, 2006: 18 (checklist); Webster, MacCorquale & Majka, 2009: 201; Holt, 2013: 243; Steury & MacRae, 2014: 9; Vlasak, 2014: 317; Webster, 2016: 484; Klingeman *et al.*, 2017: 294; Bousquet, Laplante, Hammond & Langor, 2017: 29, pl. 1; Heffern, Vlasák & Alten, 2018: 740; Vlasak & Vlasakova, 2021: 5 (hosts)

Derobrachus (Orthosoma) brunneum; Lameere, 1913: 65; Hatch, 192Drury, 1773: 83, pl. 37, fig 1

Orthosoma (Orthosoma) unicolor; Laporte, 1840: 402

Prionus unicolor; Harris, 1841: 80; 1852: 85

Orthosoma unicolor; LeConte, 1852: 177; Chenu, 1870: 307; Packard, 1872: 495

Type locality - Holotype: United States. New York (Depository unknown)

Cerambyx pennsylvanicus Degeer, 1775: 99, pl. 13, fig. 13

Prionus (Orthosoma) pennsylvanicus Drury & Westwood, 1837: 78, pl. 47, fig. 1

Prionus pennsylvanicus; Emmons, 1854: 115, pl. 34, fig. 10

Orthosoma pennsylvanicus; Fitch, 1858: 714; Thomson, 1864: 285] Lacordaire, 1868: 146

Type locality - Type: United States. Pennsylvania (NHRS)

Prionus cylindricus Fabricius. 1781: 207; 1787: 229; 1793: 247, Olivier, 1795: 23, pl. 1, fig. 6; Fabricius, 1801: 251; Harris, 1838: 88

Orthosoma cylindricum; Audinet-Serville, 1832: 156; Drapiez, 1842: 195; Haldeman, 1847a: 31; LeConte, 1851: 110; Bland, 1861: 93; Walsh, 1865: 90; Walsh & Riley, 1868: 19; 1868: 40; Redtenbacher, 1868: 201; Riley, 1870: 87, fig. 60; Pettit. 1871: 105; Riley, 1872: 139; Saunders, 1875: 29; Bethune, 1877: 23; 1877: 222; Harrington, 1881: 33

Syntypes locality - Syntypes: Amerique Boreale, (ZMUC)

Cerambyx (Prionus) cylindroides Gmelin, 1790: 1818

Type locality - Type: Amerique Boreale (NHRS)

Prionus sulcatus Olivier, 1795: 39, pl. 8, fig. 27

Orthosoma sulcatus Chevrolat, 1852: 650

Type locality - Holotype: Cayenne. (depository unknown)

Orthosoma ampliars Casey, 1912: 231; Lingafelter *et al.*, 2014: 15, fig. 14u (holotype)

Orthosoma brunneus ampliars; Casey, 1924: 226

Type locality - Holotype female: United States, Indiana. (USNM)

Orthosoma spadix Casey, 1912: 231; Lingafelter *et al.*, 2014: 322, fig. 157 o (holotype)

Type locality - Holotype female: United States, New York: Willets Point; (USNM)

***Prionus* Geoffroy, 1762**

Prionus Geoffroy, 1762: 198; Fabricius, 1775: 159; Latreille, 1802: 265; Lepeletier & Audinet-Serville *in* Latreille, 1825: 199; Stephens 1831: 226; Gray *in* Griffith & Pidgeon, 1832: 99; Audinet-Serville, 1832: 191; Mulsant, 1839: 20; Laporte, 1840: 395; Drapiez, 1844b: 171; Chevrolat *in* D'Orbigny, 1847b: 470; Blanchard, C.E. *in* Cuvier, 1849: 469; LeConte, 1851: 107; Thomson, C. G., 1859: 148; Desmarest, 1860: 305; Thomson, 1861a: 295; 1864:283; 1865: 470; Mulsant, 1863: 40, 346; Schjodte, 1865: 195, 202; Lacordaire, 1868: 60; Desmarest *in* Chenu, 1870: 305; LeConte, 1873b: 288; Girard, 1873: 716; LeConte & Horn, 1883: 273; Dimmock, 1884:159 (biol.); Leng, 1884: 57; Blatchley, 1910: 1011; Reitter, 1912: 3; Lameere, 1912a: 185 (rev.); Casey, 1912: 231, 232; Schaufuss *in* Calwer, 1916: 823; Lameere, 1919: 132; Planet, 1924: 18; Portevin, 1927: 8; Picard, 1929: 36; Bradley, 1930: 227; Plavilstshikov, 1936: 69; Knull, 1946: 145; Heyrovsky, 1955: 71; Duffy, 1960: 70 (larva, pupa); Dillon & Dillon, 1961: 577; Linsley, 1962: 37; Arnett, 1962: 855; Hatch, 1971: 92; Kerzhner, 1991: 123 (available name); Chemsak: 1996: 101; Monné, M.A., 2006: 84 (cat.); Monné, M.A. & Hovore, 2006: 18 (checklist); Monné, M. A., 2012: 132; Santos-Silva, Nearn & Swift, 2016: 11 .

Type-species - *Cerambyx coriarius* Linnaeus, 1758 (Latreille subsequent designation, Latreille, 1810) [see also Kerzhner, 1991: 123, designation under the plenary powers].

Neopolyarthron Semenov, 1899: 255

Prionus (Neopolyarthron); Casey, 1924: 222 (syn.); Linsley, 1962: 46; Chemsak, 1996: 110; Monné, M. A., 2006: 87 (cat.); Monné, M.A., 2012: 132.

Type-species - *Cerambyx imbricornis* Linnaeus, 1767 (subsequent designation, Linsley, 1962: 46).

Prionus (Riponus) Casey, 1912: 245.

Type-species - *Cerambyx imbricornis* Linnaeus, 1767 (subsequent designation, Linsley, 1962: 46).

Prionus (Antennalia) Casey, 1912: 249; 1924: 223; Linsley, 1962: 49; Chemsak, 1996: 113.

Type-species - *Prionus fissicornis* Haldeman, 1846 (subsequent designation, Linsley, 1962: 49).

Prionus (Prionus) Geoffroy, 1762

Prionus (Prionus); Casey, 1912: 232; 1924: 212; Linsley, 1962: 38; Chemsak, 1996: 102; Monné, M.A., 2006: 84 (cat.); Monné, M.A. & Hovore, 2006: 18 (checklist);

Prionus (Prionellus) Casey, 1924: 209.

Hypoprionus Santos-Silva, Nearn & Swift, 2016: 8 (replacement name for *Prionellus* Casey, 1924 not *Prionellus* Kieffer, 1895 (Diptera).

Type-species - *Cerambyx laticollis* Drury, 1773 (Santos-Silva, Nearn & Swift, 2016: 8 designation, 1962: 38).

1. *Prionus (Prionus) californicus* Motschulsky, 1845

Type locality - Holotype: Northern California. (ZMUM). **Distribution** - Pacific Coast of North America, from Alaska to Sonora, Baja California and Rocky Mountains to New Mexico. **Host plants** - *Rhus* sp. (Anacardiaceae), *Alnus incana* (Linnaeus) Moench (Betulaceae), *Arbutus menziesii* Pursh (Ericaceae), *Castanea* sp., *Quercus garryana* Douglas, *Q. lobata* Née (Fagaceae), *Juglans regia* Linnaeus (Juglandaceae), *Eucalyptus* sp. (Myrtaceae), *Abies grandis* (Lambert) Lindley, *Pinus* sp., *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae), *Amelanchier canadensis* Medikus, *Malus* sp., *Prunus* sp., *Pyrus* sp., *Rubus* sp. (Rosaceae), *Citrus* sp. (Rutaceae), *Populus deltoides* Bartram ex Marshall, *P. trichocarpa* Torrey & A. Gray, *Salix* sp. (Salicaceae), *Sequoia sempervirens* (D. Don) Endlicher (Taxodiaceae), *Humulus lupulus* Linnaeus (Urticaceae), *Vitis* sp. (Vitaceae).

Prionus californicus Motschulsky, 1845: 89, pl. 2, fig. 9; LeConte, 1852: 177; Mannerheim, 1852: 364; White, 1853: 16; Snow, 1877: 19 (distr.); 1878: 76 (distr.); 1883: 42 (distr.); Leng, 1884: 57, 58; Rivers, 1886a: 64 (biol.); Blanchard, F., 1887: 86; bb: 144 (distr.); Townsend, 1892: 38; Blaisdell, 1892: 34 (hosts); Hamilton, 1894: 30 (distr.); Townsend, 1895: 46; Beutenmuller, 1896: 74 (hosts); Wickham, 1897a: 167 (hosts); Harrington, 1899: 107 (distr.); Daggett, 1901: 319 (biol.); Fall, 1901: 142 (distr.); Snow, 1906b: 170 (distr.); Fall & Cockerell, 1907: 191 (distr.); Schaeffer, 1908: 329 (distr.); Lameere, 1912a: 242; Craighead, 1915: 20, pl. 4, figs 8, 9 (larva); Essig, 1915: 251, fig. 241 (biol.); Garnett, 1918: 173 (distr.); Craighead, 1923: 29, pl. 36, fig. 2 (pupa); Essig, 1926: 449, figs 349, 350 (biol.); Tanner, 1927: 33, figs 167, 168; Hardy & Preece, 1927a: 187 (biol.); Crawford & Eyer, 1928: 3, 5 figs (biol., control); Tanner, 1928: 277 (distr.); Knowlton, 1930: 56; Pack, 1930: 219 (distr.); Beaulne, 1932: 289 (hosts); Barrett, 1932: 289 (hosts); Herrick, 1935: 274 (biol.); Knowlton & Thatcher, 1936: 278; Doane *et al.*, 1936: 164, figs 78, 79 (biol.); Moore, 1937: 87 (distr.); Linsley, 1938: 105 (syn.); Quayle, 1938: 319 (biol.); Linsley, 1942: 26; Crawford & Eyer, 1942: 1 (biol.); Hardy, 1942: 10 (biol.); Leech, 1947: 141 (biol.); Schuh & Mote, 1948: 103 (biol.); Knowlton & Wood, 1950: 10 (distr.); Craighead, 1950: 262 (biol.); DeLeon, 1952: 80 (hosts); Keen, 1952: 193 (hosts); Leech, 1955: 52 (biol.); Nishio, 1956a: 241, figs 1, 2; Linsley, 1957: 6 (syn.); Linsley, Knull & Statham, 1961: 5 (distr.); Tyson, 1970: 298 (distr.); Swan & Papp, 1972: 441, fig. 927; Tanner & Tanner, 1974: 220 (distr.); Kirk & Balsbaugh, 1975: 96 (distr.); MacKay, Zak & Hovore, 1987: 363 (distr., hosts); Hovore, 1988: 3 (distr.); Chemsak, Linsley & Noguera, 1992: 20 (cat.); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 106, pl. 7, figs 3, 4; Linsley & Chemsak, 1997: 424 (hosts); Heffern 1998: 6 (distr.); Monné, M.A., 2002: 23 (cat. hosts); Cervantes, Hanks, Lacey & Barbour, 2006: 718 (pheromone); Barbour *et al.*, 2006: 623 (biol.); Monné, M.A. & Hovore, 2006: 19 (checklist) Maki *et al.*, 2011: 933 (biol.); Barbour, Lacey & Hanks, 2007: 333 (biol.); Rice, Merickel & MacRae, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 30, pl 1; Barbour *et al.*, 2019: 1130 (pherom)

Prionus (Prionus) californicus; Casey, 1912: 242; 1924: 216; Linsley, 1962: 41, figs 13, 14; Hatch, 1971: 93, pl. 9, fig. 2; Monné, M.A., & Giesbert, 1994: 14 (checklist); Monné, M.A., 2006: 84 (cat.); Monné, M.A. & Hovore, 2006: 18 (checklist); Santos-Silva, Nearn & Swift, 2016: 80, figs 111-113, 158-171, 173-176, 183 (syn.).

Prionus crassicornis LeConte, 1851: 108; White, 1853: 17.

Prionus (Prionus) californicus crassicornis; Casey, 1924: 216.

Syntypes locality - Syntypes male and female: United States, Oregon. (MCZN).

Prionus curvatus LeConte, 1859a: 19.

Prionus (Prionus) californicus curvatus; Casey, 1924: 217.

Type locality - Holotype female: United States, New Mexico: near Santa Fe. (MCZN).

Prionus Horni Lameere. 1912a: 243; Linsley, 1935b: 161; Damoiseau & Cools, 1987: 33 (syntypes).

Syntypes locality - Syntypes male and female: United States, Arizona. (NHMW).

Prionus (Prionus) alutaceus Casey, 1912: 237; 1924: 216; Lingafelter *et al.*, 2014: 14, figs 13k, l (holotype).

Type locality - Holotype male: United States, Arizona. (USNM).

Prionus (Prionus) solidus Casey, 1912: 237; 1924: 215; Lingafelter *et al.*, 2014: 221, figs 157i, j (holotype).

Type locality - Holotype female: United States, Colorado. (USNM).

Prionus (Prionus) spiculosus Casey, 1912: 240; 1924: 217; Lingafelter *et al.*, 2014: 322, figs 157w, x (holotype).

Type locality - Holotype female: United States, Arizona: Grand Cañon of the Colorado. (USNM).

Prionus (Prionus) consors Casey, 1912: 240; 1924: 214; Lingafelter *et al.*, 2014: 44, figs 46q, r (holotype).

Type locality - Holotype male: United States, Colorado: Boulder Co. (USNM).

Prionus (Prionus) consors acomanus Casey, 1912: 241; Lingafelter *et al.*, 2014: 44, figs 46u, v (holotype).

Prionus (Prionus) acomanus; Casey, 1924: 218.

Type locality - Holotype female: United States, New Mexico: Jemez Springs. (USNM).

Prionus (Prionus) consors proximans Casey, 1912: 241; Lingafelter *et al.*, 2014: 45, figs 46w, x (holotype).

Prionus (Prionus) acomanus proximans; Casey, 1924: 218.

Type locality - Holotype: United States, New Mexico: Fort Wingate. (USNM).

Prionus (Prionus) angustulus Casey, 1912: 241; Lingafelter *et al.*, 2014: 17, figs 16s, t (lect. designation).

Prionus (Prionus) ineptus angustulus; Casey, 1924: 220.

Type locality - Lectotype male: United States, Cañon of the Colorado River. (USNM).

Prionus (Prionus) californicus ineptus Casey, 1912: 242; Lingafelter *et al.*, 2014: 35, figs 36m, n (holotype).

Prionus (Prionus) ineptus; Casey, 1924: 219.

Type locality - Holotype male: United States, California. (USNM).

Prionus (Prionus) californicus punctulatus Casey, 1912: 243; Lingafelter *et al.*, 2014: 35, figs 36q, r (holotype).

Prionus (Prionus) punctulatus; Casey, 1924: 220.

Type locality - Holotype female: United States, Arizona. (USNM).

Prionus (Prionus) fissifrons Casey, 1912: 243; Lingafelter *et al.*, 2014: 63, figs 67k, l (holotype).

Prionus (Prionus) spiculosus fissifrons; Casey, 1924: 218.

Type locality - Holotype male: United States, Arizona. (USNM).

Prionus (Prionus) texanus Casey, 1912: 243; 1924: 214; Lingafelter *et al.*, 2014: 332, figs 170a, b (holotype).

Type locality - Holotype male: United States, Texas: near El Paso. (USNM).

Prionus (Prionus) terminalis Casey, 1912: 244; Lingafelter *et al.*, 2014: 331, figs 168q, r (holotype).

Type locality - Holotype male: United States, Arizona: Cañon of the Colorado. (USNM).

Prionus (Prionus) serriger Casey, 1924: 215; Lingafelter *et al.*, 2014: 318, figs 153m, n (holotype).

Type locality - Holotype male: United States, Utah: Ibapah Mts., Trout Creek. (USNM).

Prionus (Prionus) suspectus Casey, 1924: 215; Lingafelter *et al.*, 2014: 330, figs 166s, t (holotype).

- Type locality** - Holotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) orbiceps Casey, 1924: 216; Lingafelter *et al.*, 2014: 292, figs 123w, x (holotype).
- Type locality** - Holotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) humeralis Casey, 1924: 216; Lingafelter *et al.*, 2014: 77, figs 84k, l (holotype).
- Type locality** - Holotype male: United States, California. (USNM).
Prionus (Prionus) californicus compar Casey, 1924: 217; Lingafelter *et al.*, 2014: 35, figs 36k, l (lect. designation).
- Type locality** - Lectotype male: United States, Utah: Provo. (USNM).
Prionus (Prionus) californicus ovipennis Casey, 1924: 217; Lingafelter *et al.*, 2014: 35, figs 36o, p (holotype).
- Type locality** - Holotype male: United States, Utah: Provo Cañon, North Fork. (USNM).
Prionus (Prionus) spiculosus coloradensis Casey, 1924: 218; Lingafelter *et al.*, 2014: 322, figs 158a, b (holotype).
- Type locality** - Holotype male: United States, Colorado, Boulder Co. (USNM).
Prionus (Prionus) scutellaris Casey, 1924: 219; Lingafelter *et al.*, 2014: 316, figs 151a, b (holotype).
- Type locality** - Holotype female: United States, Utah: Provo Cañon. (USNM).
Prionus (Prionus) ineptus ambiguus Casey, 1924: 219; Lingafelter *et al.*, 2014: 80, figs 86w, x (holotype).
- Type locality** - Lectotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) ineptus uintanus Casey, 1924: 220; Lingafelter *et al.*, 2014: 80, figs 87a, b (lect. designation).
- Type locality** - Lectotype male: United States, Utah: Provo Cañon, North Fork. (USNM).
Prionus (Prionus) spaldingi Casey, 1924:221; Knowlton & Thatcher, 1936: 278 (distr.); Lingafelter *et al.*, 2014: 322, figs 157q, r (lect. designation).
- Type locality** - Lectotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) stultus Casey, 1924: 221; Lingafelter *et al.*, 2014: 326, figs 163a, b (lect. designation).
- Type locality** - Lectotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) stultus parvicollis Casey, 1924: 222; Lingafelter *et al.*, 2014: 327, figs 163c, d (holotype).
- Type locality** - Holotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) nanus Casey, 1924: 222; Lingafelter *et al.*, 2014: 103, figs 114i, j (holotype).
- Type locality** - Holotype male: United States, Utah: Eureka. (USNM).
Prionus (Prionus) validiceps Casey, 1912: 235; 1924: 214; Linsley, 1957: 8; Lingafelter *et al.*, 2015: 339, fig. 177w,x.
- Type locality** - Holotype male: unknown (USNM).
Prionus (Prionus) tumidus Casey, 1912: 235; 1924: 213; Lingafelter *et al.*, 2014: 337, fig. 175g (holotype).
- Type locality** - Holotype male: United States, New Mexico: Fort Wingate. (USNM).
Prionus (Prionus) fontinalis Casey, 1924: 236; 1924: 212; Lingafelter *et al.*, 2014: 65, fig. 69s (holotype).
- Type locality** - Holotype male: United States. New Mexico: Jemez Springs. (USNM).

2. *Prionus (Prionus) fissicornis* Haldeman, 1848

- Syntypes locality** - Syntypes: United States, Nebraska: near Platte river. (MCZN).
Distribution - United States (Great Plains east of the Rocky Mountains, from Montana to Minnesota, south to Texas). **Host plants** - *Zea mays* Linnaeus (Poaceae)
Prionus fissicornis Haldeman, 1847a: 374; 1848: 125; LeConte, 1851: 108; White, 1853: 17; LeConte, 1858a: 40; 1859: 19, pl. 1, figs 14; Lacordaire, 1868: 61; Snow, 1878: 67; Popenoe, 1878: 82; LeConte & Horn, 1883: 274; Leng, 1884: 57, pl. 2, figs 14, 15; Blanchard, 1887: 86; Evans, 1906: 99; Fall & Cockerell, 1907: 191; Wickham, 1909a: 28; Lameere, 1912a: 240; Swenk, 1922: 200; Doane *et al.*, 1936: 165; Brimley, 1938: 210; Alexander, 1958: 49; Kirk & Balsbaugh, 1975: 97; Stein & Tigestad, 1976: 31; Turnbow & Franklin, 1980: 347

Prionus (Antennalia) fissicornis; Casey, 1924: 223; Linsley, 1957: 9; 1962: 49; Hovore, Penrose & Neck, 1987: 394; Chemsak, Linsley & Noguera, 1992: 21; MacRae, 1993: 227; Lingafelter & Horner, 1995: 393; Monné, M.A., & Giesbert, 1994: 15 (checklist); Monné, M.A., 1995: 55; Chemsak, 1996: 113, pl. 8, 2 figs; Heffern, 1998: 6

Prionus (Prionus) fissicornis; Santos-Silva, Nearn & Swift, 2016: 36, figs

Prionus (Antennalia) fissicornis transversus Casey, 1912: 251; 1924: 223; Lingafelter *et al.*, 2014: 63, fig. 67i (holotype)

Type locality - Holotype female: United States, Texas, (USNM).

Prionus (Antennalia) thoracicus Casey, 1924: 223; Lingafelter *et al.*, 2014: 333, fig. 170w (holotype)

Type locality - Holotype female: United States, Colorado: Akron, (USNM)

3. *Prionus (Prionus) heroicus* Semenov, 1908

Type locality - Lectotype male: United States, Arizona. (MCZN). **Distribution** - Southwestern United States, Mexico (Tamaulipas). **Host plants** - *Quercus* sp. (Fagaceae).

Prionus heros Fall, 1905: 274 (preoccupied); Skinner, 1905: 291.

Prionus heroicus Semenov, 1908: 259 (new name); Lameere, 1912a: 241; Casey, 1912: 244; 1924: 212; Linsley, 1957: 8 (syn.); Linsley, Knull & Statham, 1961: 7 (distr.); Linsley, 1962: 45; Hovore & Giesbert, 1976: 350 (biol.); Lewis, 1979: 22 (distr.); Chemsak, Linsley & Noguera, 1992: 20 (distr.); Chemsak, 1996: 109, pl. 7, figs 7, 8; Linsley & Chemsak, 1997: 424 (hosts); García Morales *et al.*, 2015: 112 (distr.).

Prionus (Riponus) heroicus; Lameere, 1919: 138.

Prionus (Prionus) heroicus; Monné, M.A., & Giesbert, 1994: 135 (checklist); Monné, M.A. & Hovore, 2006: 19 (checklist); Santos-Silva, Nearn & Swift, 2016: 76, figs 143-157, 172, 177, 178 (lectotype).

Prionus (Prionus) vastus Casey, 1912: 236; 1924: 213; Lingafelter *et al.*, 2014: 342, fig. 180e (holotype).

Type locality - Holotype female: United States. Arizona. (USNM).

Prionus (Prionus) tristis Casey, 1912: 236; 1924: 212; Lingafelter *et al.* - United States.

Syntypes locality - Syntypes female: New Mexico: Jemez Springs. (USNM).

Prionus (Prionus) tetricus Casey, 1912: 237; Lingafelter *et al.*, 2014: 332, fig. 169g (holotype).

Type locality - Holotype female: United States. New Mexico: Jemez Springs. (USNM).

4. *Prionus (Prionus) imbricornis* (Linnaeus, 1767)

Type locality - Type: United States, Carolina. (Depository unknown). **Distribution** - Indiana to Louisiana west to Great Plains from South Dakota to Texas and Florida, Dominican Republic. **Host plants** - *Castanea dentata* (Marshall) Borkhausen, *Quercus* sp. (Fagaceae), *Zea mays* Linnaeus (Poaceae), *Pyrus* sp., *Sorbus* sp. (Rosaceae), *Vitis* sp. (Vitaceae).

Cerambyx imbricornis Linnaeus, 1767: 622.

Cerambyx (Prionus) imbricornis; Gmelin, 1790: 181.

Prionus imbricornis; Olivier, 1795: 28, pl. 13, fig. 52; Palisot de Beauvois, 1805: 242, pl. 36, fig. 2; Schoenherr, 1817: 340; Lepeletier & Audinet-Serville, 1825: 202; Audinet-Serville, 1832: 192; Haldeman, 1847a: 31; LeConte, 1851: 108; Chevrolat, 1852: 650; White, 1853: 17; LeConte, 1859a: 48 (distr.); Lacordaire, 1868: 61; Riley, 1870: 89 (biol.); 1872: 140; Popenoe, 1877: 33 (distr.); Snow, 1878: 67 (distr.); Webster, 1879: 20 (biol.); Riley, 1880: 238 (hosts); LeConte & Horn, 1883: 274; Leng, 1884: 57, pl. 2, fig. 13; Horn, 1885: 89 (syn.); 1886: 138; Hamilton, 1886: 112 (larva); Blanchard, 1887: 86; Marten, 1890: 60; Riley & Howard, 1891: 407 (biol.); Bruner, 1891: 240 (biol.); Forbes, 1894: 106, pl. 13, figs 3, 6; Hamilton, 1895: 337 (distr.); Leng & Hamilton, 1896: 164 (syn.); Beutenmuller, 1896: 74 (hosts); Lugger, 1899: 194, fig. 117; Ulke, 1903: 25 (distr.); Hebard, 1903: 261; Tucker, 1906: 12 (distr.); Wickham, 1909a: 28 (distr.); Blatchley, 1910: 1012, fig. 425; Leng, 1911: 215 (distr.); Lameere, 1912a: 239 (rev.); Fisher & Kirk, 1912: 309 (distr.); Craighead, 1915: 18, pl. 8, fig. 1 (larva); 1923: 29 (larva); Kirk & Knull, 1926: 21 (distr.); Leonard, 1928: 433 (distr.); Ware, 1929: 368 (distr.); Beaulne, 1932: 197 (hosts); Doane *et al.*, 1936: 165; Brimley, 1938: 210 (distr.); Loding, 1945: 113 (distr.); Knull, 1946: 146, pl. 2, fig. 3;

- Craighead, 1950: 261 (hosts); Beal, Haliburton & Knight, 1952: 71, pl. 10, fig. 4; Alexander, 1958: 49 (distr.); Dillon & Dillon, 1961: 580, pl. 57, No.7; Frost, 1969: 95 (distr.); Baker, 1972: 200, fig. 68; Swan & Papp, 1972: 442, fig. 929; Kirk & Balsbaugh, 1975: 96 (distr.); Payne, Lowman & Pate, 1975: 680 (biol.); Mullins, 1975: 43 (biol.); Stein & Tigestad, 1976: 31; Payne *et al.*, 1976: 9; Turnbow & Franklin, 1980: 338 (distr.); White, 1985: 281, fig. 122; Yanega, 1996: 27, pl. 2, fig. 12; Linsley & Chemsak, 1997: 424 (hosts); Lingafelter, 2007: 13, 138, 2 figs, pl. 2f (hosts); Steury & MacRae, 2014: 9 (distr.).
- Prionus (Riponus) imbricornis* Casey, 1912: 248.
- Prionus (Neopolyarthron) imbricornis*; Lameere, 1913: 77 (cat.); 1919: 137; Casey, 1924: 222; Ortenburger & Hatch, 1926: 146 (distr.); Hatch, 1930: 26; Gilmour, 1954: 45 (distr.); Linsley, 1957: 9 (syn.); 1962: 46, figs 15, 16; Arnett, 1985: 360, fig. 24.178; Hovore, Penrose & Neck, 1987: 294; Chemsak, Linsley & Noguera, 1992: 21 (cat.); Lingafelter & Horner, 1993: 164 (distr.); Monné, M.A., & Giesbert, 1994: 16 (checklist); Monné, M.A., 1995: 54 (cat.); Chemsak, 1996: 111, pl. 7, figs 9, 10; Heffern, 1998: 6 (distr.); Schiefer, 1998: 115 (distr.); Holt, 2013: 243 (distr.).
- Prionus (Prionus) imbricornis*; Santos-Silva, Nearn & Swift, 2016: 36, figs, (syn.).
- Prionus debilis* Casey, 1891: 21; Rice, 1981: 479 (distr.); Yanega, 1996: 26, pl. 1, figs 11a,b.
- Prionus (Riponus) debilis*; Casey, 1912: 249.
- Prionus (Neopolyarthron) debilis*; Casey, 1924: 222; Linsley, 1962: 47, fig. 16; Chemsak, Linsley & Noguera, 1992: 21 (cat.); MacRae, 1993: 227 (distr.); Monné, M.A., 1995: 54 (cat.); Chemsak, 1996: 112, pl. 7, figs 11, 12; Heffern, 1998: 174 (distr.); Peck & Thomas, 1998: 116 (distr.); Lingafelter *et al.*, 2014: 50, 369, fig. 53i (lectotype).
- Prionus imbricornis debilis*; Brimley, 1938: 210.
- Prionus imbricornis* var. *debilis*; Knoll, 1946: 146.
- Type locality** - Lectotype male: United States, Indiana. (USNM).
- Prionus (Riponus) imbricornis mimus* Casey, 1912: 248; Lingafelter *et al.*, 2014: 78, fig. 85 m (holotype).
- Prionus (Neopolyarthron) imbricornis mimus* Casey, 1924: 222.
- Type locality** - Holotype male: United States. (USNM).
- Prionus (Riponus) imbricornis brunneus* Casey, 1912: 248; Lingafelter *et al.*, 2014: 78, fig. 85 k (holotype).
- Prionus (Neopolyarthron) imbricornis brunneus*; Casey, 1924: 222.
- Type locality** - Holotype male: United States, North Carolina: Southern Pines. (USNM).
- Prionus (Riponus) diversus* Casey, 1912: 247; Lingafelter *et al.*, 2014: 50, 369, fig. 57o (lectotype).
- Prionus (Neopolyarthron) diversus* Casey, 1924: 222; Hatch, 1930: 26.
- Prionus (Neopolyarthron) imbricornis diversus*: Ortenburger & Hatch, 1926: 146 (distr.).
- Type locality** - Lectotype male: United States, Indiana. (USNM).
- Prionus (Riponus) diversus cuneatus* Casey, 1912: 247; Lingafelter *et al.*, 2014: 54, 369, fig. 57q (lectotype).
- Prionus (Neopolyarthron) cuneatus*; Casey, 1924: 222.
- Type locality** - Lectotype male: United States, North Carolina: Southern Pines. (USNM).
- Prionus (Antennalia) fissicornis parviceps* Casey, 1912: 250; Lingafelter *et al.*, 2014: 62, fig. 63g (lectotype).
- Prionus (Antennalia) parviceps*; Casey, 1924: 223.
- Type locality** - Holotype female: United States, Texas. (USNM).
- Prionus beauvoisi* Lameere, 1915: 60 (*nom. nov.*).
- Type locality** - Holotype female: Dominican Republic. (MNHN).
- Prionus (Neopolyarthron) robustus* Casey, 1924: 222; Lingafelter *et al.*, 2014: 311, fig. 174w (holotype).
- Prionus robustus*: Brimley, 1938: 210 (distr.).
- Type locality** - Holotype female: United States, North Carolina: Southern Pines. (USNM).

5. *Prionus (Prionus) laticollis* (Drury, 1773)

Type locality - Holotype: United States, New York, (depository unknown). **Distribution** - Southern Canada and eastern United States to Oklahoma and South Dakota, **Host plants** – *Castanea* sp., *Quercus* sp (Fagaceae), *Kalmia latifolia* (Lauraceae)... *Malus* sp (Rosaceae), *Populus* sp. (Salicaceae);

Cerambyx laticollis Drury, 1773: 83, pl. 37, fig. 2.

Prionus (Derobrachus) laticollis. Drury & Westwood, 1837: 78, pl. 37, fig. 2

Prionus laticollis; Harris, 1841: 79; Haldeman, 1847a: 31; LeConte, 1852: 177; Emmons, 1854: 115, pl. 34, fig. 5; Fitch, 1859: 845; Bland, 1861: 93; Holmes, 1869: 51; Packard, 1870: 594, fig. 120; Riley, 1870: 77, figs 61, 62; 1873: 56; Treat, 1874: 344, figs 1-5; Saunders, 1875: 29, fig. 5; Popenoe, 1877: 33; Riley, 1880: 238; Packard, 1881: 127; 1881: 118, figs 117, 118; Weed, 1884: 13; Leng, 1884: 57, pl. 2, figs 6, 12; Blanchard, 1887: 85; Packard, 1890: 82, figs 162, 163; Bruner, 1891: 185; Webster, 1892: 198, figs 21-23; Hopkins, 1893: 192; Bruner, 1894: 154; Hamilton, 1895: 337; Beutenmuller, 1896: 74; Ehrmann, 1897: 170; Wickman, 1897: 83, figs 13, 14; Bruner, 1899: 162, figs 37-39; Lugger, 1899: 193, fig. 115; Smith, 1900: 285, fig. 124; Hopkins, 1902: 60; Dury, 1902: 158; Ulke, 1903: 25; Lockhead, 1903: 111, fig. 97; Hopkins, 1904: 97; Horsfall, 1904: 7, fig. 3; Pettit, 1904: 41, fig. 40; Laurent, 1905: 62; Fyles, 1905: 92, fig. 13; Morris, 1908: 446; Smitth, 1910: 334, fig. 130; Blatchley, 1910: 1011, fig. 424; Lameere, 1912a: 236; Fisher & Kirk, 1912: 309; Craighead, 1915: 19, pl. 3, fig. 3, pl. 6, fig. 1; Johnson, 1915: 314; Nicolay, 1919: 63; Lameere, 1919: 137; Leng, 1920: 266; Britton, 1920: 266; Kempers, 1923: 102; Kirk & Knull, 1926: 21; Leonard, 1928: 433; Hatch, 1930: 26; Beaulne, 1932: 197; Goldman, 1933: 95, pl. 5, figs 45-48; Easterling, 1934: 140; Herrick, 1935: 220, figs 180, 274; Doane *et al.*, 1936: 165; Brimley, 1938: 210; Becker, 1942: 208; Loding, 1945: 113; Knull, 1946: 145; Sherman, 1946: 126; Craighead, 1950: 252, fig. 50c; Jaques, 1951: 251, fig. 593; Beal *et al.*, 1952: 71, pl. 1, fig. 2; Shenefelt & Benjamin, 1955: 99; Linsley, 1957: 8; Alexander, 1958: 49; Gibson & Carrillo, 1959: 117; Linsley, 1962: 39; Farrar & Kerr, 1968: 563; Benham, 1969: 1331, figs. 1-4; 1970: 1413, figs 1-5; 1971: 89; Payne, Tedders & Gentry, 1970: 3, fig. 3; Swan & Papp, 1972: 442, fig. 928; Baker, 1972: 200; Gosling, 1973: 67; Kirk & Balsbaugh, 1975: 96; Benham & Farrar, 1976: 569, figs 1-9; Turnbow & Franklin, 1980: 338; Chemsak, Linsley & Noguera, 1992: 21; MacRae, 1993: 227. Monné, M.A., 1995: 52; Chemsak, 1996: 103, pl. 6, fig 11, 12; Yanega, 1996: 17, pl. 1, figs 9a, 9b; Schiefer, 1998: 115; Peck & Thomas, 1998: 116; Vlasák & Vlasakova, 2002: 204; Holt, 2013: 243; Staury & MacRae, 2014: 9; Santos-Silva, Nearn & Swift, 2016: 37, figs; Klingeman *et al.*, 2017: 294; Haack, Keena & Eyre, 2017: 74; Bousquet, Laplante, Hamond & Langor, 2017: 30, pl. 1; Hanks & Wang, 2019: 32; Vlasak & Vlasakova, 2021: 5 (hosts)

Prionus (Prionus) laticollis. Casey, 1912: 234; Monné, M.A., & Giesbert, 1994: 15 (checklist); Monné, M.A. & Hovore, 2006: 19 (checklist);

Prionus (Prionellus) laticollis; Casey, 1924: 209; Gilmour, 1954: 45 *Prionus brevicornis* Fabricius, 1801: 260; Harris, 1838: 88; Haldeman, 1847a: 31; LeConte, 1851: 109; Chevrolat, 1853: 650; White, 1853: 16; Bethune, 1868: 23; Packard, 1872: 495; LeConte, 1878: 126; LeConte & Horn, 1883: 274; Clarkson, 1884: 95; Packard, 1890: 481; Zimsen, 1964: 163

Syntypes locality - Syntypes: America Boreali. (ZMCD)

Prionus (Prionus) kempfi Casey, 1912: 233; Lingafelter *et al.*, 2014: 84, fig 92c (holotype)

Type locality - Holotype female: United States. New York: Adirondacks mountains (USNM)

Prionus (Prionus) parvus Casey, 1912: 234; Lingafelter *et al.*, 2014: 296, fig 128w (holotype)

Type locality - Holotype male: United States. (USNM)

Prionus (Prionus) laticollis oblongus Casey, 1912: 234; Lingafelter *et al.*, 2014: 87, fig 95i (holotype)

Type locality - Holotype male: United States, Indiana, (USNM)

Prionus (Prionellus) frosti Casey, 1924: 210; Lingafelter *et al.*, 2014: 76, fig 71k (lectotype)

Type locality - Lectotype male: United States, Massachusetts: Framingham. (USNM)

Prionus (Prionellus) nigrescans Casey, 1924: 211; Lingafelter *et al.*, 2014: 106, fig 116s (lectotype)

Type locality - Lectotype male: United States, New York: Lake George (Adirondacks). (USNM)

Prionus (Prionellus) densus Casey, 1924: 211; Lingafelter *et al.*, 2014: 52, fig 55o (holotype)

Type locality - Holotype male: United States (USNM)

6. *Prionus (Prionus) lecontei* Lameere, 1912

Type locality - Lectotype male: California. (NHMW). **Distribution** - British Columbia to Mexico (Baja California). **Host plants** - *Quercus agrifolia* Née, *Q. dumosa* Nuttall, *Q. lobata* Née (Fagaceae).

Prionus Le Contei Lameere, 1912a: 244.

Prionus lecontei; Linsley, 1935b: 161; Hovore & Giesbert, 1976: 350 (hosts); Hovore, 1988: 3 (distr.); Chemsak, Linsley & Noguera, 1992: 21 (cat.); Noguera & Chemsak, 1996: 396 (cat.). Chemsak, 1996: 108, pl. 7, figs 5, 6; Linsley & Chemsak, 1997: 424 (hosts); Monné, M.A., 2002: 25 (cat. hosts); Monné, M.A., 2006: 87 (cat.).

Prionus (Prionus) lecontei; Monné, Monné, M.A., & Giesbert, 1994: 15 (checklist); M.A. & Hovore, 2006: 19 (checklist); Santos-Silva, Nearn & Swift, 2016: 74, figs 49-51, 130-132 (lectotype).

7. *Prionus pocularis* Dalman, 1817

Type locality - Holotype: United States, Georgia (NHRS). **Distribution** - Eastern North America to Texas, **Host plants** - *Pinus* sp (Pinaceae)

Prionus pocularis Dalman, 1817: 148; Haldeman, 1847a: 31; White, 1853: 17; Lacordaire, 1868: 61; Leng, 1884: 57; Beutenmuller, 1896: 74; Smith, 1900: 285; Ulke, 1903: 25; Evans, 1906: 99; Fall & Cockerell, 1907: 191; Leng, 1910: 77; Blatchley, 1910: 1012; Smith, 1910: 324; Lameere, 1912a: 328; Dow, 1913: 78; Frost, 1915: 207; Craighead, 1915: 20; Greene, 1918: 257; Nicolay, 1919: 63; Britton, 1920: 266; Kirk & Knul, 1926: 21; Leonard, 1928: 433; Doane *et al.*, 1936: 165; Brimley, 1938: 210; Loding, 1945: 113; Knul, 1946: 146; Craighead, 1950: 262; Beal *et al.*, 1952: 134; Linsley, 1957: 8; Dillon & Dillon, 1961: 650, pl. 57; Linsley, 1962: 41; Gosling, 1973: 67; Turnbow & Franklin, 1980: 338; Chemsak, Linsley & Noguera, 1992: 21; MacRae, 1993: 227; Monné, M.A., 1995: 53; Browne & Peck, 1996: 2158; Chemsak, 1996: 105, pl. 7, figs 1,2; Yanega, 1996: 27, pl. 1, fig. 10; Linsley & Chemsak, 1997: 425; Schiefer, 1998: 115; Peck & Thomas, 1998: 116; Vlasák & Vlasakova, 2002: 204; Bousquet, Laplante, Hammond & Langor, 2017: 30, pl. 2

Prionus (Prionus) pocularis; Casey, 1912: 238; 1924: 213; Monné, M.A., & Giesbert, 1994: 15 (checklist); Monné, M.A. & Hovore, 2006: 19 (checklist); Holt, 2013: 243; Klingeman, *et al.*, 2017: 294

Prionus laevigatus Harris, 1837: 83, pl. 1, fig. 6; Haldeman, 1847a: 31; LeConte, 1851: 109; White, 1853: 16; Bland, 1861: 93; Lacordaire, 1868: 61

Syntypes – Syntypes male and female: United States, Massachusetts (MCZN)

Prionus obliquicornis LeConte, 1851: 108; White, 1853: 17; Lacordaire, 1868: 61

Prionus (Prionus) obliquicornis; Casey, 1912: 239

Type locality - Holotype male: United States, Georgia (MCZN)

Prionus curticornis LeConte, 1851: 109; Lacordaire, 1868: 61

Prionus (Prionus) curticornis Casey, 1912: 233

Type locality - Holotype female: United States, Georgia. (MCZN)

Prionus (Prionus) validiceps Casey, 1912: 233; 1924: 214; Lingafelter *et al.*, 2014: 339, fig. 177w (holotype)

Type locality - Holotype male: United States (USNM)

Prionus (Prionus) pocularis prolixus Casey, 1912: 239; Lingafelter *et al.*, 2014: 301, fig. 134m (holotype)

Type locality - Holotype female: United States, (USNM)

Prionus (Prionus) bicolor Casey, 1912: 239; 1924: 213; Lingafelter *et al.*, 2014: 27, fig. 28a (holotype)

Type locality - Holotype female: United States, Indiana (USNM)

Prionus (Homaesthesis) LeConte, 1873

Homaesthesis LeConte, 1873b: 288; LeConte & Horn, 1883: 273. Leng, 1884: 57; Casey, 1912: 231; Craighead, 1923: 28

Prionus (Homaesthesis); Linsley, 1962: 49; Hatch, 1971: 93; Hovore & Turnbow, 1984: 4; Monné, M.A., 1995: 56; Chemsak, 1996: 114; Monné, M.A. & Hovore, 2006: 19 (checklist); Santos-Silva, Nearn & Swift, 2016: 10.

Type species - *Prionus emarginatus* Say, 1824 (Casey designation, 1912: 254).

Prionina Casey, 1912: 231

Type species - *Prionus palparis* Say, 1824 (original designation)

1. *Prionus (Homaesthesis) arenarius* Hovore, 1981

Type locality - Holotype male: United States, Texas: Ward County, Monahans Sandhills State Park. (CASC), **Distribution** - United States (Texas and New Mexico)

Prionus (Homaesthesis) arenarius Hovore, 1981: 453, fig. 1; Hovore & Turnbow, 1984: 4, figs 1, 3; Chemsak, Linsley & Noguera, 1992: 21 (cat.); Monné, M.A., & Giesbert, 1994: 15 (checklist); Monné, M.A., 1995: 56 (cat.); Chemsak, 1996: 117, pl. 8, figs 3,4; Monné, M.A. & Hovore, 2006: 19 (checklist); Santos-Silva Nearn & Swift, 2016: 12, figs 17-19; 52-53

2. *Prionus (Homaesthesis) emarginatus* Say, 1824

Type locality - Type: United States, Arkansas: Arkansas river, near the mountains. (depository unknown). **Distribution** - United States (Colorado, Arizona, Idaho, New Mexico, Kansas, Wyoming, Missouri, Nebraska, Utah, Texas).

Prionus emarginatus Say, 1824: 327; Haldeman, 1847a: 31; LeConte, 1851: 107; White, 1853: 17; LeConte, 1859a: 19, pl. 2, fig. 13; 1859b: 184; Lacordaire, 1868: 61; Packard, 1877: 803, pl. 70, fig. 6; 1881: 161, fig. 73; 1890: 703, fig. 239; Lameere, 1912a: 247; Haack, 2017: 110; Rice, MacRae & Merickel, 2017: 669 (distr.)

Homaesthesis emarginata; LeConte, 1873b: 288; 1876: 520; Snow, 1877: 19; 1883: 42; LeConte & Horn, 1883: 273; Leng, 1884: 57; Blanchard, 1887: 86; Beutenmuller, 1896: 74; Snow, 1906: 179; Fall & Cockerell, 1907: 191; Casey, 1912: 254; Craighead, 1923: 28, pl. 1, fig. 10; pl. 12, figs 10-12; 1950: 239, fig. 51a; Linsley, 1958: 110; Gwynne & Hostetler, 1978: 347

Prionus (Homaesthesis) emarginatus; Lameere, 1913: 78; Linsley, 1957: 10; 1962: 50; Hatch, 1971: 94; Hovore & Turnbow, 1984: 4; Chemsak, Linsley & Noguera, 1992: 21 (cat.); Monné, M.A., & Giesbert, 1994: 15 (checklist); Monné, M.A., 1995: 56 (cat.); Chemsak, 1996: 118, pl. 8, figs 5, 6; Heffern, 1998: 8; Monné, M.A. & Hovore, 2006: 19 (checklist); Santos-Silva, Nearn & Swift, 2016: 12, figs 1-4; 83-86

Prionus innocuous LeConte, 1862: 43. Lacordaire 1868: 61

Type locality - Holotype female: United States: New Mexico. (MCZN)

Homaesthesis pubicollis Casey, 1912: 254; Lingafelter *et al.*, 2014: 304, fig. 137 m (holotype)

Type locality - Holotype male: United States, Colorado (USNM).

Homaesthesis debiliceps Casey, 1912: 255; Lingafelter *et al.*, 2014: 50, fig. 53e (holotype)

Type locality - Holotype male: United States, Colorado (USNM)

3. *Prionus (Homaesthesis) geminus* Santos-Silva, Nearn & Swift, 2016

Type locality - Holotype male: United States, Idaho: Pocatello (Bannock County). (EMEC).

Distribution - United States (Idaho, Utah, Arizona, Texas, Colorado)

Prionus (Homaesthesis) integer; Linsley, 1962: 51; Chemsak, 1996: 119 (not LeConte)

Prionus (Homaesthesis) geminus Santos-Silva, Nearn & Swift, 2016: 12, figs 5-8, 87-90

Prionus geminus; Rice, MacRae & Merickel, 2017: 669 (distr.)

4. *Prionus (Homaesthesis) integer* LeConte, 1851

Type locality - Holotype male: United States, Colorado: near Pike's Peak, Rocky Mountains. (MCZN). **Distribution** - United States (Kansas and New Mexico to Colorado and eastern Oregon). **Host plants** - *Artemisia tridentata* Nuttall, *Chrysothamnus viscidiflorus* (Hooker) Nuttall (Asteraceae)

Prionus integer LeConte, 1851: 197; White, 1853: 17; LeConte, 1859a: 48; Lacordaire, 1868: 61; Knowlton & Thatcher, 1936: 278; Knowlton & Wood, 1950: 10
Homaesthis integra; LeConte, 1873b: 288; LeConte & Horn, 1883: 273; Casey, 1912: 253
Homaesthis integer; Popenoe, 1877: 33; Leng, 1884: 57, pl. 3, fig. 2; Blanchard, 1887: 85; Knaus, 1904: 156; Fall & Cockerell, 1907: 191; Lameere, 1912a: 246;
Prionus (Homaesthis) integer; Lameere, 1913: 78 (cat.); 1919: 139; Barr & Penrose, 1959: 92; Hatch, 1971: 94; Penrose & Westcott, 1974: 236; Hovore & Turnbow, 1984: 4; Chemsak, Linsley & Noguera, 1992: 22 (cat.); Monné, M.A., & Giesbert, 1994: 16 (checklist); Monné, M.A., 1995: 56 (cat.); Heffern, 1998: 6; Monné, M.A. & Hovore, 2006: 19 (checklist); Santos-Silva, Nearn & Swift, 2016: 12, figs 79-82; Rice, MacRae & Merickel, 2017: 669 (distr.)

5. *Prionus (Homaesthis) palparis* (Say, 1824)

Syntypes locality - Syntypes male and female: United States, Arkansas: Arkansas river near the mountains. (depository unknown). **Distribution** - United States (Arizona, Colorado, Kansas, Nebraska, New Mexico, Oklahoma, Utah).

Prionus palparis Say, 1824: 327; Haldeman, 1847a: 31; LeConte, 1851: 108; White, 1853: 15; LeConte, 1859a: 48; 1859b: 185; Lacordaire, 1868: 61; LeConte, 1873b: 288; LeConte & Horn, 1883: 274; Leng, 1884: 57; Blanchard, 1887: 86; Fall & Cockerell, 1907: 191; Lameere, 1912a: 246; Knowlton & Wood, 1950: 10;

Prionina palparis; Casey, 1912: 292

Peionus (Prionina) palparis; Lameere, 1919: 139

Prionus (Homaesthis) palparis; Linsley, 1962: 58; Hovore & Turnbow, 1984: 4; Chemsak, Linsley & Noguera, 1992: 22; Monné, M.A., & Giesbert, 1994: 16 (checklist); Monné, M.A., 1995: 57; Chemsak, 1996: 126, pl. 6, figs 9.10; Heffern, 1998: 6; Monné, M.A. & Hovore, 2006: 19 (checklist); Santos-Silva, Nearn & Swift, 2016: 22, figs 9-11, 75-76

6. *Prionus (Homaesthis) simplex* (Casey, 1912)

Type locality - Holotype male: United States, Kansas, (USNM), **Distribution** - Western Kansas and southwestern Nebraska to central Utah, Colorado, New Mexico and Arizona

Prionina simplex Casey, 1912: 253; Knaus, 2014: 91;

Prionus (Homaesthis) simplex; Linsley, 1962: 53; Hovore & Turnbow, 1984: 4; Chemsak, Linsley & Noguera, 1992: 22 (cat.); Monné, M.A., & Giesbert, 1994: 16 (checklist); Monné, M.A., 1995: 57 (cat.); Chemsak, 1996: 123, pl. 9, fig. 3; Heffern, 1998: 174; Monné, M.A. & Hovore, 2006: 19 (checklist); Lingafelter *et al.*; 2014: 320, fig. 155w (holotype); Santos-Silva, Nearn & Swift, 2016: 12, figs

Prionus (Homaesthis) rhodocerus Linsley, 1957: 10; 1962: 53; Hovore, 1981: 457; Hovore & Turnbow, 1984: 4; Chemsak, Linsley & Noguera, 1992: 22 (cat.); Monné, M.A., 1995: 57 (cat.); Chemsak, 1996: 122; Heffern, 1998: 6; Monné, M.A. & Hovore, 2006: 20 (checklist);

Type locality - Holotype male: United States, Utah: Red Creek. (CASC)

Prionus (Homaesthis) linsleyi Hovore, 1981: 455, fig. 2; Hovore & Turnbow, 1984: 4; Chemsak, Linsley & Noguera, 1992: 16; Monné, M.A., 1995: 57; Chemsak, 1996: 124, pl. 9, figs 4,5

Type locality - Holotype male: United States, California: Coconino County, 6.5 miles South of Moenkopi (CASC)

7. *Prionus (Homaesthis) spinipennis* Hovore & Turnbow, 1984

Type locality- Holotype male: United States, Texas: Ward County, Monahans Sandhills State Park. (CASC). **Distribution** - United States (Western Texas to eastern New Mexico)

Prionus (Homaesthis) spinipennis Hovore & Turnbow, 1984: 1, figs 2, 4; Chemsak, Linsley & Noguera, 1992: 23 (cat.); Monné, M.A., & Giesbert, 1994: 16 (checklist); Monné, M.A., 1995: 57 (cat.); Chemsak, 1996: 121, pl. 8, figs 11, 12; Monné, M.A. & Hovore, 2006: 20 (checklist); Santos-Silva, Nearn & Swift, 2016: 12, figs 14-14. 69-72

SOLENOPTERINI Lacordaire, 1868

Solénoptérides Lacordaire, 1868:171, 180.

Solenopterini; LeConte, 1873b: 286; LeConte & Horn, 1883: 271; Galileo & Martins, 1993: 90 (rev.); 1993: 199 (rev.); 1993: 425 (rev.); 1994: 109 (rev.); Monné, M.A., 2006: 91 (cat.); Bousquet *et al.*, 2009: 20; Bouchard *et al.*, 2011: 461.

Type-genus: *Solenoptera* Audinet-Serville, 1832

Type-species: *Prionus canaliculatus* Fabricius, 1787 designated by Villiers (1980a: 144). Availability (under Article 11.7.2).

Dérancistrines Lameere, 1912c:181; *Nom. nudum*. Comment. This name is unavailable under Article 11.7 (vernacular name proposed after 1899).

Derancistri; Lameere, 1919: 99.

Derancistrini; Melzer, 1919: 99; Knull, 1946: 143, 144; Linsley, 1961b: 629 (mimet.).

Type-genus: *Derancistrus* Audinet-Serville, 1832.

Type-species: *Prionus elegans* Palisot de Beauvois, 1819 (monotypy).

Elateropsis Chevrolat, 1862

Elateropsis Chevrolat, 1862: 269; Thomson, 1864: 276; Lacordaire, 1868: 183; LeConte, 1873b: 290; LeConte & Horn, 1883: 275; Leng, 1884: 8, 60; Gahan, 1890: 27; Galileo & Martins, 1994a: 109 (rev.); Monné, M.A., 2006: 92 (cat.); Monné, M.A., 2012: 132.

Derancistrus (*Elateropsis*); Lameere, 1909: 14; 1913: 51; 1919: 98; Zayas, 1957: 168; Linsley, 1962: 60; Zayas, 1975: 33.

Type-species - *Cerambyx lineatus* Linnaeus, 1758 (subsequent designation, Thomson, 1864: 276).

1. *Elateropsis rugosus* Gahan, 1890

Type locality - Holotype female: not stated. (BMNH). **Distribution** - United States (Florida), Bahamas, Cuba (?). **Host plants** - *Metopium toxiferum* (Linnaeus) Krug & Urban (Anacardiaceae), *Bursera simaruba* (Linnaeus) Sargent (Burseraceae), *Coccoloba diversifolia* Jacquin (Polygonaceae).

Elateropsis rugosa Gahan, 1890: 28; 1895a: 89; Leng & Mutchler, 1914: 444 (distr.); Schwarz *in* Holland, 1917: 341 (distr.).

Derancistrus (*Elateropsis*) *rugosus*; Lameere, 1909: 14; 1913: 51 (cat.); 1919: 99; Cazier & Lacey, 1952: 7 (distr.); Zayas, 1957: 170, pl. 12; 1975: 39; Turnbow & Hovore, 1979: 219 (distr., hosts); Chemsak, Linsley & Noguera, 1992: 18 (cat.).

Derancistrus rugosus; Browne, Peck & Ivie, 1993: 40 (distr.).

Elateropsis rugosus; Galileo & Martins, 1994a: 115, figs 163–166, 209; Monné, M.A., 2002: 26 (cat. hosts); Monné, M.A., & Giesbert, 1994: 11 (checklist); Peck, 2005: 165 (distr.); Monné, M.A. & Hovore, 2006: 22 (checklist); Monné, M.A., 2006: 95 (cat.).

2. *Elateropsis scabrosus* Gahan, 1890

Syntypes locality – Syntypes male and female: Cuba. (BMNH). **Distribution** - United States (Florida), Bahamas, Cuba. **Host** - *Bursera simaruba* (Linnaeus) Sargent (Burseraceae).

Elateropsis scabrosa Gahan, 1890: 29; 1895a: 89; Schaeffer, 1908: 331; Leng & Mutchler, 1914: 444 (distr.).

Derancistrus (*Elateropsis*) *scabrosus*; Lameere, 1909:18; 1913: 51 (cat.); 1919: 99; Cazier & Lacey, 1952: 7; Zayas, 1957: 170, pl. 14; Linsley, 1962: 61, fig. 22; Chemsak, 1967b: 182, fig. 2; Zayas, 1975: 40; Chemsak, Linsley & Noguera, 1992: 18 (cat.).

Derancistrus scabrosus; Chemsak, 1969a: 186 (distr.); Thomas, 1977: 86, fig. 2 (biol., distr.); Browne, Peck & Ivie, 1993: 41 (distr.).

Elateropsis scabrosus; Galileo & Martins, 1994a: 118, figs 167, 168, 170, 171, 211 (syn.); Monné, M.A., 2002: 26 (cat. hosts); Monné, M.A., & Giesbert, 1994: 11 (checklist); Peck, 2005: 166 (distr.); Monné, M.A., 2006: 95 (cat.); Monné, M.A. & Hovore, 2006: 22 (checklist); Devesa, Fonseca & Barro, 2015b: 29, 461, 76, figs 1-10.

Elateropsis fuliginosus; Chevrolat, 1862: 271 (*partim*); LeConte, 1873b: 290; LeConte *in* Schwarz, 1878: 470; LeConte & Horn, 1883: 275; Leng, 1884: 60 (not Fabricius, 1775).
Derancistrus (Elateropsis) punctatus; Zayas, 1957: 171 (not Gahan, 1890).

***Sphenostethus* Haldeman, 1848**

Sphenostethus Haldeman, 1848: 126; Le Conte, 1851: 106; Thomson, 1861: 288; 1864: 276, 467; Lacordaire, 1868: 184; LeConte 1873: 290; Le Cote & Horn, 1883: 275; Leng, 1884: 60; Lameere, 1909: 20; Blatchley, 1910: 1012; Knull, 1946: 144; Galileo & Martins, 1993: 211; 1994: 146; Monné, M.A. & Hovore, 2006: 23 (checklist)

Derancistrus (Sphenostethus); Linsley, 1962: 59

Type species - *Sphenostethus taslei* Buquet, 1841 (monotypy) (= *Sphenostethus serripennis* Haldeman, 1846)

***Hoplopteryx* Westwood, 1845: 40** (not Agassiz, 1838)

1. *Sphenostethus taslei* (Buquet, 1841)

Type locality - Holotype: United States, Louisiana (M NHN). **Distribution** - Atlantic coast Pennsylvania and norther Florida to Missouri and Texas, **Host plants** - *Cercis canadensis* Linnaeus (Caesalpiniaceae), *Sesbania drummondii* (Rydberg) C ory (Fabaceae), *Castanea dentata* (Marshall) Borkhausen, *Fagus ferrugineus* Aiton, *Quercus alba* Linnaeus (Fagaceae)

Solenoptera taslei Buquet, 1841: XXXI X; White, 1851: 54; Strauch, 1861: 126

Sphenostethus taslei; Thomson, 1864: 276; Lacordaire, 1868: 185; LeConte, 1873: 290; LeConte & Horn, 1883: 275; Leng, 1884: 6; Luggier, 1884: 203 (hosts); Hamilton, 1895: 337; Smith, 1900: 285; Dury 1902: 158; Ulke, 1903 25; Blatchley, 1910: 1012; Smith, 1910: 324; Fisher & Kirk, 1912: 309; Craighead, 1915: 21; 1923: 29; Galileo & Martins, 1993: 215; Monné, M.A., & Giesbert, 1994: 11 (checklist); Monné, M.A., 1995: 40; Chemsak, 1996: 132; Yanega, 1996: 27, pl. 2, fig. 14; Schiefer, 1998 115; Peck & Thomas, 1998: 116; MacRae & Rice, 2007: 228; Holt, 2013: 243. Klingeman *et al.*, 2017: 294

DBerancistrus (Sphenostethus) taslei; Lameere, 1909: 20; 1913: 52; 1919: 99; Linsley. 1962: 59; Chemsak, Linsley & Noguera, 1992: 18; MacRae, 1993: 226

Derancistrus taslei; Champlain, Kirk & Knull, 1925: 105; Kirk & Knull, 1926: 21; Hatch, 1930: 26; Brimley, 1938: 209; Loding, 1945: 113; Knull, 1946: 144; Sherman, 1946: 127; Alexander, 1958: 46; Solomon, Doolittle & Spilman, 1976: 290; Turnbow & Franklin, 1980: 338; Rice, 1981: 459; Arnett, 1985: 360; Linsley & Chemsak, 1995: 364 (hosts);

Hoplopteryx denticulatus Westwood, 1841: 40; Haldeman, 1847: 375

Type locality ? (OXUM)

Sphenostethus serripennis Haldeman, 1648

Type locality - Holotype: United States, Soutghern Pennsylvania (MCZN)

SPONDYLIDINAE Audinet-Serville, 1832

Spondylii Audinet-Serville, 1832: 123

Spondylidinae; Bousquet, Laplante, Hammond & Langor, 2017: 78

Type-genus: *Spondylis* Fabricius, 1775

Type-species: *Attelabus buprestoides* Linnaeus, 1758 designated by Latreille (1810: 431).

ASEMINI Thomson, 1861

Asemitae Thomson, 1861a: 139 (key), 259; 1864: 266; 1867a: 9.

Asemites; Fairmaire *in* Jacquelin DuVal, 1864a: 122.

Asemides; Lacordaire, 1868: 205.

Asemi; LeConte, 1873b: 293; LeConte & Horn, 1883: 278.

Asemini; Gahan, 1906: 94; Aurivillius, 1912: 15 (cat.); Casey, 1912: 256; Bradley, 1930: 228, 230; Linsley, 1939: 65; Knull, 1946: 148; Dillon & Dillon, 1961: 582; Linsley, 1962: 67; Arnett, 1962: 856; Hatch, 1971: 94; Monné, M.A., 1994: 18 (cat.); Monné, M.A., 2006: 107 (cat.); Bousquet, Heffern, Bouchard & Nearn., 2009: 21; Bouchard *et al.*, 2011: 464; Bousquet, Laplante, Hammond & Langor, 2017: 80 (key gen)

Type-genus: *Asemum* Eschscholtz, 1830.

Type-species: *Cerambyx striatum* Linnaeus, 1758 designated by Westwood (1838: 40).

Criomorphates Mulsant, 1863: 79, 115 (based on *Criomorphus* Mulsant, 1839). *Nomen nudum*.
Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Mulsant 1863).

Criomorphaires Planet, 1924: 232.

Criomorphini; Portevin, 1927: 20, 36.

Type-genus: *Criomorphus* Mulsant, 1839 (junior homonym of *Criomorphus* Curtis, 1831 [Hemiptera]).

Type-species: *Callidium aulicum* Fabricius, 1775 by monotypy. Comment. This name is permanently invalid because it is based on a preoccupied type genus (Article 39).

Criocéphalites Fairmaire, 1864a: 125, 192 (based on *Criocephalus* Mulsant, 1839). *Nomen nudum*. Comment. This name is unavailable under Article 11.7 (not subsequently Latinized and attributed to Fairmaire 1864).

Criocephalinae Sharp, 1905: 147.

Type-genus: *Criocephalus* Mulsant, 1839

Type-species: *Cerambyx rusticus* Linnaeus, 1758 (monotypy).

Tetropiina Seidlitz, 1891: 179; Reitter, 1912: 42 (*partim*).

Type-genus: *Tetropium* Kirby, 1837.

Type-species: *Callidium cinnamopterum* Kirby, 1837 (see ICZN 1988: 71).

Nothorhinini Zagajkevich, 1991: 110.

Type-genus: *Nothorhina* Redtenbacher, 1845.

Type-species: *Callidium muricatum* Dalman, 1817 designated by Thomson (1864: 267).

Comment. This name has been listed in synonymy with the Asemini by Vives and Alonso-Zarazaga (2000: 569).

***Arhopalus* Audinet-Serville, 1834**

Arhopalus Audinet-Serville, 1834b: 77; Linsley, 1940: 254; Knoll, 1946: 149; Linsley, 1962: 68; Arnett, 1962: 856; Hatch, 1971: 94; Chemsak, 1996: 18; Monné, M.A., 2006: 108 (cat.); Monné, M.A., 2012: 133; Bousquet. Laplante, Hammond & Langor, 2017: 80 (key spp)

Type-species - *Cerambyx rusticus* Linnaeus, 1758 (Westwood subsequent designation, Westwood, 1838: 40).

Criocephalum Dejean, 1835: 328 (cat.); Bousquet & Bouchard. 2013: 79.

Type-species - *Cerambyx rusticus* Linnaeus, 1758 (monotypy).

Criocephalus Mulsant, 1839: 63; LeConte, 1850a: 35; Thomson, 1861a: 260; Mulsant, 1863: 125, 431; Thomson, 1865: 464; Lacordaire, 1868: 210; LeConte, 1873a: 169; 1873b: 293, 294; Provancher, 1877: 578, 584; Bates, 1879: 15; LeConte & Horn, 1883: 278; Leng, 1884: 63; Wickham, 1897b: 105; Gahan, 1906: 96; Blatchley, 1910: 1014; Casey, 1912: 265; Bradley, 1930: 230; Chagnon, 1936: 256; Chagnon & Robert, 1962: 253, 256; Zayas, 1975: 48.

Type-species - *Cerambyx rusticus* Linnaeus, 1758 (subsequent designation, Thomson, 1864: 268).

Hylescopus Gistel, 1856 : 376

Type-species - *Cerambyx rusticus* Linnaeus, 1758 (monotypy).

Cephalocrius Sharp, 1905: 147.

Type-species - *Criocephalus syriacus* Reitter, 1895 (monotypy).

1. *Arhopalus asperatus* (LeConte, 1859)

Syntypes locality - Syntypes male and female: United States, Kansas. (MCZN). **Distribution** - Coniferous belt of western North America, Mexico (Baja California, Chiapas, Chihuahua, Veracruz), Honduras. **Host plants** - *Abies concolor* Gordon & Glen, *Abies magnifica* A. Murray, *Picea* sp., *Pinus ponderosa* Douglas ex Lawson & P. Lawson, *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae).

Criocephalus asperatus LeConte, 1859a: 19; 1873a: 170; Snow, 1877: 19 (distr.); Bates, 1884: 239; Leng, 1884: 63; Slossom, 1895: 6 (distr.); Harrington, 1899: 108 (distr.); Fall, 1901: 27 (distr.); Knaus, 1904: 156 (distr.); Sharp, 1905: 160; Snow, 1906b: 170 (distr.); Fall & Cockerell, 1907: 191 (distr.); Garnett, 1918: 174 (distr.); Hardy, 1926: 27; Essig, 1926: 451 (biol.); Knowlton, 1930: 56 (distr.); Beaulne, 1932: 198 (hosts); Davis, 1932: 85 (distr.);

- Mank, 1934: 80 (distr.); Knowlton & Thatcher, 1936: 278 (distr.); Doane *et al.*, 1936: 168 (biol.); Moore, 1937: 87 (distr.); Carl & Hardy, 1945: C36 (biol.).
- Arhopalus asperatus*; Knowlton & Wood, 1950: 10 (distr.); Keen, 1952: 195, fig. 93 (hosts); Linsley, 1957: 11 (syn.); Linsley, Knull & Statham, 1961: 7; Linsley, 1962: 69, figs 25, 26; Hatch, 1971: 95, pl. 9, fig. 5; Kirk & Balsbaugh, 1975: 97 (distr.); Furniss & Carolin, 1977: 291 (hosts); Lewis, 1979: 22 (distr.); Chemsak, Linsley & Mankins, 1980: 28 (distr.); Hovore, 1988: 3 (distr.); Chemsak, Linsley & Noguera, 1992: 24 (cat.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 19, pl. 1, fig. 9 (rev.); Linsley & Chemsak, 1997: 345 (hosts); Heffern 1998: 7 (distr.); Monné, M.A., 2002: 32 (cat. hosts); Turnbow, Cave & Thomas, 2003: 5 (distr.); Monné, M.A., 2006: 108 (cat.); Monné, M.A. & Hovore, 2006: 23 (checklist); Hovore, 2006: 371 (distr.); Gutiérrez & Noguera, 2015: 138 (distr.); Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet. Laplante, Hammond & Langor, 2017: 81, pl. 14
- Criocephalus honduranus* Casey, 1912: 265; Lingafelter *et al.*, 2014: 76, figs 82u, v (holotype).
Type locality - Holotype male: Honduras. (USNM).
- Criocephalus filitarsis* Casey, 1912: 266; Lingafelter *et al.*, 2014: 62, figs 66k, l (holotype).
Type locality - Holotype female: United States, Texas: El Paso. (USNM).

2. *Arhopalus foveicollis* (Haldeman, 1847)

- Syntypes localities** - Syntypes: United States, Maryland, Virginia. (MCZN). **Distribution** - Coniferous forests of North America except the Pacific Coast. **Host plants** - *Picea* sp., *Pinus rigida* Miller, *Pinus strobus* Linné (Pinaceae).
- Criocephalus foveicollis* Haldeman, 1847a: 35.
- Arhopalus foveicollis*; Linsley, 1957: 11 (syn.); 1962: 70, fig. 26; Gosling, 1973: 68, fig. 7 (biol., distr.); Stein & Tigestad, 1976: 3 (distr.); Laliberté, Chantal & LaRochelle, 1977: 91 (biol.); Furniss & Carolin, 1977: 291; Gosling, 1986: 155 (hosts); Chemsak, Linsley & Noguera, 1992: 24 (cat.); MacRae, 1993: 227 (distr.); Monné, M.A., 1994: 19 (cat.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Chemsak, 1996: 21, pl. 1, fig. 10; Yanega, 1996: 43, pl. 2, fig. 16; Linsley & Chemsak, 1997: 345 (hosts); Heffern, 1998: 7 (distr.); Vlasák & Vlasakova, 2002: 209 (distr.); Monné, M.A. & Hovore, 2006: 23 (checklist); Lingafelter, 2007: 15, 138, 2 figs, pl. 3d (hosts); Webster, 2016: 486; Klingeman *et al.*, 2017: 294 (distr.); Bousquet. Laplante, Hammond & Langor, 2017: 81, pl. 24
- Callidium agreste* Kirby, 1837: 170 (preoccupied).
- Criocephalus agrestis*; Haldeman, 1847a: 35; LeConte, 1850b: 234 (distr.); 1850a: 36; White, 1855: 311; LeConte, 1859a: 48 (distr.); Packard, 1870: 119 (distr.); Pettit, 1871: 105 (distr.); Bethune, 1872: 55; LeConte, 1873a: 170; Horn, 1873: 717; 1876: 169; Provancher, 1877: 585; Packard, 1881: 158 (biol.); Harrington, 1881: 33 (biol.); Leng, 1884: 63; Saunders, 1884: 53; 1887: 29 (distr.); Packard, 1890: 699, pl. 17, figs 3, 3a; Harrington, 1890: 185 (distr.); Evans, 1895: 173 (distr.); Chagnon, 1897: 16 (distr.); Wickham, 1897c: 106; 1899: 199 (distr.); Harrington, 1899: 108 (distr.); Smith, 1900: 285 (distr.); Ulke, 1903: 25 (distr.); Chagnon, 1905: 35 (distr.); Sharp, 1905: 160; Fall & Cockerell, 1907: 191 (distr.); Morris, 1908: 446; Davis, 1909: 42 (distr.); Wickham, 1909a: 28 (distr.); Morris, 1909: 60 (biol.); Sherman, 1910: 194 (distr.); Smith, 1910: 324; Blatchley, 1910: 1015; Fisher & Kirk, 1912: 309 (distr.); Frost, 1915: 207; Chagnon, 1917: 230 (distr.); Britton, 1920: 266 (distr.); Casey, 1924: 230 (syn.); Fall, 1926: 202 (distr.); Kirk & Knull, 1926: 21 (distr.); Leonard, 1928: 434 (distr.); Brown, 1930: 244 (distr.); Beaulne, 1932: 198 (hosts); Brown, 1934: 229 (distr.); Mank, 1934: 80 (distr.); Doane *et al.*, 1936: 268; Chagnon, 1936: 256, pl. 17, fig. 1; 1939: 86 (distr.); Procter, 1946: 177 (biol.); Clark, 1956: 41 (distr.); Chagnon & Robert, 1962: 256, pl. 17, fig. 1.
- Criocephalus rusticus* var. *agrestis*; Hamilton, 1890: 44.
- Arhopalus agrestis*; Knull, 1946: 119, pl. 3, fig. 7; Knowlton & Wood, 1950: 10 (distr.); Hicks, 1955: 165 (distr.); Gardiner, 1957: 251 (distr.).
- Syntypes locality** - Syntypes: Canada, Nova Scotia. (BMNH).
- Criocephalus lacustrinus* Casey, 1912: 267; Lingafelter *et al.*, 2014: 85, fig. 93e (holotype).
Type locality - Holotype female: United States, Wisconsin: Bayfield. (USNM).

Criocephalus cervinus Casey, 1912: 267; Lingafelter *et al.*, 2014: 39, fig. 40w (holotype).

Type locality - Holotype male: United States, Indiana. (USNM).

3. *Arhopalus hesperus* Chemsak & Linsley, 1965

Type locality - United States, Western Nevada and California **Host plants** - *Pinus monophylla* Torrey & Fremont (Pinaceae)

Arhopalus rusticus hesperus; Chemsak & Linsley, 1965: 141; Tyson, 1970: 297 (hosts); Chemsak, Linsley & Noguera, 1992: 23 (cat.); Monné, M.A., 1994: 20; Chemsak, 1996: 27, pl. 2, fig. 3; Monné, M.A. & Hovore, 2006: 23 (checklist);

Arhopalus hesperus; Bousquet, Laplante, Hammond & Langor, 2017: 18

4. *Arhopalus montanus* (LeConte, 1873)

Syntypes locality - Syntypes male and female: United States, Colorado. (MCZN).

Distribution - Rocky Mountain region from Colorado to New Mexico, Arizona and northern Mexico (Chihuahua, Chiapas, Nuevo León, Oaxaca, Tamaulipas). **Host plants** - *Pinus arizonica* Engelman ex Rothrock, *P. leiophylla chihuahuana* (Engelman) A. E. Murray, *P. ponderosa* Douglas ex Lawson & P. Lawson, *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae).

Criocephalus montanus LeConte, 1873a: 170; Lameere, 1883: 9 (cat.); Leng, 1884: 63, 64; Craighead, 1923: 33.

Arhopalus montanus; Knowlton & Wood, 1950: 10 (distr.); Linsley, 1957: 12 (syn.); Linsley, Knull & Statham, 1961: 7 (distr., hosts); Bousquet, Laplante, Hammond & Langor, 2017: 81

Arhopalus rusticus montanus; Linsley, 1972: 76, fig. 27; Furniss & Carolin, 1977: 291; Lewis, 1979: 22 (distr., hosts); Chemsak, Linsley & Noguera, 1992: 24 (cat.); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 26, pl. 2, fig. 2 (rev.); Linsley & Chemsak, 1997: 346 (hosts); Heffern, 1998: 7 (distr.); Monné, M.A., 2002: 33 (cat. hosts); Monné, M.A., & Giesbert, 1994: 22 (checklist); Monné, M.A., 2006: 109 (cat.); Monné, M.A. & Hovore, 2006: 23 (checklist); Gutiérrez & Noguera, 2015: 138 (distr.); García Morales *et al.*, 2015: 99 (distr.).

Arhopalus montanus; Bousquet, Laplante, Hammond & Langor, 2017: 81

Nothorhina spissicornis Casey, 1912: 264; Lingafelter *et al.*, 2014: 323, figs 159g, h (lect. designation).

Type locality - Lectotype male: United States, Colorado: Boulder Co. (USNM).

Nothorhina spissicornis longicornis Casey, 1912: 246; Lingafelter *et al.*, 2014: 323, figs 159m, n (holotype).

Type locality - Holotype male: United States, Colorado: Boulder Co. (USNM).

Nothorhina gracilipes Casey, 1912: 265; Lingafelter *et al.*, 2014: 70, figs 75u, v (holotype).

Type locality - Holotype male: United States, New Mexico: Las Vegas. (USNM).

Criocephalus cavicollis Casey, 1912: 266; Lingafelter *et al.*, 2014: 38, figs 39s, t (holotype).

Type locality - Holotype male: United States, Colorado. (USNM).

Criocephalus coloradensis Casey, 1924: 231; Lingafelter *et al.*, 2014: 43, figs 44m, n (holotype).

Type locality - Holotype female: United States, Colorado: Florissant. (USNM).

5. *Arhopalus nubilus* (LeConte, 1850)

Syntypes locality - Syntypes: not stated. (MCZN). **Distribution** - Southeastern United States, Bahamas, Jamaica. **Host plants** - *Pinus palustris* Miller (Pinaceae).

Criocephalus nubilus LeConte, 1850a: 36; White, 1855: 311; LeConte, 1873a: 170; Riley, 1880: 238 (hosts); Packard, 1881: 162 (biol.); Leng, 1884: 63, 64; Packard, 1890: 704 (biol.); Beutenmuller, 1896: 74 (hosts); Castle & Laurent, 1897: 8 (distr.); Sharp, 1905: 158; Fall & Cockerell, 1907: 191 (distr.); Wickham, 1909b: 402 (distr.); Davis & Leng, 1912: 121 (hosts); Dozier, 1918: 334 (distr.); Gowdey, 1926: 21 (distr.); Loding, 1933: 148 (distr.); Brimley, 1938: 210 (distr.); Loding, 1945: 114 (distr.); Sherman, 1946: 126 (distr.).

Arhopalus rusticus nubilus; Linsley, 1962: 75, fig. 27; Chemsak, 1967b: 182 (distr.); Turnbow & Franklin, 1980: 338 (distr.); Chemsak, Linsley & Noguera, 1992: 24 (cat.); Browne, Peck & Ivie, 1993: 41 (distr.); Chemsak, 1996: 25, pl. 2, fig. 1 (rev.); Linsley & Chemsak, 1997: 346

(hosts); Monné, M. A., 2002: 34 (cat. hosts); Monné, M.A., & Giesbert, 1994: 22 (checklist); Monné, M.A., 2006: 110 (cat.).

Arhopalus nubilus; Bousquet. Laplante, Hammond & Langor, 2017: 81

6. *Arhopalus obsoletus* (Randall, 1838)

Type locality - United States, Maine. (Depository unknown). **Distribution** - Boreal North America west to Rocky Mountains, Mexico, Guatemala, Honduras. **Host plants** - *Pinus banksiana* Lambert, *P. pseudo-strobus* Lindley, *P. sylvestris* Linné, *P. tenuifolia* Benth (Pinaceae).

Callidium obsoletum Randall, 1838: 27.

Criocephalus obsoletus; LeConte, 1851: 106; White, 1855: 311; Bland, 1861: 94 (distr.); Lacordaire, 1868: 211; Packard, 1872: 93 (distr.); LeConte, 1873a: 170; Provancher, 1877: 585; Leng, 1884: 63; Packard, 1888: 142 (distr.); Wickham, 1897b: 106; Chagnon, 1897: 15 (distr.); Smith, 1900: 285 (distr.); Ulke, 1903: 25 (distr.); Sharp, 1905: 154; Chagnon, 1905: 232; Fall & Cockerell, 1907: 191 (distr.); Blatchley, 1910: 1015; Smith, 1910: 325 (distr.); Fisher & Kirk, 1912: 309 (distr.); Nicolay, 1919: 63 (distr.); Britton, 1920: 266 (distr.); Craighead, 1923: 33 (larva, biol.); Hatch, 1925: 579 (distr.); Kirk & Knull, 1926: 22 (distr.); Leonard, 1928: 434 (distr.); Beaulne, 1932: 198 (hosts); Linsley, 1935a: 72 (distr.); Doane *et al.*, 1936: 168 (biol., hosts); Brimley, 1938: 210 (distr.); Loding, 1945: 114 (distr.); Craighead, 1950: 237 (biol.); Schwerdtfeger, 1955: 319 (biol.).

Arhopalus obsoletus; Knull, 1946: 150; Duffy, 1960: 82 (hosts).

Arhopalus rusticus obsoletus; Linsley, 1962: 75, fig. 27; Gosling, 1973: 68, fig. 8 (biol.); Turnbow & Franklin, 1980: 338 (distr.); Rice, 1981: 460 (distr.); Chemsak, Linsley & Noguera, 1992: 24 (cat.); MacRae, 1993: 227 (distr., hosts); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 25, pl. 1, fig. 12 (rev.); Linsley & Chemsak, 1997: 346 (hosts); Monné, M.A., 2002: 34 (cat. hosts); Turnbow, Cave & Thomas, 2003: 5 (distr.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Monné, M.A., 2006: 110 (cat.); Hovore, 2006: 371 (distr.); Holt, 2013: 246 (distr.); Spomer, 2014: 298 (distr.); Webster *et al.*, 2016: 112 (hosts); Klingeman *et al.*, 2017: 295 (distr.)

Arhopalus obsoletus; Webster, 2016: 486; Bousquet. Laplante, Hammond & Langor, 2017: 81, pl. 14

Criocephalus obscurus LeConte, 1850a: 36.

Type locality - United States, New York. (MCZN).

Criocephalus mexicanus Thomson, 1861a: 260; 1878: 7 (type); Bates, 1879: 15 (distr.); 1884: 239 (distr.); Schaeffer, 1908: 329 (distr.).

Type locality - Mexico. (MNHN).

7. *Arhopalus productus* (LeConte, 1850)

Type locality - Holotype: United States, Oregon. (MCZN). **Distribution** - Pacific coast of North America from British Columbia to Baja California and Rocky Mountains, south into Mexico, Guatemala and Honduras. **Host plants** - *Abies* sp., *Pinus* sp., *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae).

Criocephalus productus LeConte, 1850a: 36; White, 1855: 310; LeConte, 1857: 23 (distr.); 1859a: 48 (distr.); 1869: 371 (distr.); 1873a: 169; Horn *in* Hayden, 1873: 717; Packard, 1877: 804; Snow, 1877: 19 (distr.); LeConte, 1879: 505 (distr.); Packard, 1881: 159, fig. 71 (biol.); Snow, 1881: 70 (distr.); 1883: 42 (distr.); Leng, 1884: 63, 64 (not pl. 2, fig. 4); Bates, 1884: 239 (distr.); Packard, 1890: 699, fig. 235 (biol.); Townsend, 1895: 47 (distr.); Harrington, 1899: 108 (distr.); Fall, 1901: 27; Sharp, 1905: 160; Snow, 1906: 1906b: 170 (distr.); Fall & Cockerell, 1907: 191 (distr.); Wickham, 1909a: 28 (distr.); Hyslop, 1912: 100 (biol.); Garnett, 1918: 173 (distr.); Craighead, 1923: 32, pl. 2, figs 2, 3, 6, 23, 25 (larva); Hardy, 1926: 4, pl. 4, fig. 23; Knowlton, 1930: 56 (distr.); Pack, 1930: 219 (distr.); Beaulne, 1932: 198 (hosts); Mank, 1934: 80 (distr.); Knowlton & Thatcher, 1936: 278 (distr.); Doane *et al.*, 1936: 168 (biol.); Moore, 1937: 87 (distr.); Kimmey & Furniss, 1943: 23, figs 6, c-d (biol.).

Arhopalus productus; Knowlton & Wood, 1950: 10 (distr.); Keen, 1952: 195 (biol.); Eaton & Lyon, 1955: 11, figs 1-3 (biol.); Linsley, 1957: 12 (syn.); Eaton, 1959: 114 (biol.); Linsley,

1962: 72, fig. 27; Hatch, 1971: 95, pl. 16, fig. 1; Kirk & Balsbaugh, 1975: 97 (distr.); Furniss & Carolin, 1977: 291, fig. 174A (hosts); Chemsak, Linsley & Mankins, 1980: 29 (distr.); Chemsak, Linsley & Noguera, 1992: 24 (cat.); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 22, pl. 1, fig. 11 (rev.); Linsley & Chemsak, 1997: 345 (hosts); Heffern 1998: 7 (distr.); Monné, M.A., 2002: 33 (cat. hosts); Turnbow, Cave & Thomas, 2003: 5 (distr.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Monné, M.A., 2006: 109 (cat.); Hovore, 2006: 371 (distr.); Monné, M.A. & Hovore, 2006: 24 (checklist); Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 81, pl. 14
Criocephalus elongatus Casey, 1924: 230; Lingafelter *et al.*, 2014: 57, figs 60w, x (lect. designation).

Type locality - Lectotype female: United States, Utah: Stockton. (USNM).

Criocephalus propinquus Casey, 1924: 230; Lingafelter *et al.*, 2014: 303, figs 136g, h (lect. designation).

Type locality - Lectotype female: United States, Colorado. (USNM).

8. *Arhopalus rusticus rusticus* (Linnaeus, 1758)

Syntypes locality - Syntypes: Suecia. (UZIU). **Distribution** - Europe, Palearctic Asia through Japan; north of Africa, and North America (except Pacific Coast). Introduced in Argentina (Buenos Aires).

Cerambyx rusticus Linnaeus, 1758: 395.

Arhopalus rusticus; Laporte, 1840: 453; Linsley, 1962: 63; Monné, M.A., & Giesbert, 1994: 22 (checklist); Chemsak, 1996: 23; Di Iorio, 2004d: 664 (distr.); Turienzo, 2007: 321 (hosts, distr.); Monné, M.L., Monné, M.A. & Wang, 2017: 35, fig. 58 (larva); Bousquet, Laplante, Hammond & Langor, 2017: 81, pl. 14

Asemum Eschscholtz, 1830

Asemum Eschscholtz, 1830: 66; Audinet-Serville, 1834b: 79; Westwood, 1838: 40; Mulsant, 1839: 62; LeConte, 1850a: 35; Thomson, 1861a: 260; Mulsant, 1863: 119, 425; Thomson, 1864: 268; 1865: 464; Lacordaire, 1868: 206; LeConte, 1873b: 293; LeConte & Horn, 1883: 277, 278; Leng, 1884: 62; Blatchley, 1910: 1014; Casey, 1912: 257; Craighead, 1923: 31 (larva); Bradley, 1930: 230; Hopping, 1931: 234 (syn.); Chagnon, 1936: 256; Knull, 1946: 184; Dillon & Dillon, 1961: 582; Linsley, 1962: 78; Arnett, 1962: 856; Chagnon & Robert, 1962: 253, 256; Hatch, 1971: 95; Monné, M.A., 2006: 110 (cat.); Monné, M.A., 2012: 133; Bousquet, Laplante, Hammond & Langor, 2017: 82 (key spp.)

Type-species - *Cerambyx striatus* Linnaeus, 1758 (subsequent designation, Westwood, 1838).

Onychoplectes Gistel, 1856: 376.

Type-species - *Cerambyx striatus* Linnaeus, 1758 (monotypy).

Liaseum Casey, 1912: 262; Bradley, 1930: 230.

Type-species - *Asemum nitidum* LeConte, 1873 (subsequent designation, Linsley, 1962: 78).

1. *Asemum australe* LeConte, 1850

Type locality - Holotype: United States, Georgia. (MCZN). **Distribution** - Canada, Atlantic Coast from Massachusetts to Georgia and Louisiana. **Host plants** - *Pinus strobus* Linné (Pinaceae).

Asemum australe LeConte, 1850a: 35; White, 1855: 312; Linsley, 1957: 14 (syn.); 1962: 83, fig. 31; Turnbow & Franklin, 1980: 338; Chemsak, Linsley & Noguera, 1992: 24 (cat.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Chemsak, 1996: 36, pl. 2, fig. 8; Yanega, 1996: 44, pl. 2, fig. 18; Schiefer, 1998: 116 (distr.); Vlasák & Vlasakova, 2002: 209 (distr.); Monné, M.A. & Hovore, 2006: 24 (checklist); Lingafelter, 2007: 15, 139 (hosts); Bousquet, Laplante, Hammond & Langor, 2017: 82, pl. 15

Criocephalus australis; LeConte, 1862: 43; 1873a: 170; 1873b: 294; Henshaw, 1874: 22 (distr.); LeConte & Horn, 1883: 179; Leng, 1884: 63; Evans, 1906: 99 (distr.); Wickham, 1909a: 28 (distr.).

Criocephalus champlainianus Casey, 1924: 231; Leonard, 1928: 434; Lingafelter *et al.*, 2014: 39, fig. 41g (holotype).

Type locality - Holotype male: United States, New York: near Lake George (Lake Champlain region). (USNM).

2. *Asemum caseyi* Linsley, 1957

Type locality - Holotype male: United States, California: Bear Valley (St. Bernardino Mts.). (CASC). **Distribution** - Canada. Coniferous forests of Oregon and California to Mexico (Baja California). **Host plants** - *Pinus contorta* Douglas ex Loudon, *P. coulteri* D. Don, *P. jeffreyi* Grev, & Balb., *P. lambertiana* Douglas, *P. ponderosa* Douglas ex Lawson & P. Lawson, *P. sabiniana* Douglas ex D. Don (Pinaceae).

Asemum caseyi Linsley, 1957: 13; 1962: 83, fig. 31; Barr & Penrose, 1969: 88 (distr.); Hatch, 1971: 96; Furniss & Carolin, 1977: 292; Skiles, 1978b: 14 (biol.); Hovore, 1988: 4 (distr.); Chemsak, Linsley & Noguera, 1992: 24 (cat.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 34, pl. 2, fig. 7 (rev.); Linsley & Chemsak, 1997: 346 (hosts); Monné, M.A., 2002: 35 (cat. hosts); Monné, M.A., 2006: 111 (cat.); Monné, M.A. & Hovore, 2006: 24 (checklist); Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 83, pl. 15

3. *Asemum nitidum* LeConte, 1873

Type locality - Holotype male: United States, Oregon. (MCZN). **Distribution** - Pacific Coast of North America from British Columbia to southern California and New Mexico.

Host plants - *Abies* sp., *Pinus jeffreyi* Balfour, *P. ponderosa* Douglas ex Lawson & P. Lawson, *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae).

Asemum nitidum LeConte. 1873a: 169; Leng, 1884: 62; Fall, 1901: 143 (distr.); Hopkins, 1904: 21 (biol.); Garnett, 1918: 173 (distr.); Craighead, 1923: 32, pls 2,3,7; Hopping, 1931: 235; Beaulne, 1932: 197 (hosts); Doane *et al.*, 1936: 167; Keen, 1952: 196 (biol.); Linsley, 1957: 13 (syn.); Townes & Townes, 1960: 157 (paras.); Linsley, 1962: 82; Barr & Penrose, 1969: 88 (distr.); Furniss & Carolin, 1977: 292; Skiles, 1978b: 14 (distr.); Chemsak, Linsley & Noguera, 1992: 24 (cat.); Monné, M.A., 1994: 22 (cat.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Chemsak, 1996: 33, pl. 2, fig. 6; Linsley & Chemsak, 1997: 346 (hosts); Monné, M.A. & Hovore, 2006: 24 (checklist); Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 83

Liasemum nitidum; Casey, 1912: 262; Essig, 1926: 450; Moore, 1937: 87 (distr.);

Liasemum mokelumne Casey, 1912: 262; Moore, 1937: 87 (distr.); Lange, 1937: 174 (hosts); Lingafelter *et al.*, 2014: 101, 368, fig. 111g (lectotype).

Asemum mokelumne; Hopping, 1931: 235; Doane *et al.*, 1936: 167; Keen, 1952: 196 (hosts);

Type locality - Lectotype female: United States, California: Calaveras County, Mokelumne Hill. (USNM).

4. *Asemum striatum* (Linnaeus, 1758)

Syntypes locality - Syntypes: Sweden. (UZIU). **Distribution** - Coniferous belt of Europe, Asia and North America. **Host plants** - *Abies firma* Siebold & Zuccarini, *A. holophylla* Maximowicz, *A. homolepis* Siebold & Zuccarini, *A. mariesii* Masters, *A. nephrolepis* (Trautvetter) Maximowicz, *A. sachalinensis* Masters, *A. veitchii* Lindley, *Larix europaea* de Candolle, *L. kaempferi* Fortune ex Gordon, *L. koreana* Nakai, *L. laricina* (Duroi) K. Koch, *Picea abies* Linné, *P. glauca* (Moench) Voss, *P. jezoensis* Carrière, *P. koraiensis* Nakai, *Pinus armandii* Franchet, *P. contorta* Douglas ex Loudon, *P. densiflora* Siebold & Zuccarini, *P. echinata* Miller, *P. inops* Solander, *P. jeffreyi* Balfour, *P. koraiensis* Siebold & Zuccarini, *P. nigra* Arnold, *P. ponderosa* Douglas ex Lawson & P. Lawson, *P. resinosa* Aiton, *P. rigida* Miller, *P. strobus* Linné, *P. sylvestris* Linné, *P. tabulaeformis* Hortulanorum ex C. Koch, *P. thunbergii* Parlatore, *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae), *Cryptomeria japonica* D. Don (Taxodiaceae).

Cerambyx striatus Linnaeus, 1758: 396; 1761: 192; 1767: 635.

Callidium striatum; Fabricius, 1775: 191; 1801: 343; Kirby, 1837: 171; Bethune, 1872: 56.

Asemum striatum; Audinet-Serville, 1834: 69; White, 1855: 312; Thomson, 1864: 268; Heyden, 1890: 132 (syn.); Smith, 1905: 165 (hosts); Dillon & Dillon, 1961: 582, pl. 57, No. 9; Linsley, 1962: 79, figs 29, 30; Hatch, 1971: 97, pl. 9, fig. 6; Swan & Papp, 1972: 444, fig. 934; Gosling, 1973: 68 (distr.); Perry, 1975: 59 (hosts); Kirk & Balsbaugh, 1975: 97 (distr.); Furniss & Carolin, 1977: 292 (hosts); Turnbow & Franklin, 1980: 338 (distr.); Gosling, 1984: 71 (hosts); Waters & Hyche, 1984: 284 (distr.); Gosling, 1986: 255 (hosts); Chemsak, Linsley & Noguera, 1992: 24 (cat.); MacRae, 1993: 227 (distr., hosts); Monné, M.A., 1994: 23 (cat.); Monné, M.A., & Giesbert, 1994: 22 (checklist); Chemsak, 1996: 32, pl. 2, fig. 5; Yanega, 1996: 94, pl. 2, figs 17a,b; Linsley & Chemsak, 1997: 345 (hosts); Heffern, 1998: 7 (distr.); Schiefer, 1998: 116 (distr.); Peck & Thomas, 1998: 117 (distr.); Vlasák & Vlasakova, 2002: 209 (distr., hosts); Turnbow, Cave & Thomas, 2003: 5 (distr.); Sikes & Webster, 2005: 327 (distr.); Monné, M.A. & Hovore, 2006: 24 (checklist); Lingafelter, 2007: 15, 139, 2 figs, pl. 3c (hosts); Webster, McCorquodale & Majka, 2009: 221 (distr.); Holt, 2013: 246 (distr.); Webster, 2016: 486; Monné, M.L., Monné, M.A. & Wang, 2017, fig. 97; Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 82, pl. 15

Callidium agreste Fabricius, 1787: 152; 1801: 338; Zimsen, 1964: 179 (type).

Type locality - Holotype: Germany, Saxe-Anhalt: Halle. (ZMUC).

Callidium (Callidium) dichrous Gmelin, 1790: 1846.

Type locality -

Callidium buprestoide Savenius, 1825: 64.

Type locality - Holotype: Finlandia. (Depository unknown).

Asemum atrum Eschscholtz, 1830: 66; LeConte, 1851: 105; White, 1855: 312; LeConte, 1857: 23; LeConte, 1869: 371; Snow, 1877: 19 (distr.); Harrington, 1881: 22 (biol.); Leng, 1884: 62; Holland, 1888: 91 (distr.); Harrington, 1890: 185 (distr.); Packard, 1890: 809 (hosts); Keen, 1895: 219 (distr.); Beutenmuller, 1896: 74 (hosts); Wickham, 1897b: 105; Chagnon, 1897: 15 (distr.); Harrington, 1899: 107 (distr.); Warren, 1899: 196 (distr.); Ouellet, 1902: 122 (distr.); Fall & Cockerell, 1907: 191 (distr.); Casey, 1912: 259; Garnett, 1918: 173 (distr.); Craighead, 1920: 8; 1923: 32, pl. 6, fig. 1 (larva); Fall, 1926: 201 (distr.); Essig, 1926: 450; Engelhardt, 1927: 73 (distr.); Knowlton, 1930: 56; Pack, 1930: 219 (distr.); Hopping, 1931: 235; Beaulne, 1932: 197 (hosts); DeLeon, 1934: 57 (hosts); Brown, 1934: 229 (distr.); Mank, 1934: 79 (distr.); Knowlton & Thatcher, 1936: 278 (distr.); Chagnon, 1936: 256; Doane *et al.*, 1936: 167, fig. 81; Lange, 1937: 174 (hosts); Kimmey & Furniss, 1943: 26; Knull, 1946: 148, pl. 4, fig. 11; Leech, 1947: 108 (hosts); Knowlton & Wood, 1950: 10 (distr.); Howden & Vogt, 1951: 591 (hosts); Beal, Haliburton & Knight, 1951: 121 (biol.); Keen, 1952: 195 (biol.); Thomas, 1955: 340 (hosts); Hicks, 1955: 165 (distr.); Clark, 1956: 41 (distr.); Gardiner, 1957: 249; Linsley, Knull & Statham, 1961: 9 (distr.); Chagnon & Robert, 1962: 256; Laliberté, Chantal & LaRochelle, 1977: 91.

Type locality - Holotype: United States, California. (ZMUM)

Asemum moestum Haldeman, 1847a: 35; LeConte, 1850a: 35; Mannerheim, 1853: 248; White, 1855: 312; LeConte, 1857: 23; Bland, 1861: 94; Pettit, 1871: 105; Packard, 1872: 496, fig. 483; Henshaw, 1874: 22; Horn, 1876: 169; LeConte, 1879: 505; Riley, 1880: 238 (hosts); Harrington, 1881: 33 (biol.); Packard, 1881: 156, fig. 70 (biol.); Leng, 1884: 62, pl. 2, fig. 1; Knaus, 1885: 69 (distr.); Packard, 1890: 697, 829, pl. 19, fig. 1, fig. 234; Hopkins, 1893: 192 (biol.); Hamilton, 1894: 30; 1895: 337; Slosson, 1894: 3 (distr.); Evans, 1895: 173 (distr.); Beutenmuller, 1896: 74 (hosts); Wickham, 1897b: 105; 1899: 109 (distr.); Smith, 1900: 285 (distr.); Dury, 1902: 158 (distr.); Ouellet, 1902: 122 (distr.); MacGillivray & Houghton, 1902: 252 (distr.); Knaus, 1903: 176 (distr.); Ulke, 1903: 25, 50 (distr.); Hopkins, 1904: 35 (biol.); Chagnon, 1905: 35 (distr.); 1905: 232; Houghton, 1905: 51 (distr.); Fall & Cockerell, 1907: 191 (distr.); Wickham, 1909a: 28 (distr.); Smith, 1910: 324; Frost, 1912: 306 (distr.); Casey, 1912: 257; Nicolay, 1917: 94 (distr.); Chagnon, 1917: 230 (distr.); Blackman & Stage, 1918: 6 (biol.); Nicolay, 1919: 63 (distr.); Britton, 1920: 266 (distr.); Craighead, 1923: 31, pl. 6, figs 8, 12, pl. 25, figs 1,2; Hatch, 1925: 579 (distr.); Kirk & Knull, 1926: 21 (distr.); Procter, 1927: 110 (distr.); Leonard, 1928: 434 (distr.); Beaulne, 1932: 197 (hosts); Wolcott & Montgomery, 1933: 155; Easterling, 1934: 140 (hosts); Brown, 1934: 229 (distr.); Herrick, 1935: 252, 377 (biol.); Doane *et al.*, 1936: 167; Brimley, 1938: 210 (distr.); Savely, 1939:

- 333 (biol.); Parmelee, 1941: 377 (hosts); Loding, 1945: 114 (distr.); Procter, 1946: 176 (distr.); Hardy, 1948: 32 (distr.); Craighead, 1950: 237; Gardiner, 1950: 1.
- Syntypes locality** - Syntypes: United States, Pennsylvania. (MCZN).
- Asemum moestum* var. *obsoletum* Haldeman, 1847a: 35.
- Type locality** - Holotype: United States. (MCZN).
- Asemum moestum* var. *brunneum* Haldeman, 1847a: 35.
- Type locality** - Holotype: United States. (MCZN).
- Asemum substriatum* Haldeman, 1847a: 36.
- Type locality** - Holotype: United States. (MCZN).
- Asemum juvenicum* Haldeman, 1847a: 36; Casey, 1912: 257.
- Type locality** - Holotype: United States. (MCZN).
- Asemum fuscum* Haldeman, 1847a: 36; Casey, 1912: 261.
- Type locality** - United States. (MCZN).
- Asemum subsulcatum* Motschulsky, 1860: 152.
- Type locality** - Lectotype male: Russia, Siberia. (ZMUM).
- Asemum amurense* Kraatz, 1879: 97.
- Syntypes locality** - Syntypes: Russia, Ost Siberien: Amur.(SDEI).
- Asemum gracilicorne* Casey, 1912: 258; Lingafelter et al., 2014: 70, fig. 75q (holotype).
- Type locality** - Holotype male: United States, Wisconsin: Bayfield. (USNM).
- Asemum ebenum* Casey, 1912: 258; Leonard, 1928: 434 (distr.); Lingafelter et al., 2014: 56, fig. 59s (holotype).
- Type locality** - Holotype female: United States, Lake Superior. (USNM).
- Asemum curtipenne* Casey, 1912: 258; Lingafelter et al., 2014: 49, fig. 52a (holotype).
- Type locality** - Holotype female: United States, Wisconsin: Bayfield. (USNM).
- Asemum amputatum* Casey, 1912: 259; Lingafelter et al., 2014: 15, fig. 15c (holotype).
- Type locality** - Holotype male: United States, Massachusetts: Wellesley. (USNM).
- Asemum parvicorne* Casey, 1912: 260; Lingafelter et al., 2014: 296, fig. 128 m (holotype).
- Type locality** - Holotype female: United States, Colorado: Boulder County. (USNM).
- Asemum fulvipenne* Casey, 1912: 260; Knowlton & Thatcher, 1936: 278 (distr.); Lingafelter et al., 2014: 77, fig. 72c (holotype).
- Asemum atrum fulvipenne*; Casey, 1924: 228;.
- Type locality** - Holotype female: United States, Wyoming: Laramie. (USNM).
- Asemum costulatum* Casey, 1912: 260; Lingafelter et al., 2014: 47, fig. 49q (holotype).
- Type locality** - Holotype female: Canada, British Columbia: Aldermere. (USNM).
- Asemum pugetanum* Casey, 1912: 261; Lingafelter et al., 2014: 305, 365, fig. 138 g (lectotype).
- Type locality** - Lectotype male: United States, Washington: Seattle. (USNM).
- Asemum brevicorne* Casey, 1912: 261; Lingafelter et al., 2014: 32, fig. 33e (holotype).
- Type locality** - Holotype male: Canada, Ontario. (USNM).
- Asemum amurense* var. *tomentosum* Plavilstshikov, 1915: 108.
- Type locality** - Lectotype male: China, Manchuria, Prov. Girin: Chandaohedzy. (ZMUM).
- Asemum striatum* var. *limbatipenne* Pic, 1916: 10.
- Type locality** - Holotype: France, Saône-et-Loire: Marly-sous-Issy.(MNHN).
- Asemum carolinum* Casey, 1924: 227; Brimley, 1938: 210 (distr.); Lingafelter et al., 2014: 37, 365, fig. 38u (lectotype).
- Type locality** - Lectotype male: United States, North Carolina: Southern Pines. (USNM).
- Asemum stocktonense* Casey, 1924: 227; Lingafelter et al., 2014: 324, fig. 160 c (holotype).
- Type locality** - Holotype male: United States, Utah: Stockton. (USNM).
- Asemum striatum* var. *Theresae* Pic, 1945: 6
- Type locality** - Holotype: France, Saône-et-Loire: Digoïn. (MNHN).
- Megasemum Kraatz, 1879**
- Megasemum* Kraatz, 1879: 97; Sharp, 1905: 147; Van Dyke, 1937: 112; Linsley, 1962: 76; Arnett, 1962: 856; Hatch, 1971: 94; Monné, M.A., 2006: 111 (cat.); Monné, M.A. & Hovore, 2006: 25 (checklist); Monné, M.A., 2012: 133.
- Type-species** - *Megasemum quadricostulatum* Kraatz, 1879 (monotypy)

1. *Megasemum asperum* (LeConte, 1854)

Syntypes locality - Syntypes: United States, Oregon: Prairie Paso. (MCZN). **Distribution** - Coniferous belt of western North America to southern montane Arizona and northern Mexico (Sonora). **Host plants** - *Abies* sp., *Pinus contorta* Douglas ex Loudon, *P. monticola* Douglas ex D. Don, *Pseudotsuga menziesii* (Mirbel) Franco (Pinaceae).

Asemum asperum LeConte, 1854a: 18; 1857: 60; 1862: 42; 1869: 371.

Nothorhina aspera; LeConte, 1873a: 169; 1873b: 294; LeConte & Horn, 1883: 279; Leng, 1884: 63; 1886: pl. 3, fig. 24; Smith, 1888: 91; Harrington, 1899: 108 (distr.); Knaus, 1903: 176 (distr.); Snow, 1906b: 170 (distr.); Fall & Cockerell, 1907: 191 (distr.); Casey, 1912: 263; Garnett, 1918: 173 (distr.); Craighead, 1923: 33, pl. 6, fig. 10 (larva); Hardy & Preece, 1927b: 61 (biol.); Beaulne, 1932: 198 (hosts); Mank, 1934: 79 (distr.).

Megasemum asperum; DeLeon, 1934: 57 (hosts); Doane *et al.*, 1936: 167 (biol.); Van Dyke, 1937: 112; Hardy, 1942: 10 (biol.); Keen, 1952: 196 (hosts); Linsley, 1957: 12 (syn.); Linsley, Knull & Statham, 1961: 9 (distr.); Linsley, 1962: 77, fig. 28; Ross, 1968: 11, 12 (biol.); Hatch, 1971: 96; Furniss & Carolin, 1977: 293; Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 29, pl. 2, fig. 4 (rev.); Linsley & Chemsak, 1997: 396 (hosts); Heffern, 1998: 7 (distr.); Monné, M.A., 2002: 35 (cat. hosts); Monné, M.A., 2006: 111 (cat.); Monné, M.A. & Hovore, 2006: 25 (checklist); Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 82, pl. 15

Nothorhina aspera impressa Casey, 1912: 264; Lingafelter *et al.*, 2014: 20, figs 20o, p (lect. designation).

Type locality - Lectotype male: United States, Colorado: Boulder Co. (USNM).

Nothorhina aspera rectipennis Casey, 1912: 264; Lingafelter *et al.*, 2014: 20, figs 20q, r (holotype).

Type locality - Holotype male: United States, New Mexico: Cloudcroft. (USNM).

Criocephalus uteanus Casey, 1924: 231; Lingafelter *et al.*, 2014: 339, figs 177q, r (holotype).

Type locality - Holotype female: United States, Utah: Eureka. (USNM).

Tetropium Kirby, 1837

Callidium (*Tetropium*) Kirby in Richardson, 1837: 174;

Tetropium; Haldeman, 1847b: 372; LeConte, 1850a: 34; 1850b: 35; Thomson, 1861a: 258; Kraatz, 1863: 106; Thomson, 1864: 266; 1865: 462; Lacordaire, 1868: 208; Bethune, 1872: 93; LeConte, 1873b: 293, 294; Provancher, 1877: 578, 585; LeConte & Horn, 1883: 277, 278; Leng, 1884: 95; Casey, 1891: 22; 1896: 165; Wickham, 1897b: 105; Gahan, 1906: 95; Casey, 1912: 268; Craighead, 1923: 33 (larva); Bradley, 1930: 230; Chagnon, 1936: 256; Knull, 1946: 248; Linsley, 1962: 85; Arnett, 1962: 856; Chagnon & Robert, 1962: 253, 256; Hatch, 1971: 95; Mroczkowski, 1986: 188; Svacha & Danilevsky, 1987: 153 (larva); Monné, M.A., 2006: 111 (cat.); Monné, M.A. & Hovore, 2006: 24 (checklist); Monné, M.A., 2012: 133; Bousquet, Laplante, Hammond & Langor, 2017: 83 (key spp.)

Type-species - *Tetropium cinnamopterum* Kirby, 1837 (subsequent designation, Thomson, 1864: 266).

Isarthron Dejean, 1835: 229.

Type-species - *Callidium aulicum* Fabricius, 1775 (Linsley designation, 1962: 85).

Criomorphus Mulsant, 1839: 58; 1863: 115, 421; Planet, 1924: 236; Portevin, 1927: 26.

Type-species - *Callidium aulicum* Fabricius, 1775 (monotypy).

1. *Tetropium abietis* Fall, 1912

Syntypes locality - Syntypes male and female: United States, California: Fresno County, Huckleberry Meadow. (MCZN), **Distribution** - Pacific Coast from Washington south to southern California. **Host plants** - *Juniperus* sp. (Cupressaceae), *Abies concolor* Gordon & Glen, *A. magnifica* A. Murray (Pinaceae).

Tetropium abietis Fall, 1912: 320; Craighead, 1923: 34, pl. 3, fig. 8, pl. 8, fig. 7 (larva); Essig, 1926: 450 (biol.); Craighead & Middleton, 1930: 9 (biol.); Beaulne, 1932: 198 (hosts); Doane

et al., 1936: 168 (biol.); Hatch, 1938: 187; Keen, 1952: 173 (biol.); Anderson, 1960: 255 (biol.); Linsley, 1962: 87, figs 32,33; Tyson, 1966: 207 (hosts); Ruetter, 1970: 22 (paratype); Hatch, 1971: 97, pl. 9, fig. 7; Swan & Papp, 1972: 444, fig. 933; Furniss & Carolin, 1977: 293 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 25 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 39, pl. 2, fig. 9; Linsley & Chemsak, 1997: 443 (hosts); Monné, M.A. & Hovore, 2006: 24 (checklist); Bousquet, Laplante, Hammond & Langor, 2017: 84, pl. 15

2. *Tetropium auripilis* Bates, 1885

Type locality - Holotype male: Mexico, Veracruz: Las Vigas. (MNHN). **Distribution** - United States (southern Arizona), Mexico (Chihuahua, Veracruz)

Tetropium auripilis Bates, 1885: 435; Linsley, 1934c: 162; Linsley, Knull & Statham, 1961: 9 (distr.); Linsley, 1962: 91, fig. 33; Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Monné, M.A., 2006: 112 (cat.); Monné, M.A. & Hovore, 2006: 24 (checklist); Gutiérrez & Noguera, 2015: 138 (distr.).

3. *Tetropium cinnamopterum* Kirby, 1837

Syntypes locality - Syntypes: America Borealis, Latitude 65 degrees. (BMNH).

Distribution - Northeastern United States to Alaska. **Host plants** - *Abies* sp., *Picea glauca* (Moench) Voss, *Pinus inops* Solander, *P. ponderosa* Douglas ex Lawson & P. Lawson (Pinaceae).

Callidium (Tetropium) cinnamopterum Kirby, 1837: 174, pl. 5, fig. 8.

Callidium cinnamopterum; Haldeman, 1847a: 37.

Tetropium cinnamopterum; LeConte, 1850b: 234; 1850a: 35; White, 1855: 327; LeConte, 1857: 6, 23 (distr.); Thomson, 1864: 266; Bethune, 1872: 94 (distr.); Horn, 1876: 169; Provancher, 1877: 586; Snow, 1877: 19 (distr.); Leng, 1884: 96; Snow, 1885: 68 (distr.); Leng, 1885: 35, pl. 1, fig. 1; Harrington, 1890: 185 (distr.); Casey, 1891: 22; Slosson, 1893: 290 (distr.); 1894: 3 (distr.); Hopkins, 1893: 192; Hamilton, 1895: 337 (distr.); 1896: 165; Evans, 1895: 173 (distr.); Wickham, 1897b: 105; MacGillivray & Houghton, 1902: 252 (distr.); Skinner, 1903: 40 (distr.); Hopkins, 1904: 27 (biol.); Houghton, 1905: 51 (distr.); Snow, 1906: 179 (distr.); Fall & Cockerell, 1907: 191 (distr.); Davis, 1909: 42 (distr.); Blatchley, 1910: 1015; Morris, 1916: 19 (hosts); Chagnon, 1917: 230 (distr.); Gibson, 1917: 150 (distr.); Britton, 1920: 266 (distr.); Rohwer, 1921: 442 (paras.); Craighead, 1923: 34, pl. 2, figs 1-2, pl. 6, fig. 9 (larva); Fall, 1926: 201 (distr.); Kirk & Knull, 1926: 21 (distr.); Leonard, 1928: 434 (distr.); Beaulne, 1932: 198 (hosts); Mank, 1934: 79 (distr.); Chagnon, 1936: 256; Doane *et al.*, 1936: 168; Chagnon, 1939: 86 (distr.); Morley, 1939: 244 (biol.); Townes, 1944: 772 (paras.); Knull, 1946: 149; Procter, 1946: 177 (distr.); Craighead, 1950: 268 (biol.); Howden & Vogt, 1951: 591 (hosts); Belyea, 1952: 325 (hosts); Duffy, 1953: 259, fig. 121 (larva); Gardiner, 1957: 249, figs 1, 10, 15; Townes & Townes, 1960: 109, 159, 375 (paras.); Linsley, 1962: 89; Roff, 1967: 27 (hosts); Ross, 1968: 12 (biol.); Ross & Vanderwal, 1969: 10, figs. 1-5; Baker, 1972: 199; Swan & Papp, 1972: 444; Gosling, 1973: 68, fig. 10 (biol.); Raske, 1973: 745, figs 1-6; Furniss & Carolin, 1977: 293 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 25 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 42; Yanega, 1996: 44, pl. 2, fig. 21; Androw & Keeney, 1999: 2 (distr.); Vlasák & Vlasakova, 2002: 209 (distr., hosts); Webster, McCorquodale & Majka, 2009: 291 (distr.); Costello, Jacobi & Negrón, 2013: 151 (hosts); Webster, 2016: 486; Goczal, Sweeney & Tofilskim 2017: 497, fig. 3 (morph.); Bousquet, Laplante, Hammond & Langor, 2017: 85, pl. 15

Tetropium cinnamopterum cinnamopterum; Chemsak, 1996: 43, pl. 2, fig. 11; Linsley & Chemsak, 1997: 443 (hosts); Monné, M.A. & Hovore, 2006: 24 (checklist);

Criomorpha cinnamopterus; Mannerheim, 1853: 247.

4. *Tetropium fuscum* (Fabricius, 1787)

Syntypes locality - Syntypes: Germany, Saxe-Anhalt: Halle. (ZMUC). **Distribution** - Europe, introduced in Canada. **Host plants** - *Abies* sp., *Picea abies* Linné, *P. excelsa* Link,

P. glauca (Moench) Voss, *P. mariana* Britton, Sterns & Poggenburg, *P. orientalis* Carrière, *P. rubens* Sargent, *Pinus sylvestris* Linné (Pinaceae).

Callidium fuscum Fabricius, 1787: 154.

Tetropium fuscum; Smith & Hurley, 2001: 540 (distr., hosts); Silk *et al.*, 2007: 697 (pherom); Haack, Keena & Eyre, 2017: 75; Webster, 2016: 486; Haack, 2017: 120 (hosts); Hanks & Wang, 2017: 134 (biol.); Bousquet, Laplante, Hammond & Langor, 2017: 85

5. *Tetropium parallelum* Casey, 1891

Type locality - Lectotype female: United States, Colorado. (USNBM). **Distribution** - Canada, Colorado to Arizona and New Mexico. **Host plants** - *Abies concolor* Gordon & Glen (Pinaceae).

Tetropium parallelum Casey, 1891: 23; Leng & Hamilton, 1896: 165; Linsley, Knull & Statham, 1961: 9 (distr.); Linsley, 1962: 91; Raske, 1973: 754; Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 26 (cat.); Chemsak, 1996: 45, -pl. 3, fig. 2; Heffern, 1998: 8 (distr.); Monné, M.A. & Hovore, 2006: 24 (checklist); Lingafelter *et al.*, 2014: 295, 370, fig. 127 m (lectotype); Bousquet, Laplante, Hammond & Langor, 2017: 85

6. *Tetropium parvulum* Casey, 1801

Type locality - Holotype male: United States, Indiana. (USNM). **Distribution** - North Central States to Alaska, and Colorado. **Host plants** - *Picea engelmanni* Engelmann, *P. glauca* (Moench) Voss, *Pinus resinosa* Aiton, *P. strobus* Linné (Pinaceae).

Tetropium parvulum Casey, 1891: 24; Leng & Hamilton, 1896: 165; Raske, 1973: 745 (biol.); 1973: 757; Furniss & Carolin, 1977: 294 (hosts); Yanega, 1996: 44, pl. 2, fig. 22; Lingafelter *et al.*, 2014: 296, fig. 128q (holotype).

Tetropium cinnamopterum parvulum; Linsley, 1962: 89, fig. 33; Hatch, 1971: 98; Gosling, 1976: 158 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 25 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 44, pl. 2, fig. 12; Linsley & Chemsak, 1997: 443 (hosts); Heffern, 1998: 7 (distr.); Monné, M.A. & Hovore, 2006: 24 (checklist);

Tetropium parvulum; Bousquet, Laplante, Hammond & Langor, 2017: 85

Tetropium alaskanum Fall, 1926: 201; Raske, 1973: 754.

Type locality - Holotype male: United States, Alaska: Sheep Creek on the Yukon River below Eagle. (MCZN).

7. *Tetropium schwarzianum* Casey, 1891

Type locality - Holotype female: United States, Michigan: Marquette. (USNM). **Distribution** - Minnesota to Nova Scotia to North Carolina. **Host plants** - *Picea* sp., *Pinus resinosa* Aiton, *P. strobus* Linné (Pinaceae).

Tetropium schwarzianum Casey, 1891: 24; Leng & Hamilton, 1896: 165; Linsley, 1962: 89; Raske, 1973: 754; Gosling, 1973: 68 (distr.); 1986: 158 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 26 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 94, pl. 3, fig. 1; Yanega, 1996: 44, pl. 2, fig. 19; Linsley & Chemsak, 1997: 443 (hosts); Androw & Keeney, 1999: 2 (distr.); Vlasák & Vlasakova, 2002: 209 (distr.); Morris, 2002: 209 (distr., hosts); Monné, M.A. & Hovore, 2006: 25 (checklist); Lingafelter, 2007: 15, 139, 2 figs, pl. 3h (hosts); Webster, McCorquodale & Majka, 2009: 292 (distr.); Lingafelter *et al.*, 2014: 316, fig. 150q (holotype); Webster, 2016: 486; Klingeman *et al.*, 2017: 294 (distr.); DiGirolamo & Dodds, 2017: 410 (hosts); Bousquet, Laplante, Hammond & Langor, 2017: 85

7. *Tetropium velutinum* LeConte, 1869

Syntypes localities - Syntypes female: Canada, British Columbia: Vancouver's Island. United States, Oregon. (MCZN). **Distribution** - Pacific Coast and northern Rocky Mountains, to southern Arizona. **Host plants** - *Abies* sp., *Larix laricina* (Duroi) K. Koch, *L. occidentalis* Nuttall, *Pseudotsuga menziesii* (Mirbel) Franco, *P. mucronata* Rafinesque, *Tsuga heterophylla* (Rafinesque) Sargent (Pinaceae).

Tetropium velutinum LeConte, 1869: 382; Leng, 1884: 95; Holland, 1888: 91 (distr.); Casey, 1891: 23; Leng & Hamilton, 1896: 165; Fall, 1901: 143 (distr.); Fall & Cockerell, 1907: 191 (distr.); Garnett, 1918: 174 (distr., hosts); Rohwer, 1921: 442 (paras.); Craighead, 1923: 34, pl. 6, fig. 5, pl. 14; Essig, 1926: 450 (biol.); Hardy & Preece, 1927a: 187 (hosts); Leonard, 1928: 434 (distr.); Craighead & Middleton, 1930: 9 (biol.); Beaulne, 1932: 198 (hosts); Mank, 1934: 79 (distr.); DeLeon, 1934: 58 (hosts); Doane *et al.*, 1936: 168 (hosts); Hardy, 1942: 10 (biol.); Townes, 1944: 772 (paras.); Knull, 1946: 149, pl. 4, fig. 10; Leech, 1947: 108 (hosts); Hardy, 1948: 32 (distr.); Knowlton & Wood, 1950: 10 (distr.); Keen, 1952: 173 (biol.); Clark, 1956: 41 (distr.); Linsley, 1957: 15 (syn.); 1962: 88, fig. 33; Ross, 1967a: 23; 1967b: 25, figs 1-3; Ross, 1968: 11; Ross & Geistlinger, 1968: 14 (insecticides); Finlayson, 1969: 62 (paras.); Hatch, 1971: 97; Furniss & Carolin, 1977: 294 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 26 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 40, pl. 2, fig. 10; Yanega, 1996: 44, pl. 2, fig. 20; Linsley & Chemsak, 1997: 443 (hosts); Heffern 1998: 8 (distr.); Monné, M.A. & Hovore, 2006: 25 (checklist); Maier, 2017: 426 (distr.); Rice, MacRae & Merickel, 2017: 669 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 85

Tetropium columbianum Casey, 1912: 268; Clark, 1956: 421 (distr.); Lingafelter *et al.*, 2014: 43, fig. 44 q (holotype).

Type locality - Holotype male: Canada, British Columbia: Inverness. (USNM).

Tetropium hexagonum Casey, 1924: 228; Lingafelter *et al.*, 2014: 75, fig. 80u (holotype).

Type locality - Holotype male: United States, California: Humboldt County. (USNM).

Tetropium boreale Casey, 1924: 228; Lingafelter *et al.*, 2014: 31, fig. 31o (holotype).

Type locality - Holotype male: United States, Idaho: Priest Lake, Bonner County. (USNM).

ATIMIINI LeConte, 1873

Atimiini LeConte, 1873b: 292, 322; LeConte & Horn, 1883: 307; Leng, 1884: 61; Bradley, 1930: 228, 240; Linsley, 1939: 65 (rev.); Knull, 1946: 147; Linsley, 1962: 92; Arnett, 1962: 856; Hatch, 1971: 95; Monné, M.A., 1994: 27 (cat.); Monné, M.A., 2006: 113 (cat.); Monné, M.A. & Hovore, 2006: 25 (checklist); Bousquet, Heffern, Bouchard & Nearn., 2009: 21; Bouchard *et al.*, 2011: 465.

Type-genus: *Atimia* Haldeman, 1847.

Type-species: *Atimia tristis* Haldeman, 1847 (monotypy).

***Atimia* Haldeman, 1847**

Atimia Haldeman, 1847a: 56; 1847b: 373; LeConte, 185a: 25; Thomson, 1864: 224; 1865: 441; Lacordaire, 1868: 144; LeConte, 1873b: 322; LeConte & Horn, 1883: 307; Gahan, 1908: 144; Blatchley, 1910: 1041; Craighead, 1923: 34 (larva); Bradley, 1930: 240; Linsley, 1934a: 23; 1939: 65 (rev.); Knull, 1946: 147; Linsley, 1962: 92; Arnett, 1962: 856; Hatch, 1971: 95; Svacha & Danilevsky, 1987: 146; Monné, M.A., 2006: 113 (cat.); Monné, M.A. & Hovore, 2006: 25 (checklist); Monné, M.A., 2012: 133; Bousquet, Laplante, Hammond & Langor, 2017: 86

Type-species - *Atimia tristis* Haldeman, 1847 (monotypy) [= *Clytus confusus* Say, 1826].

Myctus Semenov & Plavilstshikov, 1937: 252.

Type-species - *Myctus maculipunctus* Semenov & Plavilstshikov, 1937 (original designation).

1. *Atimia confusa confusa* (Say, 1827)

Syntypes localities - Syntypes: United States, Pennsylvania, New Jersey, N.W. Territories. (Depository unknown). **Distribution** - Eastern and central United States and Canada. **Host plants** - *Chamaecyparis thyoides* (Linné) Britton, Sterns & Poggenburg, *Cupressus* sp., *Juniperus virginiana* Linné, *Thuja occidentalis* Linné (Cupressaceae), *Taxodium distichum* (Linné) Richard (Taxodiaceae).

Clytus confusus Say, 1827: 276; LeConte, 1859b: 333.

Clytus ? confusus; Haldeman, 1847a: 41.

Atimia confusa; Haldeman, 1847b: 373; LeConte, 1850a: 25; Thomson, 1861a: 373; 1864: 224; Lacordaire, 1868: 145; LeConte. 1873: 322; 1879:505; Harrington, 1881: 33; LeConte & Horn, 1883: 307; Leng, 1890: 10; 1890: 110 (distr.); Packard, 1890: 809 (hosts); Schwarz, 1896: 44 (biol.); Beutenmuller, 1896: 77 (hosts); Wickham, 1897c: 169; Smith, 1900: 290 (distr.); Schwarz, 1901: 44; Fall, 1901: 247 (distr.); Ulke, 1903: 26, 50 (distr.); Smith, 1910: 329; Blatchley, 1910: 1042, fig. 443; Garnett, 1918: 211 (distr.); Craighead, 1923: 35 (larva); Kirk & Knull. 1926: 41 (distr.); Leonard, 1928: 448 (distr.); Beaulne, 1932: 203 (hosts); Davis, 1932: 213 (distr.); Linsley, 1934: 24; Doane et al., 1936: 185; Brimley. 1938: 216 (distr.); Linsley, 1939: 68, pl. 14, fig. 7; Knull, 1946: 147, pl. 4, fig. 14; Craighead, 1950: 237 (biol.); Beal, Haliburton & Knight, 1952: 112 (biol.); Linsley, 1962: 95; Baker. 1972: 198; Swan & Papp, 1972: 444, fig. 935; Gosling, 1973: 68 (distr.); Laliberté, Chantal & LaRochelle, 1977: 91 (biol.); Furniss & Carolin, 1977: 292 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Chemsak, 1996: 51; Monné, M.A. & Hovore, 2006: 25 (checklist); Lingafelter, 2007: 76, 139, 2 figs, pl. 18e (hosts); Bousquet, Laplante, Hammond & Langor, 2017: 86, pl. 15; Klingeman *et al.*, 2017: 295 (distr.)

Atimia confusa confusa; Linsley, 1962: 96, pl. 1, fig. 5; Turnbow & Franklin, 1980: 338 (distr.); Rice, 1981: 460 (distr., hosts); MacRae, 1993: 227 (distr.); Monné, M.A., 1994: 27 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 53, pl. 3, fig. 4; Yanega, 1996: 44, pl. 25, fig. 284; Linsley & Chemsak, 1997: 348 (hosts); Heffern, 1998: 8 (distr.); Schiefer, 1998: 116 (distr.); Peck & Thomas, 1998: 117 (distr.); Vlasák & Vlasakova, 2002: 209 (distr., hosts); Holt, 2013: 246 (distr.); Spomer, 2014: 298, fig. 2 (distr.); Webster et al., 2016: 12 (distr., hosts).

Atimia tristis Haldeman, 1847: 56.

Syntypes locality - Syntypes: United States, Pennsylvania. (MCZN).

1a. *Atimia confusa maritima* Linsley, 1939

Type locality - Holotype male: United States, California: Coastal Central California. (CASC).

Distribution - United States (Coastal central California). **Host plants** - *Cupressus goveniana* Gordon, *C. macrocarpa* Hartweg ex Gordon (Cupressaceae).

Atimia confusa maritima Linsley, 1939: 70; Linsley, 1962: 97, pl. 1, fig. 7, fig. 34; Tyson, 1966: 202 (hosts); Swan & Papp, 1972: 445; Furniss & Carolin, 1977: 292 (hosts); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Monné, M.A., 1994: 28 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 54, pl. 3, fig. 6; Linsley & Chemsak, 1997: 349 (hosts); Monné, M.A. & Hovore, 2006: 25 (checklist);

Atimia maritima; Barr, 1947: 58 (hosts); Keen, 1952: 176 (hosts).

2. *Atimia dorsalis* LeConte, 1869

Syntypes locality - Syntypes: Canada, Vancouver's Island. (MCZN). **Distribution** - Pacific coast of North America, from British Columbia to the San Pedro Martir Mts. of Baja California to Utah. **Host plants**- *Calocedrus decurrens* (Torrey) Florin, *Cupressus forbesii* Jepson, *C. macrocarpa* Hartweg ex Gordon, *Juniperus* sp., *Thuja plicata* Donn ex D.Don (Cupressaceae), *Sequoia sempervirens* (D. Don) Endlicher (Taxodiaceae).

Atimia dorsalis LeConte, 1869: 385; 1873b: 322; Horn, 1876: 199 (distr.); LeConte & Horn, 1883: 307; Leng, 1890: 10; Horn, 1894: 339 (distr.); Fall, 1901: 147 (distr.); Wright & Coolidge, 1908: 68 (distr.); Garnett, 1918: 211 (distr.); Craighead, 1923: 34, pl. 3, fig. 4, pl. 6, fig. 2, pl. 8, fig. 5, pl. 23, fig. 5. pl. 25, figs 3, 4 (larva, hosts); Hardy, 1926: 32, pl. 4, fig. 6; Essig, 1926: 457 (hosts); Beaulne, 1932: 203 (hosts); Linsley, 1934a: 24; 1936: 199 (biol.); Doane *et al.*, 1936: fig. 94; Moore, 1937: 90 (distr.); Linsley, 1939: 70, pl. 14, fig. 4 (rev.); 1942: 28 (distr.); Leech, 1944: 30 (biol.); Hardy, 1944: 17 (distr., hosts); DeLeon, 1952: 81 (hosts); Keen, 1952: 176 (hosts); Ross, 1968: 12 (biol.); Barr & Penrose, 1969: 89 (distr.); Bousquet, Laplante, Hammond & Langor, 2017: 87, pl. 15

Atimia confusa dorsalis; Linsley, 1962: 96, pl. 1, fig. 6, fig. 34; Tyson, 1966: 202 (hosts); Hatch, 1971: 98, pl. 9, fig. 8; Swan & Papp, 1972: 445; Furniss & Carolin, 1977: 292 (hosts); Hovore, 1988: 27 (distr.); Chemsak, Linsley & Noguera, 1992: 25 (cat.); Napp, 1994: 277, figs 20, 75, 104, 133, 146, 147, 171, 172, 228, 241; M.A., & Giesbert, 1994: 23 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 54, pl. 3, fig. 5 (rev.); Linsley & Chemsak, 1997: 349 (hosts); Monné, M.A., 2002: 36 (cat. hosts); Monné, M.A., 2006: 113 (cat.); Monné, M.A. & Hovore, 2006: 25 (checklist); Rice, MacRae & Merickel, 2017: 669 (distr.)

3. *Atimia gannoni* Hovore & Giesbert, 1974

Type locality - Holotype male: United States, California: Los Angeles County, Little Mt. Gleason 5 miles West of Mill Creek Summit. (CACS). **Distribution** - Southern California northward to Tuolumne Co. **Host plants** - *Calocedrus decurrens* (Torrey) Florin. (Cupressaceae).

Atimia gannoni Hovore & Giesbert, 1974: 139, figs 1, 3; Furniss & Carolin, 1977: 293 (hosts); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., 1994: 28 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 57, pl. 3, fig. 8; Monné, M.A. & Hovore, 2006: 25 (checklist);

4. *Atimia helena* Linsley, 1934

Type locality - Holotype male: United States, California: Cypress Ridge, Marin County. (CASC). **Distribution** - Dry foothills of southern, central and northern California; San Diego, Marin, Sonoma, Napa and Lake Counties. **Host plants** - *Cupressus forbesii* Jepson, *C. goveniana* Gordon, *C. pygmaea* Sargent, *C. sargentii* Jepson (Cupressaceae).

Atimia helena Linsley, 1934a: 25; Doane *et al.*, 1936: 185; Linsley & Usinger, 1936: 55 (distr.); Linsley, 1939: 72, pl. 14, fig. 5; 1962: 97, pl. 1, fig. 2; Frankie & Jensen, 1971: 287, figs 1-4 (biol.); Furniss & Carolin, 1977: 293 (hosts); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., 1994: 28 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 61, pl. 3, fig. 11; Linsley & Chemsak, 1997: 349 (hosts); Monné, M.A. & Hovore, 2006: 25 (checklist);

5. *Atimia hoppingi* Linsley, 1939

Type locality - Holotype male: United States, Washington: Mt. Rainier, White R. Camp. (CASC). **Distribution** - Washington to northern California. **Host plants** - *Chamaecyparis nootkatensis* (D. Don) Spach, *Cupressus macnabiana* Andrew Murray (Cupressaceae).

Atimia hoppingi Linsley, 1939: 75, pl. 14, fig. 6; Keen, 1952: 176 (hosts); Linsley, 1962: 99, pl. 1, fig. 3, fig. 34; Hatch, 1971: 98, pl. 9, fig. 9; Hovore & Giesbert, 1974: 141, fig. 2; Furniss & Carolin, 1977: 293 (hosts); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., 1994: 28 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 58, pl. 3, fig. 9; Linsley & Chemsak, 1997: 349 (hosts); Monné, M.A. & Hovore, 2006: 25 (checklist); Bousquet, Laplante, Hammond & Langor, 2017: 87

6. *Atimia huachucae* Champlain & Knoll, 1922

Type locality - Holotype female: United States, Arizona: Huachuca Mts. (FMNH). **Distribution** - United States (Southern Nevada, Colorado to Texas), northern Mexico

(Chihuahua). **Host plants** - *Cupressus arizonica* Greene, *Juniperus deppeana* Steudel (Cupressaceae).

Atimia huachucae Champlain & Knull, 1922: 148; Linsley, 1934a: 24; Doane *et al.*, 1936: 185; Linsley, 1939: 67, pl. 14, fig. 1 (rev.); Keen, 1952: 176 (hosts); Linsley, Knull & Statham, 1961: 9 (distr.); Linsley, 1962: 93, pl. 1, fig. 4, fig. 34; Chemsak, 1977: 173 (type); Furniss & Carolin, 1977: 293 (hosts); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Noguera & Chemsak, 1996: 396 (cat.); Chemsak, 1996: 55, pl. 3, fig. 7 (rev.); Linsley & Chemsak, 1997: 349 (hosts); Heffern, 1998: 8 (distr.); Monné, M. A., 2002: 36 (cat. hosts); Monné, M.A., 2006: 113 (cat.); Monné, M.A. & Hovore, 2006: 25 (checklist); Noguera & Gutierrez, 2016: 657 (distr.)

7. *Atimia vandykei* Linsley, 1939

Type locality - Holotype male: United States, New Mexico: Fort Wingate. (CASC).

Distribution - New Mexico to California. **Host plants** - *Juniperus occidentalis* Hooker (Cupressaceae).

Atimia vandykei Linsley, 1939: 74, pl. 14, fig. 8; 1962: 98, pl. 1, fig., 1, fig. 34; Furniss & Carolin, 1977: 293 (hosts); Cope, 1984: 28 (hosts); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., 1994: 28 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 59, pl. 3, fig. 10; Linsley & Chemsak, 1997: 349 (hosts); Monné, M.A. & Hovore, 2006: 25 (checklist);

Paratimia Fisher, 1915

Paratimia Fisher, 1915: 78; Bradley, 1930: 240; Linsley, 1939: 77; Arnett, 1962: 856, 876; Linsley, 1962: 99; Hatch, 1971: 95; Monné, M.A., 1994: 29; Chemsak, 1996: 62; Monné, M.A. & Hovore, 2006: 25 (checklist);

Type-species - *Paratimia conicola* Fisher, 1915 (original designation).

1. *Paratimia conicola* Fisher, 1915

Type locality - Holotype male; United States, California: Monumental Mines. (USNM).

Distribution - United States (Coast Range Mts. of Oregon and California). **Host plants** - *Pinus contorta* Douglas ex Loudon, *P. tuberculata* Gordon (Pinaceae).

Paratimia conicola Fisher, 1915: 78; Rohwer, 1921: 451 (paras.); Craighead, 1923: 35, pl. 6, fig. 7, pl. 40, figs 6, 7; Essig, 1926: 795 (paras.); Doane *et al.*, 1936: 185, fig. 95; Linsley, 1939: 78, pl. 14, fig. 2; Townes, 1944: 773 (paras.); Keen, 1952: 23 (biol.); Ruckes, 1959: 134 (biol.); Townes & Townes, 1960: 384 (paras.); Linsley, 1962: 100, pl. 1, fig. 8; Hatch, 1971: 98; Chemsak, 1972: 150 (hosts); Furniss & Carolin, 1977: 293 (biol.); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., 1994: 29 (cat.); Monné, M.A., & Giesbert, 1994: 23 (checklist); Chemsak, 1996: 63, pl. 3, fig. 12; Linsley & Chemsak, 1997: 415 (hosts); Monné, M.A. & Hovore, 2006: 25 (checklist); Lingafelter *et al.*, 2014: 44, fig. 46g (holotype); Haack, 2017: 114 (hosts)

SAPHANIINI Gistel, 1848

Saphanidae Gistel, 1848: (8).

Saphanites; Fairmaire, 1868: 127.

Saphanides; Lacordaire, 1869: 211.

Saphanini; Aurivillius, 1912: 23; Bouchard *et al.*, 2011: 465.

Type-genus: *Saphanus* Audinet-Serville, 1834.

Type-species: *Callidium spinosum* Fabricius, 1792 (= *Callidium piceum* Laicharting, 1784) (monotypy).

Michthysomini LeConte, 1873b: 332. Comment: incorrect original stem formation, not in prevailing usage.

Michthysomini; LeConte & Horn, 1883: 317; Leng & Hamilton, 1896: 103.

Michthysomini; Linsley, 1962: 101; Arnett, 1962: 856.

Type-genus: *Michthysoma* J. L. LeConte, 1850 [as *Michthysoma*, incorrect subsequent

spelling of type genus name, not in prevailing usage].

Type-species: *Michthisoma heterodoxus* LeConte, 1850 (monotypy).

***Michthisoma* LeConte, 1850**

Michthisoma LeConte, 1850a: 30; Thomson, 1861a: 229; Linsley, 1962: 101; Arnett, 1962: 856; Monné, M.A., 1993: 1 (cat.).

Michthisoma; Thomson, 1864: 131; Leng in Hamilton, 1896: 103; Davis & Leng, 1912: 13; Craighead, 1923: 103; Bradley, 1930: 228; Linsley, 1959: 127.

Type-species - *Michthisoma heterodoxus* LeConte, 1850 (monotypy).

1. *Michthisoma heterodoxum* LeConte, 1850

Type locality - Holotype: United States, Georgia: Yona Mountain (near the top). (MCZN).

Distribution - United States (Mountains of North Carolina to Georgia). **Host plants** - *Castanea* sp., *Quercus* sp. (Fagaceae), *Tilia* sp. (Malvaceae).

Michthisoma heterodoxum LeConte, 1850a: 30; Linsley, 1962: 102; Perry, 1977: 97 (distr.); Turnbow & Franklin, 1980: 338 (distr.); Chemsak, Linsley & Noguera, 1992: 26 (cat.); Monné, M.A., 1993: 1 (cat.); Monné, M.A., & Giesbert, 1994: 24 (checklist); Yanega, 1996: 45, pl. 17, fig. 208; Lingafelter, 2007: 68, 147, 2 figs, pl. 16 g (hosts); Vlasák, 2014: 318 (hosts); Klingeman *et al.*, 2017: 295 (distr.)

Michthisoma heterodoxum; Melsheimer, 1853: 103 (cat.); Thomson, 1864: 131; LeConte, 1873b: 332; LeConte & Horn, 1883: 317; Leng, 1896: 103 (distr.); Beutenmuller, 1903: 519 (biol., distr.); Leng, 1910: 77 (distr.); Mutchler, 1912: 213 (distr.); Craighead, 1923: 103, figs (larva); Brimley, 1938: 216 (distr.); Sherman, 1946: 127 (distr.); Fattig, 1947: 27 (distr.).

SPONDYLIDINI Audinet-Serville, 1832

Spondylii Audinet-Serville, 1832: 123.

Spondylidae; LeConte, 1851: 99. 1873: 279.

Spondylitae; Thomson, 1860: 261; 1864: 9.

Spondylinae; LeConte & Horn, 1883: 266; Casey, 1912: 218; Linsley, 1962: 62; Arnett, 1962: 854.

Spondylini; Aurivillius, 1912: 13; Bradley, 1930: 228.

Spondylidini; Bousquet, Heffern, Bouchard & Nearn., 2009: 22; Bouchard *et al.*, 2011: 465.

Type-genus: *Spondylis* Fabricius, 1775.

Type-species: *Attelabus buprestoides* Linnaeus, 1758 designated by Latreille (1810: 431).

***Neospondylis* Sama, 2005**

Neospondylis Sama, 2005: 8; Monné, M.A., 2006: 114 (cat.); Monné, M.A. & Hovore, 2006: 23 (checklist); Monné, M.A., 2012: 134.

Type-species - *Spondylis upiformis* Mannerheim, 1843 (original designation).

Spondylis; LeConte, 1873b: 281; Bates, 1879: 14; LeConte & Horn, 1883: 266; Casey, 1912: 218; Bradley, 1930: 230; Linsley, 1962: 63; Arnett, 1962: 856; Napp, 1994: 272; Chemsak, 1996: 12. (not Fabricius, 1775).

Type-species - *Attelabus buprestoides* Linnaeus, 1758 (Thomson designation, 1864: 272).

1. *Neospondylis upiformis* (Mannerheim, 1843)

Type locality - Holotype: United States, Alaska: Sitka. (FMUH). **Distribution** - Coniferous forest of North America from Alaska southeast to the Great Lakes region, south into the Rocky Mountains and along the Pacific coast, Mexico (Durango). **Host plants** - *Picea glauca* (Moench) Voss, *Pinus contorta* Douglas ex Loudon, *P. monticola* Douglas ex D. Don (Pinaceae).

Spondylis upiformis Mannerheim, 1843: 304; LeConte, 1851: 99; White, 1853: 4; LeConte, 1857: 23; 1873b: 282; LeConte & Horn, 1883: 267; Harrington, 1890: 185 (distr.); Fall, 1901: 142; Fall & Cockerell, 1907: 190 (distr.); Casey, 1912: 218; Tanner, 1927: 33 (distr.); 1928: 277 (distr.); Knowlton, 1930: 56 (distr.); Beaulne, 1932: 107 (hosts); Knowlton, 1934: 86 (distr.);

- DeLeon, 1934: 57 (hosts); Mank, 1934: 79 (distr.); Doane, Van Dyke, Chamberlin & Burke, 1936: 166, fig. 80; Saalas, 1936: 54, pl. 5, fig. 38 (morphol.); Linsley, 1938: 105 (syn.); Leech, 1947: 108 (hosts); Hardy, 1948: 101 (distr.); Knowlton & Wood, 1950: 10 (distr.); Keen, 1952: 196 (hosts); Clark, 1956: 41 (distr.); Linsley, Knull & Statham, 1961: 7 (distr.); Linsley, 1962: 63, figs. 23, 24; Gardiner, 1970b: 33, figs 1-3 (larva); Hatch, 1971: 92, pl. 9, fig. 4; Gosling, 1973: 68, fig. 6 (biol.); Kirk & Balsbaugh, 1975: 97 (distr.); Laliberté, Chantal & LaRochelle, 1977: 99 (biol.); Furniss & Carolin, 1977: 290, fig. 172; Gosling, 1986: 157 (hosts); Chemsak, Linsley & Noguera, 1992: 23 (distr.); Monné, M.A., & Giesbert, 1994: 21 (checklist); Noguera & Chemsak, 1996: 396 (distr.); Chemsak, 1996: 12, pl. 1, fig. 7; Yanega, 1996: 28, pl. 1, fig. 4; Hefern, 1998: 7 (distr.); O'Neill, Fultz & Ivie, 2008: 33 (distr.).
- Spondylus upiformis*; Smith & Hurley, 2005: 448 (distr., error).
- Spandylis (Spondylis) upiformis*; Lameere, 1902b: 328 (revis.).
- Neospondylis upiformis*; Sama, 2005: 39, figs 1, 4-8, 15-17, 19, 21, 25-30, 36; Monné, M.A. & Hovore, 2006: 23 (checklist); Webster, MaCorquodale & Majka, 2009: 291 (distr.); Majka & Ogden, 2010: 12, figs 1, 2 (distr.); Webster *et al.*, 2016: 311 (distr.); Gutiérrez & Noguera, 2015: 139 (distr.); Webster, 2016: 486; Rice, MacRae & Merickel, 2017: 669 (distr.).
- Spondylis laticeps* LeConte, 1850b: 233; 1851: 99; White, 1853: 4.
- Type locality** - Holotype: United States, Michigan: Lake Superior (Eagle Harbor). (MCZN).
- Spondylis collaris* Casey, 1912: 218; Lingafelter *et al.*, 2014: 42, fig. 44c (holotype).
- Type locality** - Holotype male: United States: Utah. (USNM).
- Spondylis robustula* Casey, 1912: 219; Lingafelter *et al.*, 2014: 310, fig. 144k (holotype).
- Type locality** - Holotype male: United States, Washington State. (USNM).
- Spondylis subpubescens* Casey, 1912: 219; Lingafelter *et al.*, 2014: 328, fig. 165g (lectotype).
- Type locality** - Lectotype male: United States, California. (USNM).
- Spondylis basalis* Casey, 1912: 220; Lingafelter *et al.*, 2014: 26, fig. 26a (holotype).
- Type locality** - Holotype female: United States, Washington State (USNM).
- Spondylis parva* Casey, 1924: 226; Lingafelter *et al.*, 2014: 296, fig. 128c (holotype).
- Type locality** - Holotype female: United States, Washington: Port Columbia (Pudget Sound). (USNM).

***Scaphinus* LeConte, 1851**

- Scaphinus* LeConte, 1851: 100; Thomson, 1861a: 272; 1864: 272, 465; Lacordaire, 1868: 199; LeConte, 1873b: 281; LeConte & Horn, 1883: 266; Bradley, 1930: 230; Linsley, 1962: 66; Arnett, 1962: 856; Monné, M.A., 1994: 17; Chemsak, 1996: 14; Monné, M.A. & Hovore, 2006: 22 (checklist);
- Spondylis (Scaphinus)*; Lameere, 1902b: 330.
- Type-species** - *Spondylis sphaericollis* LeConte, 1847 (= *Prionus muticus* Fabricius, 1801) (monotypy).

1. *Scaphinus muticus* (Fabricius, 1801)

- Type locality** - Holotype: America Boreali. (BMNH). **Distribution** - Southeastern United States from Florida to Virginia and Arkansas. **Host plants** - *Taxodium distichum* (Linné) Richard (Taxodiaceae).
- Prionus muticus* Fabricius, 1801: 265; Zimsen, 1964: 164 (type).
- Scaphinus muticus*; Aurivillius, 1912: 15 (cat., syn.); Doane *et al.*, 1936: 266; Brimley, 1938: 210 (distr.); Loding, 1945: 113 (distr.); Sherman, 1946: 126 (distr.); Linsley, 1962: 66, fig. 24; Monné, M.A., & Giesbert, 1994: 21 (checklist);
- Spondylis sphaericollis* LeConte, 1847: 93; Perry, 1974: 215 (distr.); Turnbow & Franklin, 1980: 338 (distr.); Chemsak, Linsley & Noguera, 1992: 23 (cat.); Monné, M.A., 1994: 18 (cat.); Chemsak, 1996: 14; Yanega, 1996: 28, pl. 1, fig. 3; Linsley & Chemsak, 1997: 432 (hosts); Schiefer, 1998: 115 (distr.); Peck & Thomas, 1998: 116 (distr.); Sama, 2005: 3, 6, 9, figs 3, 23, 38; Monné, M.A. & Hovore, 2006: 22 (checklist); Holt, 2013: 246 (distr.).
- Scaphinus sphaericollis*; LeConte, 1851: 100; White, 1853: 5; Thomson, 1861a: 272; 1864: 272; Lacordaire, 1868: 199; LeConte, 1873b: 282; Lacordaire, 1876: 29, pl. 83, figs 5-5b; LeConte & Horn, 1886: 267.

Spondylis (Scaphinus) sphaericollis; Lameere, 1902b: 331.

Type locality - Holotype: United States, Georgia: Savannah. (MCZ)

References

- Abdullah. M. (1968) *Parandra (Neandra) brunnea quebecensis*, a new subspecies of the Parandrinae (Col., Cerambycidae). *The Entomologist's Monthly Magazine*, 104, 75.
- Alexander, D.R. (1958) A preliminary survey of the cerambycids (Insecta: Coleoptera) of Oklahoma. *Proceedings of the Oklahoma Academy of Sciences*, 38, 43–52.
- Amith, J.D. & Lingafelter, S.W. (2017) Ethnoentomological and distributional notes on Cerambycidae and other Coleoptera of Guerrero and Puebla, Mexico. *The Coleopterists Bulletin*, 71 (2), 301-314.
doi.org/10.1649/0010-065X-71.2.301
- Anderson R. F. (1960) *Forest and Shade Tree Entomology*. John Wiley & Sons Inc. New York: i-vi + 1-428.
- Androw, R.A. & Keeney, G.D. (1999) New records of Ohio long-horned beetles (Coleoptera: Cerambycidae) *The Great Lakes Entomologist*, 32 (1-2), 1-7
- Arnaud, P.H.Jr. (1968) Rearing of *Ergates spiculatus spiculatus* (LeConte) from a house beam at Saratoga, California (Coleoptera: Cerambycidae). *The Pan-Pacific Entomologist*, 44(1), 81.
- Arnett, R.H. (1962) *The beetles of the United States*. Catholic Univ. America Press, Washington, D. C., 1112 pp.
- Audinet-Serville, J.G. (1832) Nouvelle classification de la famille des longicornes. *Annales de la Société Entomologique de France*, 1(1), 118–201.
- Audinet-Serville, J.G. (1834) Nouvelle classification de la famille des longicornes.(suite). *Annales de la Société Entomologique de France*, 3(1), 5–110.
- Audureau, A. (2008) Contribution a la connaissance des Cerambycidae de la reserve privée forestiere de Domitila (Nicaragua). *Lambillionea*, 108(3), supplement, 3-21.
- Audureau, A. & Roguet, J-P. (2018) Contribution à la Connaissance des Cerambycidae de la Région Nord du Nicaragua (Coleoptera, Cerambycidae). *Les Cahiers Magellanes*, 30, 56-104.
- Aurivillius, C. (1912) *Coleopterorum Catalogus*, pars 39, Cerambycidae: Cerambycinae. W. Junk, Berlin, 574 pp.
- Baker, W.L. (1972) Eastern Forest Insects. *Miscellaneous Publications*, United States Department of Agriculture, Forest Service, 1175, 1–642.
- Barbour, J.D., Cervantes, D.E., Lacey, E.S. & Hanks, L.M. (2006) Calling behavior in the primitive longhorned beetle *Prionus californicus* Mots. *Journal of Insect Behavior*, 19(5), 623-629.
- Barbour, J.D., Lacey, E.S. & Hanks, L.M. (2007) Cuticular hydrocarbons mediate mate recognition in a species of longhorned beetle (Coleoptera: Cerambycidae) of the primitive Subfamily Prioninae. *Annales of the Entomological Society of America*, 100(2), 333-338.
0013-8746/07/0333D0338\$04.00/0
- Barbour, JD, Alston DG, Walsh DB, Pace M, Hanks LM (2019) Mating disruption for managing *Prionus californicus* (Coleoptera: Cerambycidae) in hop and sweet cherry. *Journal of Economic Entomology*, 112,1130–1137 [PDF](#)
- Barr, W. F. (1947) Parasites of two species of Coleoptera associated with Monterey Cypress. *The Pan-Pacific Entomologist*, 23 (2), 58.
- Barr, W.F. & Penrose, R.L. (1969) Notes on the distribution and bionomics of some Idaho Cerambycidae. *Great Basin Naturalist*, 29(2), 88–95.
- Barrett, R.E. (1932) An annotated list of the insects and arachnids affecting the various species of walnuts or members of the genus *Juglans* Linn. *University of California, Publications in Entomology*, 5, 275–310.
- Bates, H.W. (1872) On the longicorn Coleoptera of Chontales, Nicaragua. *The Transactions of the Entomological Society of London*, 1872, 163–238.
<https://doi.org/10.1111/j.1365-2311.1872.tb01888.x>
- Bates, H.W. (1879) *Biologia Centrali-Americana*, Insecta, Coleoptera. London, 5, 1–16.

- Bates, H.W. (1884) *Biologia Centrali-Americana*, Insecta, Coleoptera, suppl. to Longicornia. London, 5, 225–248.
- Bates, H.W. (1885) *Biologia Centrali-Americana*, Insecta, Coleoptera, suppl. to Longicornia. London, 5, 249–436.
- Bates, H.W. (1892) Additions to the Longicornia of Mexico and Central America, with remarks on some of the previously recorded species. *The Transactions of the Entomological Society of London*, 1892, 143–183.
- Beal, J.A., Haliburton, W. & Knight, F.B. (1952) Forest insects of the southeast: with special reference to species occurring in the Piedmont Plateau of North Carolina. *Bulletin of the Duke University School of Forestry* 14, 3-168
- Beaulne, J.T. (1932) Longicornes nuisibles aux végétaux ligneux du Canada. *Naturaliste Canadien*, 59(10), 196–203; (11), 219–222.
- Becker, W.B. (1942) *Prionus laticollis* Drury in a subterranean wooden duct for telephone cables *Journal of Economic Entomology* 35 (4), 608.
- Belyea, R.M. (1952) Death and deterioration of balsam fir weakened by spruce budworm defoliation in Ontario. Part 1. *The Canadian Entomologist*, 84 (11), 325-335.
- Benham, G.S.Jr. (1969) The Pupa of *Prionus laticollis* (Coleoptera: Cerambycidae) *Annals of the Entomological Society of America*, 62 (6), 1331-1335, 4 figs.
- Benham, G. S. Jr. (1970) Gross morphology and transformation of the digestive tract of *Prionus laticollis* (Coleoptera: Cerambycidae) *Annals of the Entomological Society of America*, 63 (5), 1413-1419, 5 figs.
- Benham, G. S. Jr. (1971) Microorganisms associated with immature *Prionus laticollis* (Col. Cerambycidae) *Journal of Invertebrate Pathology*, 18 (1), 89-93.
- Benham, G, S. Jr. & Farrar R. J. (1976) Notes on the biology of *Prionus laticollis* (Coleoptera: Cerambycidae) *The Canadian Entomologist*, 108 (6), 569-576, 9 figs.
- Bense, U. (1995) 1995. Longhorn Beetles. Illustrated Key to the Cerambycidae and Vesperidae of Europe Weikersheim : Margraf, 1995: 1-512, !260 figs
- Bethune, C.J.S. (1872) Insects of the northern parts of British America (from Kirby's "Fauna Boreali-Americana, Insecta). *The Canadian Entomologist*, 4(3), 52–57; (5); 93–96; (6), 111–118.
- Bethune, C.J.S. (1877a) A few common wood-boring beetles *7th Annual Report of the Entomological Society of the Province of Ontario 1877: 22-30*
- Bethune, C.J.S. (1877b) A few common wood-boring beetles *The Canadian Entomologist*, 9 (12), 221-226.
- Beutenmuller, W. (1896) Food habits of North American Cerambycidae. *Journal of the New York Entomological Society*, 4, 73–81.
- Biffi, G. & Fuhrmann, J. (2013) Immatures of *Parandra (Tavandra) longicollis* Thomson, 1861 and comments on the larvae of Parandrinae (Coleoptera: Cerambycidae). *Insecta Mundi*, 0323, 1-14.
- Blackman, M.W. & Stage, H.H. (1918) on insects bred from the bark and wood of the American larch, *Larix laricina* (Du Roc) Koch *Technical Publications of the New York State College of Forestry* 10: 11-115, 9 pls.
- Blaisdell, F.E. (1892) Notes on the habits of some species of Coleoptera observed in San Diego County. *Insect Life*, 5(1), 33–36.
- Blanchard, C.E. (1845) *Histoire des insectes, traitant de leurs moeurs et de leurs métamorphoses en général, et comprenant une nouvelle classification fondée sur leurs rapports naturels*. Didot, Paris, 2, 1–524.
- Blanchard, F. 1887. Notes on Coleoptera. *Entomologia Americana*, 3(5), 85–88.
- Bland, J.H.B. (1861) Catalogue of the longicorn Coleoptera taken at the vicinity of Philadelphia. *Proceedings of the Entomological Society of Philadelphia*, 1, 93–101.
- Blatchley, W.S. (1910) On the Coleoptera known to occur in Indiana. An illustrated descriptive catalogue of the Coleoptera or beetles (exclusive of the Rhynchophora) known to occur in Indiana. *Bulletin of the Indiana Department of Geology and Natural Resources*, 1, 1–1386.
- Bouchard, P. (2014) *The book of Beetles. A life size guide to six hundred of nature's gems*. Ivy Press Limited, London, 656 p.

- Bouchard, P., Bousquet, Y., Davies, A.E., Alonso-Zarazaga, M.A., Lawrence, J.F., Lyal, C.H.C., Reid C.A.M., Schmitt M., Ślipiński S.A., & Smith A.B.T., (2011) Family-group names in Coleoptera (Insecta). *ZooKeys*, 88, 1–972.
- Bousquet, Y. (2007) Nomenclature and bibliographical notes on Cerambycidae (Coleoptera). *The Coleopterists Bulletin* 61(4), 616-631.
- Bousquet, Y., Heffern, D.J., Bouchard, P. & Nearn, E.H. (2009) Catalogue of family-group names in Cerambycidae (Coleoptera). *Zootaxa* 2321, 1-80.
- Bousquet, Y. & Bouchard, P. (2013) The genera in the second catalogue (1833-1836) of Dejean's Coleoptera collection. *Zookeys*, 282, 1-219.
- Bousquet, Y., Laplante, S., Hammond, H.E.J. & Langor, D. W. (2017) *Cerambycidae (Coleoptera) of Canada and Alaska: identification guide with nomenclatural, taxonomic, distributional, host-plant, and ecological data*. Prague, Nakladatelství Jan Farkac, 300 pages
- Bouyer, T., Drumont, A. & Santos-Silva, A. (2012) Revision of African Parandrinae (Coleoptera, Cerambycidae). *Insecta Mundi*, 0241, 1-85, 222 figs.
- Bouyer, T. & Santos-Silva, A. (2016) On *Hovatoma gramreta* Gilmour, 1956, a junior synonym of *Strongylaspis corticaria* (Erichson, 1848) (Coleoptera, Cerambycidae, Prioninae) *Les Cahiers Magellanes* (NS) 23, 87-90, 4 figs.
- Bradley, J.C. (1930) *A manual of the genera of beetles of America north of Mexico. Keys for the determination of the families, tribes and genera of Coleoptera with a systematic list of the genera and higher groups*. Daw, Illston & Co., Ithaca, 360 pp.
- Brimley, C.S. (1938) *The insects of North Carolina, being a list of the insects of North Carolina and their close relations*. North Carolina Department of Agriculture, Division of Entomology, Raleigh, 560 pp.
- Britton, W.E. (1920) Check-list of the insects of Connecticut. *Connecticut State Geological and Natural History Survey Bulletin*, 31, 1–397
- Brooks F. E. (1915) The *Parandra* borer as an orchard enemy *USDA Bulletin* 262, 1-7.
- Brown. W.J. (1930) Coleoptera of the north shore of the gulf of the St. Lawrence. *The Canadian Entomologist*, 62 (11), 239-246.
- Brown. A. W. (1934) A contribution to the insect fauna of Timagami. *The Canadian Entomologist*, 66 (9), 220-231
- Browne, D.J. & Pck, S.B. (1996) The long-horned beetles of south Florida (Cerambycidae Coleoptera): biogeography and relationships with the Bahama Islands and Cuba. *Canadian Journal of Zoology*, 74(12), 2154-2169.
- Browne, D.J., Peck, S.B. & Ivie, M.A. (1993) The longhorn beetles (Coleoptera, Cerambycidae) of the Bahama Islands with an analysis of species-area relationships, distribution patterns, origin of the fauna and an annotated species list. *Tropical Zoology*, 6, 27-53.
- Bruner, L. (1891) Some insects of special interest to the fruit growers of Nebraska *Annual Report of the Nebraska Horticulture Society*, 1891: 191-266.
- Bruner, L. (1894) Insects enemies of the apple tree and its fruit *Annual Report of the Nebraska Horticulture Society*, 1894, 154-223.
- Buquet, J.B.L. (1840) G. Mécosarthron. *Mecosarthron* Buquet. *Magasin de Zoologie*, 1840, pl. 52.
- Cardona-Duque, J.C., Santos-Silva, A. & Wolff, M. (2010) Parandrinae (Coleoptera, Cerambycidae) de Colombia. *Revista Colombiana de Entomologia*, 36(1), 135-157.
- Carl, G.C. & Hardy, G.A. (1945) Flora and fauna of the Paradise Mine area, British Columbia. *Report of the Provincial Museum of Natural History & Anthropology of British Columbia*, 1944, C18–C38.
- Casey, T.L. (1890) Coleopterological notices. II. *Annals of the New York Academy of Sciences*, 5, 307–504.
- Casey, T.L. (1891) Coleopterological notices. III. *Annals of the New York Academy of Sciences*, 6, 9–214.
- Casey, T.L. (1893) Coleopterological notices. V. *Annals of the New York Academy of Sciences*, 7, 281–606, 1 pl.
- Casey, T.L. (1899) New species of *Pemphus* and *Tragosoma* (Coleoptera). *Entomological News*, 10(4), 97–99.

- Casey, T.L. (1912) Studies in the Longicornia of North America. *Memoirs on the Coleoptera*, 3, 215–376.
- Casey, T.L. (1924) Additions to the known Coleoptera of North America. *Memoirs on the Coleoptera*, 11, 1–347.
- Castle, D.M. & Laurent, P. (1897) April collecting in Georgia and Florida. *Entomological News*, 8(1), 7–9.
- Cerda, M. (1974) Revisión de los Prioninae de Chile. *Revista Chilena de Entomología*, 8, 41–46.
- Cervantes, D.E., Hanks, L.M., Lacey, E.S. & Barbour, J.D. (2006) First documentation of a volatile pheromone in a longhorned beetle (Coleoptera, Cerambycidae) of the primitive subfamily Prioninae. *Annals of the Entomological Society of America* 99(4), 718–722.
- Chagnon, G. (1897) Faune coléoptérologique au Manitoba. *Naturaliste Canadien*, 24(4), 14–16; 122–125.
- Chagnon, G. (1905) Canadian Cerambycidae. The species of the past season. *Entomological News*, 16(2), 35–36.
- Chagnon, G. (1917) *A preliminary list of the insects of the Province of Quebec*. Part III. Coleoptera. Québec Société des Protections des Plantes, Montreal, Suppl. pp. 161–277.
- Chagnon, G. (1936) Contribution à l'étude des Coléoptères de la Province de Québec. Famille XLI Cerambycides. *Naturaliste Canadien*, 63, 210–210; 241–251; 64, 22–30; 101–117.
- Chagnon, G. (1938) Contribution à l'étude des Coleoptères de la Province de Québec. *Naturaliste Canadien*, 65(5), 257–320.
- Chagnon, G. (1939) A preliminary list of the insects collected in the Laurentide Provincial Park, Montmorency County, Quebec. *70th Annual Report of the Entomological Society of Ontario*, 1939, 83–87.
- Chagnon, G. & Robert, A. (1962) *Principaux coléoptères de la Province de Québec*. Les Presses de l'Université de Montreal, Montreal, 440 pp.
- Champlain, A. B., (1909) Coleoptera Collecting by Gas-light. *Entomological News*, 20 (4) 179–181.
- Champlain, A.B., Kirk, H.B. & Knull, J.N. (1925) Notes on Cerambycidae. *Entomological News*, 36(4), 105–109; (5), 139–142.
- Champlain, A.B. & Knull, J.N. (1922) New North American Coleoptera. *Entomological News*, 33(5), 144–149.
- Chemsak, J.A. (1965) Notes on the habits of *Trichoderes pini* Chevrolat. *The Pan-Pacific Entomologist*, 41(2), 93–95.
- Chemsak, J.A. (1967) Notes on Cerambycidae of Grand Bahama Islands. *The Pan-Pacific Entomologist*, 43(3), 181–188.
- Chemsak, J.A. (1972) John A., 1972 A new seed inhabiting cerambycid from Costa Rica. *The Pan-Pacific Entomologist*, 48 (2), 150–152, 1 fig.
- Chemsak, J.A. (1977) A list of types of Cerambycidae at the Field Museum of Natural History, Chicago (Coleoptera). *The Coleopterists' Bulletin*, 31(2), 173–179.
- Chemsak, J. A. (1996) *Illustrated revision of the Cerambycidae of North America. Parandrinae, Spondylidinae, Aseminae, Prioninae*. Wolfsgarden Books, Burbank, 1, 150 pp.
- Chemsak, J.A. & Linsley, E.G. (1970) Additional designations of lectotypes of neotropical Cerambycidae in the collections of the British Museum (Natural History) (Coleoptera). *Journal of the Kansas Entomological Society*, 43(4), 404–417.
- Chemsak, J.A. & Linsley, E.G. (1977) Review of the species of *Derobrachus* Audinet-Serville occurring in the United States (Coleoptera: Cerambycidae). *Journal of the Kansas Entomological Society*, 50(4), 508–514.
- Chemsak, J.A. & Linsley, E.G. (1978) Additions to the Cerambycidae of the Revilla Gigedo Islands (Coleoptera). *Bulletin of the Southern California Academy of Sciences*, 77(3), 138–142.
- Chemsak, J.A. & Linsley, E.G. & Mankins, J.V. (1980) Records of some Cerambycidae from Honduras (Coleoptera). *The Pan-Pacific Entomologist*, 56(1), 26
- Chemsak, J.A. & Linsley, E.G. & Noguera, F.A. (1992) *Listados faunísticos de Mexico. II. Los Cerambycidae y Disteniidae de Norteamérica, Centroamérica y las Indias Occidentales (Coleoptera)*. Universidad Nacional Autónoma, Mexico, D. F., 204 pp.

- Chemsak, J.A. & Noguera, F.A. (1995) Annotated checklist of the Cerambycidae of the Estación de Biología Chamela, Jalisco, Mexico (Coleoptera), with descriptions of a new genera and species. *Folia Entomologica Mexicana*, 89, 55–102.
- Chenu, J.C. (1870) *Encyclopédie d'histoire naturelle, ou traité complet de cette science d'après les travaux des naturalistes les plus éminents de tous les pays et de tous les époques; Buffon, Daubenton, Lacépède, G. Cuvier, F. Cuvier, Geoffroy Saint-Hilaire, Latreille, de Jussieu, Brongniart, etc., etc. Ouvrage résumant les observations des auteurs anciens et comprenant toutes les découvertes modernes jusqu'à nos jours*. Firmin-Didot, Paris, Coléoptères, 3, 360 pp., (2e. ed.).
- Chevrolat, L.A. In: D'Orbigny, C.V.D. (1844) *Dictionnaire universel d'histoire naturelle résumant et complétant tous les faits présentés par les encyclopédies, les anciens dictionnaires scientifiques, les oeuvres complètes de Buffon, et les meilleurs traités spéciaux sur les diverses branches des sciences naturelles; donnant la description des êtres et des divers phénomènes de la nature, l'étymologie et la définition des noms scientifiques, les principales applications des corps organiques et inorganiques, à l'agriculture, à la médecine, aux arts industriels, etc.; dirigé par M. Charles d'Orbigny, et enrichi d'un magnifique atlas de planches gravées sur acier*. Langlois & Leclercq, Fortin, Masson & Cie., Paris, 4, 1–752 (livraison 43-49, Evenhuis, 1990)
- Chevrolat, L.A. In: D'Orbigny, C.V.D. (1846) *Dictionnaire universel d'histoire naturelle résumant et complétant tous les faits présentés par les encyclopédies, les anciens dictionnaires scientifiques, les oeuvres complètes de Buffon, et les meilleurs traités spéciaux sur les diverses branches des sciences naturelles; donnant la description des êtres et des divers phénomènes de la nature, l'étymologie et la définition des noms scientifiques, les principales applications des corps organiques et inorganiques, à l'agriculture, à la médecine, aux arts industriels, etc.; dirigé par M. Charles d'Orbigny, et enrichi d'un magnifique atlas de planches gravées sur acier*. Langlois & Leclercq, Fortin, Masson & Cie., Paris, 7, 1-868.
- Chevrolat, L.A. In: D'Orbigny, C.V.D. (1847) *Dictionnaire universel d'histoire naturelle résumant et complétant tous les faits présentés par les encyclopédies, les anciens dictionnaires scientifiques, les oeuvres complètes de Buffon, et les meilleurs traités spéciaux sur les diverses branches des sciences naturelles; donnant la description des êtres et des divers phénomènes de la nature, l'étymologie et la définition des noms scientifiques, les principales applications des corps organiques et inorganiques, à l'agriculture, à la médecine, aux arts industriels, etc.; dirigé par M. Charles d'Orbigny, et enrichi d'un magnifique atlas de planches gravées sur acier*. Langlois & Leclercq, Fortin, Masson & Cie., Paris, 10, 1-760.
- Chevrolat, L.A. In: D'Orbigny, C.V.D. (1849) *Dictionnaire universel d'histoire naturelle résumant et complétant tous les faits présentés par les encyclopédies, les anciens dictionnaires scientifiques, les oeuvres complètes de Buffon, et les meilleurs traités spéciaux sur les diverses branches des sciences naturelles; donnant la description des êtres et des divers phénomènes de la nature, l'étymologie et la définition des noms scientifiques, les principales applications des corps organiques et inorganiques, à l'agriculture, à la médecine, aux arts industriels, etc.; dirigé par M. Charles d'Orbigny, et enrichi d'un magnifique atlas de planches gravées sur acier*. Langlois & Leclercq, Fortin, Masson & Cie., Paris, 12, 1-816.
- Chevrolat, L.A. (1852) Rappel des coléoptères décrits par Palisot de Beauvois aux genres actuellement adoptés, avec synonymie. *Annales de la Société Entomologique de France*, (2)10, 629–651.
- Chevrolat, L.A. (1862) Coléoptères de l'Ile de Cuba. Notes, synonymies et descriptions d'espèces nouvelles. Familles des cérambycides et des parandrides. *Annales de la Société Entomologique de France*, (4)2, 245–280.
- Chittenden, F.H. (1894) On the habits of some longicorns. *Proceedings of the Entomological Society of Washington* 3 (2), 95-102.
- Clark, M.E. (1956) An annotated list of the Coleoptera taken at or near Terrace, British Columbia, part III. *Proceedings of the Entomological Society of British Columbia*, 52, 39–43.

- Cockerell, T.D.A. (1890) Notes on the insect fauna of high altitudes in Custer County, Colorado. *The Canadian Entomologist*, 22(2), 37–39.
- Cope, J.S. (1984) Notes on the ecology of western Cerambycidae. *The Coleopterists Bulletin*, 38(1), 27–36.
- Costello, S.L., Jacobi W.R. & Negrón, J.F. (2013) Emergence of Buprestidae, Cerambycidae, and Scolytinae (Coleoptera) from Mountain Pine Beetle-Killed and Fire-Killed Ponderosa Pines in the Black Hills, South Dakota, USA *The Coleopterists' Bulletin* 67 (2), 149-154, 1 fig.
- Craighead, F.C. (1915) Larvae of the Prioninae. *Report of the Official Secretary*, United States Department of Agriculture, 107, 1–24.
- Craighead, F.C. (1923) North American cerambycid larvae. *Bulletin of the Canada Department of Agriculture*, (n.s.) 27, 1–239.
- Craighead, F.C. (1950) Insect enemies of eastern forests. *Miscellaneous Publications*, United States Department of Agriculture, 657, 1–679.
- Craighead, F. C. & Middleton, W. 1930 An annotated list of the important North American forest insects. *USDA Miscellaneous Publications*, 74, 1-30.
- Crawford, R.F. & Eyer, J.R. (1928) The Giant Apple Root Borer. *Bulletin of the New Mexico Agricultural Experiment Station*, 168, 1–8.
- Crawford, R. F. & Eyer, J. R. (1928) The Giant Apple Root Borer. *Bulletin of the New Mexico Agricultural Experiment Station* 168, 1-8
- Crawford, R.F. & Eyer, J.R. (1942) Life history and control of the giant apple root borer. *Bulletin of the New Mexico Agricultural Experiment Station*, 295, 1–14.
- Daggett, F.S. (1901) Habits of *Prionus californicus* Mots. *Entomological News*, 12(10), 319.
- Dalman, J.W. (1817) In: Schoenherr, C. J. *Synonymia insectorum, oder: Versuch einer synonymie aller bisher bekannten Insecten; nach Fabricii Systema Eleutheratorum & c. geordnet*. Eleutherata oder Käfer. *Lewerentzischen Buch.*, Skara, 1(3), 506 pp.
- Damoiseau, R. & Cools, J. (1987) Liste du matériel typique dans les collections entomologiques de l'Institut royal des Sciences naturelles de Belgique. Coleoptera Cerambycoidea Cerambycidae : Aseminae, Cerambycinae, Disteniinae, Lepturinae, Parandrinae, Prioninae et Spondyliinae *Documents de Travail de L'Institut Royal des Sciences Naturelles de Belgique* 42: 1-39.
- Davis, A.C. (1932) A list of Coleoptera of Ft. Tejon, California. *Bulletin of Southern California Academy of Sciences*, 31(3), 75–87.
- Davis, W. T. (1909) *Proceedings of the New York Entomological Society*. Meeting of October 6, 1908. *Journal of the New York Entomological Society* 17, 41-43.
- Davis, W.T. & Leng, C.W. (1912) Insects on a recently felled tree. *Journal of the New York Entomological Society*, 20(2), 119–121.
- Dean, H.A. (1953) Long-horned beetles that attack citrus in the Lower Rio Grande Valley of Texas. *Journal of Economic Entomology*, 46(1), 174.
- Degeer, C. (1775) *Mémoires pour servir à l'histoire des insectes*. Pierre Hesselberg, Stockholm, 4, 455 pp.
- Dejean, P.F.M.A. (1835) *Catalogue des Coléoptères de la collection de M. le comte Dejean*. Crevot, Paris, 2 ed., livr. 4, 257–360.
- DeLeon, D. (1934) An annotated list of the parasites, predators, and other associated fauna of the mountain pine beetle in western white pine and lodge-pole pine. *The Canadian Entomologist*, 66(3), 51–61.
- DeLeon, D. (1952) Insects associated with *Sequoia sempervirens* and *Sequoia gigantea* in California. *The Pan-Pacific Entomologist*, 23(2), 75–91.
- Desmarest, E. (1860) *Coléoptères buprestiens, scarabéiens, pimeliens, curculioniens, scolytiens, chrysoméliens, etc.* In Chenu, J. G., *Encyclopédie d'histoire naturelle ou traité complet de cette science d'après les travaux des naturalistes les plus éminents de tous les pays et de toutes les époques; Buffon, Daubenton, Lacépède, G. Cuvier, F. Cuvier, Geoffroy Saint-Hilaire, Latreille, de Jussieu, Brongniart, etc., etc. Ouvrage résumant les observations des auteurs anciens et comprenant toutes les découvertes modernes jusqu'à nos jours. Troisième partie.* Marescq et Compagnie, Paris, [3] + 360pp. + 48 pls.

<https://doi.org/10.5962/bhl.title.1959>

- Desmarest, E. in Chenu, J.C. (1870) *Encyclopédie d'histoire naturelle, ou traité complet de cette science d'après les travaux des naturalistes les plus éminents de tous les pays et de tous les époques; Buffon, Daubenton, Lacépède, G. Cuvier, F. Cuvier, Geoffroy Saint-Hilaire, Latreille, de Jussieu, Brongniart, etc., etc. Ouvrage résumant les observations des auteurs anciens et comprenant toutes les découvertes modernes jusqu'à nos jours*. Firmin-Didot, Paris, Coléoptères, 3, 360 pp., (2e. ed.).
- Devesa Regueiro S., Fonseca E. & Barro A. (2015b) Longicornios de Cuba (Coleoptera: Cerambycidae). Vol. 1. Parandrinae, Prioninae, Spondylidinae, Cerambycinae. Greta Editores, Barcelona, 1: 1-353.
- DiGirolamo, M.F & Doods, K.J. (2017) Comparison of the species richness and abundance of Cerambycidae (Coleoptera) and Scolytinae (Coleoptera: Curculionidae) captured in aerial malaise traps with and without a bottom collector. *The Canadian Entomologist*, 149 (3), 408-412.
- Dillon, E.S. & Dillon, L.S. (1961) *A manual of common beetles of eastern North America*. Row & Peterson, Evanston, 1, 434 pp.
- Dimmock, G. (1884) Sexual attraction in *Prionus*. *Psyche*, 4(120), 159.
- Doane, R.W., Van Dyke, E.C., Chamberlin, W.J. & Burke, H.E. (1936) *Forest Insects*. McGraw-Hill, New York & London, 463 pp.
- Dow, R.P. (1913) Rare or new Coleoptera from California. *Bulletin of the Brooklyn Entomological Society*, 8(5), 77–79.
- Dozier, H.L. (1918) An annotated list of Gainesville, Florida, Coleoptera. *Entomological News*, 29(9), 331–335.
- Dozier, H.L. (1920) An ecological study of hammock and piney woods insects in Florida. *Annals of the Entomological Society of America*, 13 (4), 325-380
- Drapiez, M. (1844) *Dictionnaire classique des Sciences Naturelles présentant la définition, l'analyse et l'histoire de tous les êtres qui component les trois règnes, leur application générale aux arts, à l'agriculture, à la médecine, à l'économie domestique, etc.; resumant tous les faits présentés par les dictionnaires d'histoire naturelle, augmenté des nombreuses découvertes acquises depuis la publication de ces ouvrages*. Meline, Cans et Compagnie, Bruxelles, 1839, 9, 704 pp.
- Drapiez, M. (1845) *Dictionnaire classique des Sciences Naturelles présentant la définition, l'analyse et l'histoire de tous les êtres qui component les trois règnes, leur application générale aux arts, à l'agriculture, à la médecine, à l'économie domestique, etc.; resumant tous les faits présentés par les dictionnaires d'histoire naturelle, augmenté des nombreuses découvertes acquises depuis la publication de ces ouvrages*. Meline, Cans et Compagnie, Bruxelles, 10, 747 pp., 200 pls
- Drumont A., Komiya Z. (2010) Family Cerambycidae. Subfamily Prioninae Latreille, 1802 [pp. 86–95]. In: Löbl I., Smetana A. (Eds) *Catalogue of Palaearctic Coleoptera*. Volume 6. Chrysomeloidea. Apollo Books, Stenstrup, 924 pp.
- Drury, D. (1773) *Illustrations of Natural History, wherein are exhibited upwards of two hundred and twenty figures of exotic insects according to their different genera; very few of which have hitherto been figured by any author, being engraved and coloured from nature, with the greatest accuracy, and under the author's own inspection, on fifty copper-plates. With a particular description of each insect: interspersed with remarks and reflections on the nature and properties of many of them*. D. Drury, London, 2: vii + 90 pp, 50 pls.
- Drury, D. & Westwood, J.O. (1837) *Illustrations of exotic entomology, containing upwards of six hundred and fifty figures and descriptions of foreign insects, interspread with remarks and reflections on their nature and properties*. A new edition, brought down to the present state of the science, with the systematic characters of each species, synonyms, indexes, and other additional matter. Henry G. Bohn, London, 1, 151 pls
- Duffy, E.A.J. (1953) *A monograph of the immature stages of British and imported timber beetles (Cerambycidae)*. British Museum (Natural History), London, 350 pp.
- Duffy, E.A.J. (1960) *A monograph of the immature stages of Neotropical timber beetles (Cerambycidae)*. British Museum (Natural History), London, 327 pp.

- Duges, E.D. (1884) Métamorphoses du *Mallodon angustatum* Thoms. *Annales de la Société Entomologique de Belgique*, 28: 12-87, pl. II, figs 1-15.
- Dury, C. (1902) A revised list of the Coleoptera observed near Cincinnati, Ohio, with notes on localities, bibliographical references and descriptions of six new species. *Journal of the Cincinnati Society of Natural History*, 20, 107–196.
- Eaton, C.B. (1959) Observations on the survival of *Arhopalus productus* (LeConte) larvae in Douglas fir lumber (Coleoptera, Cerambycidae). *The Pan-Pacific Entomologist*, 35(2), 114–116.
- Easterling G. R. (1934) A study of the insect fauna of a coniferous reforestation area in Southeastern Ohio *The Ohio Journal of Science*, 34 (3), 129-146.
- Eaton, C.B. & Lyon, R.L. (1955) *Arhopalus productus* (LeC.), a borer in new buildings. *Technical Papers of the California Forest Range Experiment Station*, 11, 1–11.
- Ehrmann E. C. (1897) Coleopterological notes from my brother's diary. *Entomological News*, 8 (7), 168-170.
- Ehrmann, G. A. (1900) Notes on Coleoptera *Entomological News*, 11 (10), 619-621.
- Emmons E., (1854) Agriculture of New York ; comprising an account of the classification, composition and distribution of the soils... ; together with descriptions of the more common and injurious species of insects The Natural History of New York. *Agricultural Report 5*: 1-276, 47 pls. Albany, C. Van Benthuysen
- Erichson, W.F. (1849) Insecten. Pp. 553–617 in: *Reisen in Britisch-Guiana in den Jahren 1840–1844. Im Auftrag Sr. Majestat des Königs von Preussen ausgeführt von Richard Schomburgk. Nebst einer Fauna und Flora Guiana's nach Vorlagen von Johannes Müller, Ehrenberg, Erichson, Klotzsch, Troschel, Cabanis und Andern. Mit Abbildungen und einer Karte von Britisch-Guiana aufgenommen von Sir Robert Schomburgk. Dritter Theil.* J.J. Weber, Leipzig. viii + [1] + pp. 533–1260.
- Essig, E.O. (1915) Injurious and beneficial insects of California. *Supplement of the Monthly Bulletin of the California Commission of Horticulture*, 4(4), 541 pp.
<https://doi.org/10.5962/bhl.title.9364>
- Essig, E.O. (1926) *Insects of Western North America*. MacMillan, New York, 1035 pp.
- Essig, E.O. (1958) *Insects and mites of Western North America*. MacMillan, New York, 1050 pp.
- Evans, J. D. (1895) The insect fauna of the Sudbury District, Ontario. *The Canadian Entomologist*, 27 (7), 173-175.
- Evans, J.D. (1906) List of Coleoptera in the collection of J.D. Evans, Trenton, Ont., which have not heretofore been recorded as having been taken in Canada. *The Canadian Entomologist*, 38 (3). 96-100.
- Eyer, J.R. 1942. Life history and control of the giant apple root borer (*Prionus californicus* Mots.). *Bulletin of the New Mexico Agricultural Experiment Station*, 295, 1–14.
- Fabricius, J.C. (1775) *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. Korte, Flensburgi et Lipsiae, 832 pp.
<https://doi.org/10.5962/bhl.title.36510>
- Fabricius, J.C. (1776) *Genera insectorum eorumque characteres naturales secundum numerum figuram, situm et proportionem omnium partium oris adiecta mantissa specierum nuper detectarum*. Bartsch, Chilonii, 310 pp.
<https://doi.org/10.5962/bhl.title.65743>
- Fabricius, J.C. (1781) *Species insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus*. Bohn, Hamburgi et Kilonii, 1, 552 pp.
<https://doi.org/10.5962/bhl.title.11658>
- Fabricius, J.C. (1787) *Mantissa insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus*. Proft, Hafniae, 1, 348 pp.
<https://doi.org/10.5962/bhl.title.36471>

- Fabricius, J.C. (1793) *Entomologia systematica, emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus*. Proft, Hafniae, 1, 538 pp.
<https://doi.org/10.5962/bhl.title.36516>
- Fabricius, J.C. (1794) *Entomologia systematica, emendata et aucta*. Proft, Hafniae, 1, 472 pp.
<https://doi.org/10.5962/bhl.title.36516>
- Fabricius, J.C. (1798) *Supplementum entomologia systematica*. Proft & Storch, Hafniae, 572pp.
- Fabricius, J.C. (1801) *Systema eleutheratorum secundum ordines, genera, species: adiectis synonymis, locis, observationibus, descriptionibus*. Bibliopoli Academici Novi, Kiliae, 2, 687 pp
- Fairmaire, L.M.H. (1868) In P. N. Jacquelin du Val. *Genera des Coléoptères d'Europe, comprenant leur classification en famille naturelle*
 Paris 4, 1-295, 376 figs, 78 pls.
- Fall, H.C. (1901) List of the Coleoptera of southern California with notes on habits and descriptions of new species *Occasional Papers of the California Academy of Sciences*, 8, 1-282.
- Fall H.C. (1905) New species of Coleoptera, chiefly from the Southwest. *The Canadian Entomologist*, 37 (8), 270-276.
- Fall, H.C. (1907) Descriptions of new species. *Transactions of the American Entomological Society*, 33, 218–272.
- Fall, H.C. (1912) A new *Tetropium*, two new Bruchides, with brief notes on other Coleoptera. *Entomological News*, 23, 320–323.
- Fall, H. C. (1926) A list of the Coleoptera taken in Alaska and adjacent parts of the Yukon territory in the summer of 1924. *The Pan-Pacific Entomologist*, 2 (4), 191-208
- Fall, H.C. & Cockerell, T.D.A. (1907) The Coleoptera of New Mexico. *Transactions of the American Entomological Society*, 33, 145–272.
- Farrar, R.J. & Kerr, T.W. (1968) A preliminary study of the life history of the Broad-Necked Root Borer in Rhode Island *Journal of Economic Entomology* 61 (2), 563-564.
- Fattig, P.W. (1947) The Cerambycidae or long-horned beetles of Georgia. *Emory University Museum Bulletin*, 5, 1–48.
- Ferreira, M.C. & Veiga Ferreira, G. (1959) Catálogo dos Cerambycídeos da Região Etiópica. I Parte - Supertribos Parandrina & Prionina *Memórias do Instituto de Investigação científica de Moçambique* 1: 1-76.
- Ferrer J., Barclay M,V.L. & Hancock, E.G (2004) Discovery of lost J.C. Fabricius (1775) and A.G. Olivier (1795) types of Coleoptera in the Hunterian Museum, University of Glasgow, Scotland *Annales Zoologici*, 54 (4), 775-781.
- Finlayson, T. (1969) Final-instar larvae of two hymenopterous parasites of a wood-boring beetle, *Tetropium velutinum* LeConte (Coleoptera: Cerambycidae). *Journal of the Entomological Society of British Columbia*, 66, 62–65.
- Fisher, W.S. (1915) One new genus and two new species of Cerambycidae. *Proceedings of the Entomological Society of Washington*, 17, 77–79.
- Fisher, W.S. (1944) Cerambycidae (Coleoptera) of Caripito, Venezuela. *Zoologica*, 29(1), 3–12.
- Fisher, W.S. & Kirk, H.B. (1912) Cerambycidae of Harrisburg, Pennsylvania and vicinity with notes. *Entomological News*, 23, 308–316.
- Fitch, A. (1858) Fourth report on the noxious and other insects of the State of New York. Insects infesting evergreen forest trees. [1857] 17, 687-753.
- Fletcher, J. (1926) A preliminary biological survey of the Lloyd-Cornell Reservation. *Bulletin of the Lloyd Library* 5 (27), 1-221.
- Forbes S.A. (1894) A monograph of insect injuries to Indian corn. Part I Eighteenth Report of the State Entomologist on the Noxious and Beneficial Insects of the State of Illinois, 1-149, 15 pls.
- Fox, R.C. (1975) A case of longevity of the brown prionid, *Orthosoma brunneum* (Forster) (Cerambycidae: Coleoptera). *Proceedings of the Entomological Society of Washington* 77 (2). 237.

- Fragoso, S.A. & Monné, M.A. (1982) Notas sobre Prioninae neotropicais (Coleoptera, Cerambycidae). *Pesquisa Agropecuaria Brasileira*, 17(4), 519–523.
- Fragoso, S.A. & Monné, M.A. (1995a) Notes on Macrotomini (Coleoptera, Cerambycidae, Prioninae). *Revista Brasileira de Biologia*, 55(2), 215–225.
- Frankie, G.W. & Jensen, G.L. (1971) Studies on the distribution and biology of *Atimia helenae* Linsley on two California *Cupressus* species (Coleoptera, Cerambycidae). *The Pan-Pacific Entomologist*, 47(4), 287–293.
- Franz, E. (1954) Cerambycidae (Ins., Col.) aus El Salvador. *Senckenbergiana*, 34(4-6), 213–229.
- Frost, C.A. (1912) Collecting Coleoptera in a Maine sawmill Yard *The Canadian Entomologist*, 44, 304-308.
- Frost, C.A. (1915) Remarks on collecting at light, with a list of the Coleoptera taken *Psyche*. 22, 207-211.
- Frost, S.W. (1969) Supplement to Florida insects taken in light traps *Florida Entomologist*, 52 (2), 91-101.
- Furniss, R.L. & Carolin, V.M. (1977) Western Forest Insects. *Miscellaneous Publications*, United States Department of Agriculture, 1339, 654 pp.
- Fyles, T.W. (1896) How the forest in the District of Bedford was swept away. *26th Annual Report of the Entomological Society of the Province of Ontario* 1895, 21-25.
- Gahan, C.J. (1890) Notes on some West Indian longicorn Coleoptera, with descriptions of new genera and species. *The Annals and Magazine of Natural History*, (6)6, 23–34.
- Gahan, A.B. (1911) Some notes on *Parandra brunnea*. *Journal of Economic Entomology* 4, 299-301.
- Gahan, C.J. (1895a) On the longicorn Coleoptera of the West Indian islands. *The Transactions of the Entomological Society of London*, 1895, 79–140.
- Gahan, C.J. (1906) Coleoptera (Cerambycidae). In: *The Fauna of British India, including Ceylon and Burma*. C. T. Bingham, London, 329 pp.
- Gahan, C.J. (1908) Notes on North American Longicornia, with descriptions of some new species. *The Annals and Magazine of Natural History*, (8)1, 140–145.
- Galileo, M.H.M. (1987a) Sistemática das tribos Meroscelisini e Anacolini (Coleoptera, Cerambycidae, Prioninae) nas Américas. I. Meroscelisini. *Revista Brasileira de Entomologia*, 31(2), 141–267.
- García Morales, L.J., García Jiménez, J., Toledo, V.H. & Cantú Ayala, C.M. (2015) Lista anotada preliminar de los Cerambycidae (Coleoptera) de Tamaulipas, México. *Biodiversidade Tamaulipeca*, Instituto Tecnológico de Ciudad Victoria, 2(2), 97-115.
- Gardiner, L.M. (1957) Deterioration of fire-killed pine in Ontario and the causal wood-boring beetles. *The Canadian Entomologist*, 89 (6), 241-263, 15 figs
- Gardiner L.M. (1970) Immature stages and habits of *Spondylis upiformis* Mannerheim (Coleoptera: Cerambycidae). *The Pan-Pacific Entomologist*, 46 (1), 33-36.
- Garnett, R.T. (1918) An annotated list of the Cerambycidae of California (Coleoptera). *The Canadian Entomologist*, 50(5), 172–177; (6), 205–213; (8), 248–353; (9), 281–284.
- Gemminger, M. (1872) [Homonymy of *Malodon gnatho*]. *Coleopterologische Hefte*, 9-10, 254.
- Gibson, A. (1917) The Entomological Record, 1916. *47th Annual Report of the Entomological Society of Ontario*, 1916, 137–171.
- Gilmour, E.F. (1954) Notes on a collection of Prionidae (Col., Cerambycidae) from the Institut Royal des Sciences Naturelles de Belgique. *Bulletin du Institut de Sciences Naturelles de Belgique*, 30(24), 1–48.
- Gilmour, E.F. (1956) Revision of the Prioninae of tropical and South Africa. *Longicornia*, 3, 1–252.
- Gilmour, E.F. (1965) Some new species of *Hovatoma* Lameere (Coleoptera, Cerambycidae, Prioninae, Macrotomini) *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 41 (37), 1-15, 1 pl.
- Girard, M. (1873) *Les insectes. Traité élémentaire d'entomologie comprenant l'histoire des espèces utiles et de leurs produits, des espèces nuisibles et des moyens de les détruire, l'étude des métamorphoses et des mœurs, les procédés de chasse et de conservation. Avec planches coloriées. Introduction -Coléoptères*. J.-B. Baillièere et Fils, Paris. viii + 840 pp.

- Gistel, J.N.F. (1848a) Faunula monacensis cantharologica. *Isis von Oken* 1848: (6) [unn. pp. "1–3"]; (7): [unn. pp. "4–6"]; (8): [unn. p. "7"]; (9): [unn. pp. "8–9"]; (10): [unn. pp. "10–11"]; (11): [unn. pp. «12–13»] (inserted with fascicles).
- Gistel, J.N.F. (1848b) *Naturgeschichte der Thierreiche für höhere Schulen*. Hoffmann, Stuttgart, 8(1), xvi + 216 pp., 30 pls.
- Gmelin, J.F. (1790) *Caroli a Linnaeus Systema Naturae*. Ed. 13. Beer, Lipsiae, 1(4), 1517–2224.
- Goczal, J., Rossa, R., Sweeney, J. & Tofilski, A. (2017) Citizen monitoring of invasive species: wing morphometry as a tool for detection of alien *Tetropium* species *Journal of Applied Entomology* 141 (6), 496–506, 5 figs.
- Goldman, E.H. (1933) Comparisons of the mouthparts of adult longhorn beetles with reference to their food (Coleoptera: Cerambycidae). *Transactions of the American Entomological Society*, 59, 85–102.
- Gosling, D.C.L. (1973) An annotated list of the Cerambycidae of Michigan (Coleoptera). Part I, Introduction and the subfamilies Parandrinae, Prioninae, Spondyliinae, Aseminae and Cerambycinae. *Great Lakes Entomologist*, 6(3), 65–84.
- Gosling D. C. L. (1984) Cerambycid host plants in a southwestern Michigan woodland (Coleoptera: Cerambycidae). *The Great Lakes Entomologist*, 17 (2), 69–78.
- Gosling D. C. L. (1986) Ecology of the Cerambycidae (Coleoptera) of the Huron Mountains in Northern Michigan. *The Great Lakes Entomologist*, 19 (3), 153–162.
- Gounelle, P.E. (1911) Pierre Émile, 1911 Communications. Note sur *Halycidocrius Philippii* Berg et *Prionapterus staphylinus* Serv. [Col. Cerambycidae] (avec la planche no II) *Bulletin de la Société Entomologique de France*, (16), 319–327, 2 pls.
- Gowdey, C.C. (1926) Catalogus insectorum Jamaicensis. *Entomological Bulletin of the Department of Agricultura of Jamaica*, 4(1), 114 pp.; (2), 10 pp.
- Gray, G. (1832) In: Griffith, E. & Pidgeon, E. *The animal kingdom arranged in conformity with its organization by the Baron Cuvier, with supplementary addition to each order*. Insects. London, 2, 796 pp., (1831: pls. 6, 14, 65, 73; 1832: pls. 85, 88, 95, 100, 119).
- Greene, G.M. (1918) 1A rare Coleoptera paper of T. W. Harris *Transactions of the American Entomological Society*, 44, 251–261
- Griffith, H.G. (1900) Coleopterous fauna of Phoenix, Arizona and surrounding regions. *Entomological News*, 11(8), 561–570
- Grossbeck, J.A. (1912) List of insects collected by the "Albatross" expedition Lower California in 1911, with description of a new species of wasp. *Bulletin of the American Museum of Natural History*, 31, 323–326.
- Gutiérrez, N. & Noguera, F.A. (2015) New distributional records of Cerambycidae (Coleoptera) from Mexico *The Pan-Pacific Entomologist*, 91 (2), 135–147.
- Gwynne, D. T. & Hostetler, B. B. (1978) nMass emergence of *Prionus emarginatus* (Say) (Coleoptera: Cerambycidae) *The Coleopterists' Bulletin* 32 (4), 347–348.
- Haack, R.A. (2017) Feeding Biology of Cerambycids In Qiao Wang, *Cerambycidae of the World : Biology and Pest Management* (3): 105–132, CRC Press
- Haack R.A., Keena M.A. & Eyre, D. (2017) Life History and Population Dynamics of Cerambycids. In Qiao Wang, *Cerambycidae of the World : Biology and Pest Management* (2): 71–103, 11 figs. CRC Press
- Haldeman, S.S. (1847a) Material towards a history of the Coleoptera Longicornia of the United States. *Transactions of the American Philosophical Society*, 10, 27–66.
<https://doi.org/10.2307/1005263>
- Haldeman, S.S. (1847b) Corrections and additions to his paper on the Longicornia of the United States. *Proceedings of the American Philosophical Society*, 4, 371–376.
- Haldeman S. S. (1848) On several new genera and species of insects. *Proceedings of the Academy of Natural Sciences of Philadelphia* 3 [1846] 124–128
- Hamilton, J. (1886) Natural History notes on Coleoptera. No. 2. *The Canadian Entomologist*, 18 (6), 111–115.
- Hamilton, J. (1889) Catalogue of the Coleoptera common to North America, northern Asia and Europe, with the distribution and bibliography. *Transactions of the American Entomological Society*, 16, 88–162.

- Hamilton, J. (1890) Proposed corrections of specific names to harmonize Mr. Henshaw's Catalogue of the North American Coleoptera, with the generally accepted European nomenclature, with relation to the species common to the two continents. *Entomologica Americana*, 6 (3), 41-44.
- Hamilton, J. (1892) Notes on Coleoptera - No. XI. *The Canadian Entomologist*, 24 (12), 293-300
- Hamilton, J. (1894) Catalogue of the Coleoptera of Alaska, with the synonymy and distribution. *Transactions of the American Entomological Society*, 21, 1-38.
- Hamilton, J. (1894) Catalogue of the Coleoptera common to North America, northern Asia and Europe, with the distribution and bibliography (2nd edition). *Transactions of the American Entomological Society*, 21, 345-416.
- Hamilton, J. (1895) Catalogue of the Coleoptera of southwestern Pennsylvania, with notes and descriptions. *Transactions of the American Entomological Society*, 22, 317-381.
- Hanks, L.M. & Wang, Q. (2017) Reproductive Biology of Cerambycids. In Qiao Wang, Cerambycidae of the World: Biology and Pest Management (4): 133-159. CRC Press
- Hardy, G.A. (1926) Cerambycidae of Vancouver Island (Preliminary annotated list). *Report of the Museum of Natural History of British Columbia*, 1925, 24-33.
- Hardy, G.A. (1927) Buprestidae and Cerambycidae of Vancouver Island. *Report of the Provincial Museum of Natural History of British Columbia*, 1926, C32-C37.
- Hardy, G.A. (1942) Notes on some wood-boring beetles of Saanich, Vancouver Island, B. C. (Coleoptera: Cerambycidae and Buprestidae). *Proceedings of the Entomological Society of British Columbia*, 39, 9-13.
- Hardy, G.A. (1944) Further notes on the Cerambycidae of Vancouver Island (Coleoptera). *Proceedings of the Entomological Society of British Columbia*, 41, 15-18.
- Hardy, G.A. (1945) Some longhorn beetles of Vancouver Island. *Victoria Naturalist*, 2, 88-92.
- Hardy, G.A. (1948) Some beetles of the families Cerambycidae and Buprestidae from Manning Park, British Columbia. *Proceedings of the Entomological Society of British Columbia*, 44, 31-34.
- Hardy, G.A. (1955) The natural history of the Forbidden Plateau area, Vancouver Island, British Columbia. *Report of the Provincial Museum of Natural History of British Columbia*, 1954, B24-B63.
- Hardy, G.A. (1957) Notes on the flora and fauna of the Blenkinshop Lake area on southern Vancouver Island, British Columbia. *Report of the Provincial Museum of Natural History of British Columbia*, 1956, D25-D66.
- Hardy, G.A. & Preece, W.H.A. (1926) Notes on some species of Cerambycidae from the southern portion of Vancouver Island. *The Pan-Pacific Entomologist*, 3(1), 34-40.
- Hardy, G.A. & Preece, W.H.A. (1927a) Further notes on some species of Cerambycidae (Col.) from the southern portion of Vancouver Island, B. C., with descriptions of some new varieties. *The Pan-Pacific Entomologist*, 3(4), 187-193.
- Hardy, G.A. & Preece, W.H.A. (1927b) Additional notes on some Cerambycidae of Vancouver Island, B. C. *The Pan-Pacific Entomologist*, 4(2), 61-67.
- Harrington, W.H. (1881) On some Coleoptera injurious to our pines. *Transactions of the Ottawa Field Naturalists' Club* 2, 28-33.
- Harrington, W.H. (1887) Insects infesting maple trees. *17th Annual Report of the Entomological Society of the Province of Ontario 1886*, 22-33
- Harrington, W.H. (1899) A few Canadian longicorns. *The Canadian Entomologist*, 31(5), 107-108.
- Harrington W.H. (1890) On the lists of Coleoptera published by the Geological Survey of Canada, 1842-1888. *The Canadian Entomologist*, 22 (9), 184-191.]
- Harris, T.W. (1837) Characteristics of some previously described North American Coleopterous Insects, and descriptions of others which appear to be new in the Collection of Mr. Abraham Halsey *Transactions of the Hartford Natural History Society* 1 [1836], 65-91.
- Harris. T.W. (1838) Habits of some of the insects injurious to vegetation in Massachusetts Reports of the Commissioners on the Zoological Survey of the State 1838: 57-104. Boston, Dutton & Wentworth, State printers.

- Harris, T.W.(1841) A report on the insects of Massachusetts injurious to vegetation Cambridge, Massachusetts 459 pp.
- Harris, T.W. (1852) A treatise on some of the insects of New England which are injurious to vegetation Boston. White & Potter printers. Second edition : 1-513.
- Hatch, M.H. (1925) A list of Coleoptera from Charlevoix County, Michigan. *Papers of the Michigan Academy of Sciences, Arts and Letters*, 4, 543–586.
- Hatch, M.H. (1930) Records and new species of Coleoptera from Oklahoma and western Arkansas, with subsidiary studies. *Publications of the University of Oklahoma Biological Survey*, 2 (1), 15-26
- Hatch, M.H. (1938) A bibliographical catalogue of the injurious arachnids and insects of Washington. *University of Washington Publications in Biology*, 1, 163-223.
- Hatch, M.H. (1971) The beetles of the Pacific Northwest. Part V: Rhipicerioidea, Sternoxi, Phytophaga, Rhynchophora and Lamellicornia. *University of Washington, Publications in Biology*, 16, 1–662.
- Hatch, M.H. & Ortenburger, A.I. (1930) Records and new species of Coleoptera from Oklahoma. *Publications of the University of Oklahoma, Biological Survey*, 2(1), 7–14.
- Hebard, M. (1903) Notes on the collecting around Thomasville, Georgia. *Entomological News*, 14(8), 260–261.
- Heffern, D.J. (1998) *Insects of Western North America. 1. A survey of the Cerambycidae (Coleoptera), or longhorned beetles, of Colorado*. Contributions of the C. P. Gillette Museum of Arthropod Diversity, Colorado State University, Fort Collins, 32 pp.
- Heffern, D.J., Drumont, A. Sama, G., Komiya, Z., Tavakilian, G.L. & Santos-Silva, A. (2006). Reversal of precedence using the International Code of Zoological Nomenclature to maintain prevailing usage of *Macrotoma* Audinet-Serville, 1832 (Coleoptera, Cerambycidae). *Bulletin de la Société Entomologique de France*, 111(1), 123-137.
- Heffern, D.J. & Santos-Silva, A. (2016) New species and new record of Cerambycidae (Coleoptera) from Mexico, with updates to type depository for two species of Disteniinae. *Zootaxa*, 4072(5), 569-578.
<http://doi.org/10.11646/zootaxa.4072.5.4>
- Heffern, D. J., Vlasák, J. & Alten R.L. (2018) Larval host plant records, distributional records, and biological information on North American Cerambycidae (Coleoptera) *The Coleopterists' Bulletin* 72 (4), 739-750.
- Henshaw, S. (1874) List of Coleoptera collected in the vicinity of Cliftondale, Mass., June 12, 1873 *Psyche*. 1 (6), 22-23.
- Herbst, J.F.W. (1799) *Natursystem aller bekannten in-und Ausländischen Insecten*. Käfer. Pauli, Berlin, 8, 1–420.
- Herrick, G. W. (1904) Insects injurious to pecans. *Bulletin of the Mississippi Agricultural Experiment Station* 86, 1-42.
- Herrick, G.W. (1935) *Insect enemies of shade-trees*. Comstock Publ., Ithaca, 417 pp
- Heyrovsky, L. (1955) Faune CSR. Svazek 5. Tesarokiviti (Cerambycidae). Ceskosl. Akad. Ved Sekce biol., Praha, 346 pp.
- Hicks, S D. (1955) Natural History Survey of Coppermine, Northwest Territories, 1951 *Canadian Field-Naturalist* 69 (4), 162-166.]
- Holland, W.J. (1888) Captures made while travelling from Winnipeg to Victoria, B. C. *The Canadian Entomologist*, 20 (5), 89-92.
- Holmes, W.C. (1869) Gigantic root-borer *The American Entomologist* 2, 51
- Holt, B. (2013) A preliminary checklist of the Cerambycidae and Disteniidae (Coleoptera) of Alabama, *The Coleopterists' Bulletin* 241-256.
- Hopkins, A. D. (1893) Catalogue of West Virginia forest and shade tree insects collected in 1890-1893, including injurious, beneficial and other insects taken on or in some part of the tree examined *Bulletin of the West Virginia University Agricultural Experiment Station* 32 (8), 171-251.
- Hopkins, A.D. (1902) Insects detrimental and destructive to forest products used for constructing material. *USDA Division of Entomology Bulletin* (n. s.) 31: 60-62.

- Hopkins, A.D. (1904) Catalogue of the exhibits of insect enemies of forests and forest product at the Louisiana Purchase Exposition, St. Louis, Mo, 1904. *Bulletin of the Division of Entomology*, United States Department of Agriculture, 48(n.s.), 1–56.
- Hopping, R. (1931) New Coleoptera from Western Canada. III. *The Canadian Entomologist*, 63(10), 233–238.
- Horn, G.H. (1861) Observations on the habits of some coleopterous larvae and pupae. *Proceedings of the Entomological Society of Philadelphia* 1 (2), 43–44.
- Horn, G.H. (1867) Notes on the habits of a few California Coleoptera. *Proceedings of the Entomological Society of Philadelphia*, 6, 289–293.
- Horn, G.H. (1873) In: Hayden, F. V. 6th Ann. Report of the U. S. Geological survey of the territories embracing portions of Montana, Idaho, Wyoming, and Utah, being a report of progress of the exploration for the year 1872. Coleoptera. Washington, p. 717.
<https://doi.org/10.1002/ardp.18782120154>
- Horn, G.H. (1875) Synonymical notes and descriptions of new species of North American Coleoptera. *Transactions of the American Entomological Society*, 5, 126–156, 8 figs
<https://doi.org/10.2307/25076300>
- Horn, G.H. (1876) Notes on the coleopterous fauna of Guadalupe Island. *Transactions of the American Entomological Society*, 5, 198–201.
<https://doi.org/10.2307/25076305>
- Horn, G.H. (1880a) List of the Coleoptera described by Mr. J. H. B. Bland with their synonymy. *Transactions of the American Entomological Society*, 8, x–xii.
- Horn, G.H. (1880b) Coleoptera collected in the Florida keys. *Transactions of the American Entomological Society*, 8, xvii.
- Horn, G.H. (1885) Synonymical notes No. 2. *Entomologica Americana*.1 (5), 88–90.
- Horn, G.H. (1886a) A review of the species described by Olivier in the "Entomologiae". *Transactions of the American Entomological Society*, 13, 135–144.
- Horn, G.H. (1886b) Notes from the Museum at Cambridge (Col.). *Transactions of the American Entomological Society*, 13, xi–xvi.
- Horn, G.H. (1889) Synonymical notes (Col.). *Entomologia Americana*, 5(10–11), 198–199, 212.
- Horn, G.H. (1890) Color variation in *Ergates spiculatus* var. *marmoratus* Ckll. *Entomological News*, 1, 161.
- Horn, G.H. (1891) New species and miscellaneous notes. *Transactions of the American Entomological Society*, 18, 32–48.
- Horn, G.H. (1894) The Coleoptera of Baja California. *Proceedings of the California Academy of Sciences*, (2)4, 302–449.
- Horn, G.H. in Leng C. W. (1884) Synopses of Coleoptera (Cerambycidae). *Bulletin of the Brooklyn Entomological Society* 7 (1), 7–11.
- Horsfall, T. (1904) Orchard enemies *Bulletin of the Missouri Fruit Experiment Station* 9, 1–31, 17 figs.
- Houghton, C.O. (1905) A List of Coleoptera. Taken on the Summit of Mt. Seward, N. Y. *Entomological News*, 16, 50–51.
- Hovore, F.T. 1981. Two new species of *Prionus* (*Homaesthis*) from the southwestern United States, with notes on other species (Coleoptera: Cerambycidae). *The Coleopterists Bulletin*, 35(4): 453–457.
- Hovore, F.T. (1983) Taxonomic and biological observations on southwestern Cerambycidae (Coleoptera). *The Coleopterists Bulletin*, 37(4), 379–387.
- Hovore, F.T. (1988) Additions to the cerambycid beetle fauna of Baja California, Mexico: records and descriptions (Coleoptera: Cerambycidae). *Wasmann Journal of Biology*, 46(1–2), 1–29
- Hovore, F.T. (2006) The Cerambycidae (Coleoptera) of Guatemala. In Enio Cano. Biodiversidad de Guatemala 1: i–vi + 1–674. Universidad del Valle de Guatemala
- Hovore, F.T. & Giesbert, E.F. (1974) Two new species of Cerambycidae from southern California (Coleoptera). *The Pan-Pacific Entomologist*, 50(2), 139–144.
- Hovore, F.T. & Giesbert, E.F. (1976) Notes on the ecology and distribution of western Cerambycidae. *The Coleopterists Bulletin*, 30(4), 349–360.

- Hovore, F.T., Penrose, R.L. & Giesbert, E.F. (1978) Notes on North American Cerambycidae (Coleoptera). *Entomological News*, 89(2–3), 95–100.
- Hovore, F.T., Penrose, R.L. & Neck, R.W. (1987) The Cerambycidae or longhorned beetles of Southern Texas: a faunal survey (Coleoptera). *Proceedings of the California Academy of Sciences*, 44(13), 283–334.
- Hovore, F.T. & Santos-Silva, A. (2004) Notes and description of Neotropical Macrotonini (Coleoptera, Cerambycidae, Prioninae). *Revista Brasileira de Entomologia*, 48(1), 49–54. <https://doi.org/10.1590/s0085-56262004000100009>
- Hovore, F. T. & Turnbow, Jr. R.H. (1984) A new species of *Prionus* (*Homaesthis*) from the Monahans sandhills of western Texas (Coleoptera: Cerambycidae) *Entomological News*, 95 (1), 1-4
- Howard, L.O. (1889) Scientific results of explorations by the U. S. Fish Commission steamer Albatross N. V. Annotated catalogue of the insects collected in 1887-88. *Proceedings of the United States National Museum*, 12 (771), 185–207.
- Howden, H.F. & Vogy, G;B. (1951) Insect communities of standing dead pine (*Pinus virginiana* Mill.) *Annals of the Entomological Society of America*, 44 (4), 581-595, 4 figs.
- Hubbard, H.G. (1886) Habits of *Hypotrachia spissipes*, the attacks of *Mallodon melanopus*. *Proceedings of the Entomological Society of Washington*, 1(1), 13.
- Hummel. A.D. (1825) Arvid David, 1825 *Essais Entomologiques N° IV. Insectes de 1825. V. Novae Species Variorum*. Essais Entomologiques 4: 1-72. St. Petersburg (Savenius)
- Hyslop, J.A. (1912) [Notes from the Pacific Northwest]. *Proceedings of the Entomological Society of Washington*, 14(2), 100–101
- Illiger, J.C.W. (1802) Aufzählung der Käfergattungen nach der Zahl der Fussglieder *Magazin für Insektenkunde*, 1, 285-305.
- Ingles, L.G. (1933) The succession of insects in tree trunks as shown by collections from various stages of decay. *Journal of Entomology and Zoology*, 25, 57–59.
- Johnson H.L. (1915) Coleoptera found in the vicinity of Meriden, Connecticut *Entomological News*, Philadelphia 26 (7): 307-319.
- Keen, F.P. (1929) Insect enemies of California pines and their control. California Department of Natural Resources, *Research Division of Forestry Bulletin*, 7, 1–113.
- Keen, F.P. (1952) Insect enemies of Western Forest. *Miscellaneous Publications*, United States Department of Agriculture, 278, 1–280. (revised edition).
- Keen, J.H. (1895) List of Coleoptera collected at Massett, Queen Charlotte Islands, B. C. *The Canadian Entomologist*, 27 (8), 217-220.
- Kempers. B.J.W. (1923) Bernet K. J. W., 1923 Abbildungen von Flügelgeäder der Coleopteren *Entomologische Mitteilungen*, 12, 71-115, 33 pls, 791 figs.
- Kerzhner, I.M. (1991) Histoire abrégée des insectes qui se trouvent aux environs de Paris (Geoffroy, 1762): proposed conservation of some generic names (Crustacea and Insecta) *Bulletin of Zoological Nomenclature*, 48 (2), 107-134.
- Kimmey, J.W. & Furniss, R.L. (1943) Deterioration of fire killed Douglas fir. *Technical Bulletin*, United States Department of Agriculture, 851, 1–6.
- Kirby, W. (1837) Insects. Coleoptera. In: Richardson J., *Fauna Boreali-Americana; or the Zoology of the Northern Parts of British America*. Josiah Fletcher, Norwich, 325 pp.
- Kirk, H.B. & Knull, J.N. (1926) Annotated list of the Cerambycidae of Pennsylvania. *The Canadian Entomologist*, 58(1), 21–26; (2), 39–46.
- Kirk, V.M. & Balsbaugh, E.U. (1975) A list of the beetles of South Dakota. *Technical Bulletin of the South Dakota Agricultural Experiment Station*, 42, 1–139.
- Klages, E.A. (1901) Supplement to Dr. John Hamilton's list of the Coleoptera of southwestern Pennsylvania *Annals of the Carnegie Museum*, 1, 265-294.
- Klingeman, W. E., Youseff, N. N., Oliver, J. B. & Basham J. P. (2017) The Longhorned Beetles (Coleoptera: Cerambycidae) of Tennessee: Distribution of Species, seasonal adult activity, and new State records *Florida Entomologist* 100 (2): 292-302, 1 carte.
- Knaus, W. (1885) Additions to the catalogue of Kansas Coleoptera for 1883 and 1884. *Transactions of the Kansas Academy of Science*, 9, 57-61.

- Knaus, W. (1903) The Coleoptera of the Sacramento Mountains of New Mexico. *Entomological News*, 14(6), 172–180.
- Knaus, W. (1904) The Coleoptera of the Sacramento Mountains of New Mexico. II. *Entomological News*, 15(4), 152–157.
- Knowlton, G.F. (1930) Notes on Utah Coleoptera. *Florida Entomologist*, 14(3), 53–56; (4), 75–77.
- Knowlton, G.F. (1934) Distributional notes on Utah Coleoptera. *Journal of the Kansas Entomological Society* 7 (3), 79–86.
- Knowlton, G.F. & Thatcher, T.O. (1936) Notes on wood-boring beetles. *Utah Academy of Sciences, Arts and Letters*, 13, 277–281.
- Knowlton, G.F. & Wood, S.L. (1950) An annotated list of Utah Cerambycidae. *Bulletin of the Brooklyn Entomological Society*, 45(1), 10–13.
- Knoll, J.N. (1946) The long-horned beetles of Ohio (Coleoptera, Cerambycidae). *Bulletin of the Ohio Biological Survey*, 39, 133–354.
- Korotyaev, B.A., Konstanstinov, A.S., Lingafelter, S. W., Mandelshtam, M.Y & Volkovitsh, M.G. (2005) Gall-inducing Coleoptera *In* Biology, Ecology, and Evolution of Gall-inducing Arthropods. Anantanarayanan Raman, Carl W. Schaefer & Toni M. Withers Editors. Science Publishers, Inc. Enfield (NH): 239-271, 19 figs, 2 tables.
- Kratz, G. (1863) Ueber einige zum Theil neue Cerambyciden-Gattungen. *Berliner Entomologische Zeitschrift*, 7, 97–108.
- Kratz, G. (1879) Ueber die Bockkäfer Ost-Sibiriens, namentlich die von Christoph am Amur gesammelten. *Deutsches Entomologische Zeitschrift*, 23, 77–177.
- Lackerbeck, K. (1998) Neue und wenig bekannte Prioninae (Coleoptera, Cerambycidae). *Entomofauna Zeitschrift für Entomologie*, 19(32), 517–524.
- Lacordaire, J.T. 1868. *Histoire naturelle des insectes. Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes. Tome huitième*. Librairie Encyclopédique de Roret, Paris. 552 pp. [Although dated “1869” this volume was published by November 28, 1868 (*Bibliographie de la France*)]
<https://doi.org/10.5962/bhl.title.67686>
- Lacordaire, J.T. (1869) *Histoire Naturelle des Insectes. Genera des Coléoptères, ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes*. Librairie Encyclopédique de Roret, Paris, 9(1), 409 pp.
<https://doi.org/10.5962/bhl.title.67686>
- Lacordaire, J.T. (1872) *Histoire Naturelle des Insectes. Genera des Coléoptères, ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes*. Librairie Encyclopédique de Roret, Paris, 9(2), 411–930.
<https://doi.org/10.5962/bhl.title.67686>
- Lacordaire, J.T. (1876) *Histoire Naturelle des Insectes. Genera des Coléoptères, ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes*. Atlas. Librairie Encyclopédique de Roret, Paris, 47 pp., 134 pls.
- Laliberté J. L., Chantal C. & Larochelle A. (1977) Ecologie des longicornes du Québec. *Faberies* 3, 88-102.
- Lameere, A.A. (1902a) Révision des prionides (Premier mémoire - Parandrides). *Annales de la Société Entomologique de Belgique*, 46, 59-111.
- Lameere, A.A. (1902b) Révision des prionides (Troisième mémoire - Spondylines). *Annales de la Société Entomologique de Belgique*, 46, 303–334.
- Lameere, A.A. (1902c) Révision des prionides (Quatrième mémoire - Sténodontines). *Mémoires de la Société Entomologique de Belgique*, 9, 63–110.
- Lameere, A.A. (1903a) Révision des prionides (Septième mémoire). *Mémoires de la Société Entomologique de Belgique*, 11, 1–216.
- Lameere, A.A. (1903b) Révision des prionides (Huitième mémoire - Mécosarthrines). *Annales de la Société Entomologique de Belgique*, 47, 307–320.
- Lameere, A.A. (1911) Révision des prionides (Dix-huitième mémoire - Prionines. V). *Annales de la Société Entomologique de Belgique*, 55, 260–277.

- Lameere, A.A. (1912a) Révision des prionides (Vingtième mémoire - Prionines VII). *Annales de la Société Entomologique de Belgique*, 56, 185–260.
- Lameere, A.A. (1912b) Révision des prionides (Vingt-déuxième mémoire - Addenda et Corrigenda). *Mémoires de la Société Entomologique de Belgique* 21, 113-188.
- Lameere, A.A. (1913) *Coleopterorum Catalogus*. Pars 52: Cerambycidae: Prioninae. W. Junk, Berlin, 108 pp.
- Lameere, A.A. (1915) Note sur les Prioninae du Muséum National d'Histoire Naturelle de Paris. *Bulletin du Museum National d' Histoire Naturelle*, 21, 51–63.
- Lameere, A.A. (1919) *Genera Insectorum*. Coleoptera, Fam. Cerambycidae, subfam. Prioninae. P. Wytsman, Bruxelles, 172, 189 pp.
- Lange, W.W.H. (1937) An annotated list of the insects, mostly Coleoptera, associated with Jeffrey pine in Lassen National Forest, California *The Pan-Pacific Entomologist*, 13 (4), 172-175.
- Laplante, S. (2017) Description of a new Nearctic species of *Tragosoma* Audinet-Serville (Coleoptera: Cerambycidae: Prioninae), with species validations, new synonymies and a lectotype designation. *Insecta Mundi*, 0578, 1-17.
- Laporte, F.L.N. (1840) *Histoire Naturelle des Insectes Coléoptères avec une introduction renfermant l'anatomie et la physiologie des animaux articulés, par M. Brullé; ouvrage accompagné de 155 planches gravées sur acier représentant plus de 800 sujets*. P. Duménil, Paris, 2, 563 pp.
- Latreille, P.A. (1802) *Histoire naturelle, générale et particulière, des crustacés et des insectes. Ouvrage faisant suite à l'histoire naturelle générale et particulière, composée par Leclerc de Buffon, et rédigée par C.S. Sonnini, membre de plusieurs sociétés savantes. Tome troisième. Familles naturelles des genres*. Dufart, Paris. 3: xii + pp. 13-467 (+ 1 p. errata, unnumbered).
- Laurent, P. (1905) Feldman Collecting Social. Meeting of December 21, 1904 *Entomological News*, 16 (2), 62.
- LeConte, J.L. (1847) Fragmenta Entomologica. *Journal of the Academy of Natural Sciences of Philadelphia* (n. s.) (2)1, 71-93.
- LeConte, J.L. (1850a) An attempt to classify the Longicorn Coleoptera of the part of America North of Mexico. *Journal of the Academy of Natural Sciences of Philadelphia* (ser. 2, 2, 5-38).
- LeConte, J.L. (1850b) General remarks upon the Coleoptera of Lake Superior *In Agassiz. Lake Superior, its physical character, vegetation, and animals*. Boston : 461 pp.
- LeConte, J.L. (1851) An attempt to classify the longicorn Coleoptera of the part of America, north of Mexico. *Journal of the Academy of Natural Sciences of Philadelphia*, (2)2, 99–112.
- LeConte, J.L. (1852) An attempt to classify the longicorn Coleoptera of the part of America, north of Mexico. *Journal of the Academy of Natural Sciences of Philadelphia*, (2)2, 139–178.
- LeConte, J.L. (1853) Descriptions of twenty new species of Coleoptera inhabiting the United States. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 6, 226–235.
- LeConte, J.L. (1854a) Descriptions of some new Coleoptera from Oregon, collected by Dr. J. G. Cooper on the North Pacific R. R. Expedition under Gov. J. J. Stevens. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 7, 16–20
- LeConte, J.L. (1854b) Notice of some coleopterous insects, from the collection of the Mexican Boundary Commission. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 7, 79–85.
- LeConte, J.L. (1854c) Some corrections in the nomenclature of Coleoptera found in the United States. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 7, 216–220.
- LeConte, J.L. (1857) Report upon the insects collected on the survey, pp. 1–72, 2 pls. *In: Report of explorations and surveys to ascertain the most practicable and economical route for a railroad from the Mississippi River to the Pacific Ocean*. Made under the direction of the Secretary of War in 1853-5, according to Acts of Congress of March 3, 1853, May 31, 1854 and August 5, 1854. Washington, House of Representatives, 12, Book 2, Part III.
- LeConte, J.L. (1858) John Lawrence, 1858 Catalogue of Coleoptera of the Regions adjacent to the Boundary Line between the United States and Mexico. *Journal of the Academy of Natural Sciences of Philadelphia* (ser. 2) 4, 9-42.

- LeConte, J.L. (1859a) The Coleoptera of Kansas and Eastern New Mexico. *Smithsonian Contributions to Knowledge*, 2(6), 1–58.
- LeConte, J.L. (1859b) *The complete writings of Thomas Say on the entomology of North America*. J. L. LeConte, New York, 1, 412 pp.; 2, 814 pp.
- LeConte, J.L. (1861a) Notes on the coleopterous fauna of Lower California. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 13, 335–338.
- LeConte, J.L. (1861b) New species of Coleoptera inhabiting the Pacific district of the United States. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 13, 338–359.
- LeConte, J.L. (1862) Notes on the classification of Cerambycidae with descriptions of new species. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 14, 38–43.
- LeConte, J.L. (1869) List of Coleoptera collected in Vancouver's Island by Henry and Joseph Matthews, with Descriptions of some new Species. *The Annals and Magazine of Natural History*, London (4) 4, 368–385.
- LeConte, J.L. (1873a) New species of North American Coleoptera. Prepared for the Smithsonian Institution. Part II. *Smithsonian Miscellaneous Collections*, 11(264), 169–238.
<https://doi.org/10.5962/bhl.title.17758>
- LeConte, J.L. (1873b) Classification of the Coleoptera of North America. Part II. *Smithsonian Miscellaneous Collections*, 11(265), 279–348.
- LeConte, J.L. (1878) Stridulation of Coleoptera *Psyche*. 2, 126.
- Leech, H.B. (1944) Spring flight of *Atimia dorsalis* (Coleoptera, Cerambycidae). *The Pan-Pacific Entomologist*, 20(1), 30.
- Leech, H.B. (1947) *Sarcophaga rapax* reared from *Prionus californicus*. *The Canadian Entomologist*, 79(7-8), 141.
- Leech, H.B. (1955) Orr's records of beetles eaten by the pallid bat. *The Coleopterists' Bulletin*, 9(4), 52.
- Leech, H.B. (1957) [Trip to Junipero Serra Peak, Santa Lucia Mountains, Monterey Co., California, August 8-16]. *The Pan-Pacific Entomologist*, 33(1), 50–51.
- Leech, H.B. (1968) Distribution of the cerambycid beetle *Ergates pauper* Linsley. *The Pan-Pacific Entomologist*, 44, 86.
- Leng, C.W. (1884) Synopses of the Coleoptera (Cerambycidae). *Bulletin of the Brooklyn Entomological Society*, 7(1), 7–11; (4), 57–64; (7), 95–101; (8), 112–116.
- Leng, C.W. (1885) Synopses of Cerambycidae. *Entomologia Americana*, 1(2): 28–35; (6), pl. 3; (7), 130–136.
- Leng, C.W. (1890) Synopses of Cerambycidae. *Entomologia Americana*, 6(1), 9–13; (4), 65–69; (5), 97–98; (6), 104–110; (8), 156–160; (10), 185–200; (11), 213–214.
- Leng, C.W. in Leng & Hamilton, 1896: 103, Synopsis of the Cerambycidae of North America. Part III, The Lamiinae. *Transactions of the American Entomological Society*, 23, 101–178.
- Leng, C.W. (1910) Notes on Coleoptera collected in northern Georgia. *Journal of the New York Entomological Society*, 18(2), 71–82.
- Leng, C.W. (1911) Notes on Coleoptera collected in northern Georgia - II *Journal of the New York Entomological Society* 19 (4): 209–216.
- Leng, C.W. & Hamilton, J. (1896) Synopsis of the Cerambycidae of North America. Part III, The Lamiinae. *Transactions of the American Entomological Society*, 23, 101–178.
- Leng, C.W. & Mutchler, A.J. (1914) A preliminary list of the Coleoptera of the West Indies as recorded to January 1, 1914. *Bulletin of the American Museum of Natural History*, 33, 391–493.
- Leonard, M.D. (1928) A list of the insects of New York with a list of the spiders and certain other allied groups. *Memoirs of the Cornell University Agricultural Experimental Station*, 101, 1–1121.
- Lepelletier de Saint Fargeau, A.L.M. & Audinet-Serville, J.G. (1825) In: *Encyclopédie méthodique. Histoire naturelle. Entomologie, ou histoire naturelle des Crustacés, des Arachnides et des insectes*. Paris, 10(1), 344 pp.
- Lewis, A.E. (1979) A list of Cerambycidae from the Hualpai Mountains, Mojave County, Arizona (Coleoptera). *The Pan-Pacific Entomologist*, 55(1), 21–26.

- Lingafelter, S.W. (1998) Observations of interactive behavior in *Parandra glabra* (Coleoptera: Cerambycidae). *Entomological News*, 109(1), 75–80.
- Lingafelter, S. W. (2007) Illustrated key to the Longhorned beetles of the Eastern United States. Coleopterists Society Special Publication, 3, 1-206
- Lingafelter, S.W. & Horner, N.V. (1993) The Cerambycidae of North Central Texas. *The Coleopterists' Bulletin*, 47(2), 159–191.
- Lingafelter, S.W. & Micheli, C.J. (2004) New species of Cerambycidae (Coleoptera) from Puerto Rico with records and notes for other species. *Journal of the New York Entomological Society*, 112(1), 37–55.
[https://doi.org/10.1664/0028-7199\(2004\)112%5B0037:nsoccf%5D2.0.co;2](https://doi.org/10.1664/0028-7199(2004)112%5B0037:nsoccf%5D2.0.co;2)
- Lingafelter, S.W., Nearns, E. H., Tavakilian. G.L., Monné, M.A. & Biondi, M. (2014) *Longhorned Woodboring Beetles (Coleoptera, Cerambycidae and Disteniidae) Primary types of the Smithsonian Institution*. Smithsonian Institution Scholarly Press, Washington. D. C., 390 pp.
- Linnaeus, C. (1758) *Systema naturae per regne tria naturae secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Salvius, Holmiae, 1, 823 pp.
<https://doi.org/10.5962/bhl.title.559>
- Linnaeus, C. (1761) *Fauna Suecica sistens animalia Sueciae regni: Mammalia, Aves, Amphibia, Pisces, Insecta, Vermes. Distributa per classes et ordines, genera et species, cum differentiis specierum, synonymis auctorum, nominibus incolarum, locis natalium descriptionibus insectorum*. Salvius, Holmiae, 578 pp.
<https://doi.org/10.5962/bhl.title.46380>
- Linnaeus, C. (1767) *Systema natura* editio duodecima reformata. Salvius, Holmiae; Thomas, Vindobonae, 1(2), 533–1327
- Linsley, E.G. (1934a) A short review of the genus *Atimia*, with the description of two new species (Coleoptera, Cerambycidae). *The Pan-Pacific Entomologist*, 10(1), 23–26.
- Linsley, E.G. (1934b) Studies in the Cerambycidae of Lower California (Coleoptera). *The Pan-Pacific Entomologist*, 10(2), 59–63.
- Linsley, E.G. (1934c) Notes and descriptions of West American Cerambycidae (Coleoptera). *Entomological News*, 45(6), 161–165; (7), 181–185.
- Linsley, E.G. (1935a) Studies in the Longicornia of Mexico (Coleoptera: Cerambycidae). *Transactions of the American Entomological Society*, 61, 67–102.
- Linsley, E.G. (1935b) Notes and descriptions of West American Cerambycidae II (Coleoptera). *Entomological News*, 46(6), 161–166.
- Linsley, E.G. (1938) Synonymical notes on some North American Cerambycidae (Coleoptera). *The Pan-Pacific Entomologist*, 14(3), 105–109.
- Linsley, E.G. (1939) The longicorn tribe Atimiini (Coleoptera: Cerambycidae). *Bulletin of the Southern California Academy of Sciences*, 38(2), 63–80.
- Linsley, E.G. (1940) Notes and descriptions of West American Cerambycidae IV (Coleoptera). *Entomological News*, 51(9), 253–258; (10), 285–290.
- Linsley, E.G. (1942) Contributions towards a knowledge of the insect fauna of Lower California No. 2 Coleoptera: Cerambycidae. *Proceedings of the California Academy of Sciences*, (4)24, 21–96.
- Linsley, E.G. (1957) Descriptive and synonymical notes on some North American Cerambycidae (Coleoptera). *American Museum Novitates*, 1828, 1–21.
- Linsley, E.G. (1958) The role of Cerambycidae in forest, urban and agricultural environments *The Pan-Pacific Entomologist*, 34 (3). 105-124.
- Linsley, E. G. (1959) A new subspecies of *Tragosoma* from southeastern Arizona (Coleoptera: Cerambycidae). *The Pan-Pacific Entomologist*, 35(3), 127–128.
- Linsley, E.G. (1961) Lycidlike Cerambycidae (Coleoptera). *Annals of the Entomological Society of America*, 54(5), 628–635.
- Linsley, E.G. (1962) The Cerambycidae of North America. Part II. Taxonomy and classification of the Parandrinae, Prioninae, Spondyliinae and Aseminae. *University of California, Publications in Entomology*, 19, 1–102.

- Linsley, E.G. & Chemsak, J.A. (1966c) Cerambycidae of the Revilla Gigedo Islands (Coleoptera). *Proceedings of the California Academy of Sciences*, (4) 33(10), 249–254.
- Linsley, E.G. & Chemsak, J.A. (1997) The Cerambycidae of North America. Part VIII. Bibliography, Index and Host Plant Index. *University of California, Publications in Entomology*, 117, 534 pp.
- Linsley, E.G., Knull, J.N. & Statham, M. (1961) A list of Cerambycidae from the Chiricahua Mountain area, Cochise County, Arizona. *American Museum Novitates*, 2050, 1–34.
- Linsley, E.G. & Ross, E.S. (1940) Records of some Coleoptera from the San Jacinto Mountains, California. *The Pan-Pacific Entomologist*, 16(2), 75–76.
- Lockhead, W. (1903) A key to orchard insects *33rd Annual Report of the Entomological Society of the Province of Ontario* 1902: 101-114, figs 64-103.
- Loding, H.P. (1933) Alabama Coleoptera not generally listed from the Gulf Coast States of the Mississippi River., Fla., Ga., Ala., and Miss. *Bulletin of the Brooklyn Entomological Society*, 28(4), 139–151.
- Loding, H.P. (1945) Catalogue of the beetles of Alabama. *Monograph of the Geological Survey of Alabama*, 11, 1–172.
- Lozada Piña, A, Fernández García & Trujillo Anaya, M. (2004) Lista preliminar de los coleópteros (Insecta, Coleoptera) de Topes de Collarte, Trinidad, Sancti Spiritus, Cuba. *Boletín de la Sociedad Entomológica Aragonesa*, 34, 101-106.
- Lugger, O. (1899) Beetles injurious to fruit-producing plants *Bulletin of the Minnesota Agricultural Experiment Station* 66, 85-331.
- MacGillivray, A.D. & Houghton, C.O. (1902) A List of insects taken in the Adirondack Mountains, New York. - I. *Entomological News*, 13 (8), 247-253.
- MacKay, W.P., Zac, J.C. & Hovore, F.T. (1987) Cerambycid beetles (Coleoptera: Cerambycidae) of the northern Chihuahuan Desert (South Central New Mexico). *The Coleopterists' Bulletin*, 41(4), 361–369.
- MacRae, T.C. (1993) Annotated checklist of the longhorned beetles (Coleoptera: Cerambycidae and Disteniidae) occurring in Missouri. *Insecta Mundi*, 7, 223–252.
- Maes, J.-M. (1998) Catálogo de los insectos y artrópodos terrestres de Nicaragua. Setab Bosawas, Marena, Managua, Nicaragua, 1899 pp.
- Maes, J.-M., Allen, A., Monné, M.A. & Hovore, F.T. (1994) Catálogo de los Cerambycidae (Coleoptera) de Nicaragua. *Revista Nicaraguense de Entomología*, 27, 1–58.
- Maes, J.-M., Berghe, E., Dauber, D., Audureau, A., Nearn, E., Skilman, F., Heffern, D. & Monné, M.A. (2010a). Catalogo ilustrado de los Cerambycidae (Coleoptera) de Nicaragua. Parte I - Parandrinae, Prioninae, Aseminae. *Revista Nicaraguense de Entomología*, 70 (Suplemento 1-1), 1-102.
- Maier, C.T. (2017) Cerambycidae (Coleoptera) accidentally introduced into Connecticut from China or from Other areas in the United States *Proceedings of the Entomological Society of Washington* 119 (3), 423-429
- Majka C.G. & Ogden J. (2010) New records of Cerambycidae in Nova Scotia. *Acadian Entomological Society* 6, 12-15.
- Maki, E.C., Millar, J.G., Rodstein, J., Hanks, L.M. & Barbour, J.D. (2011) Evaluation of mass trapping and mating disruption for managing *Prionus californicus* (Coleoptera, Cerambycidae) in hop production yards. *Journal of Economic Entomology*, 104(3), 933-938.
- Mank, E. (1934) The Coleoptera of Glacier Park, Montana. *The Canadian Entomologist*, 66(4), 73–81.
- Mankin, R.W., Burman, H., Menocal, O. & Carrillo, D. (2018). Acoustic detection of *Mallodon dasystemus* (Coleoptera: Cerambycidae) in *Persea americana* (Laurales: Lauraceae) branch stumps. *Florida Entomologist*, 101 (2), 321-323.
- Manley, G.V. & French, J.V. (1976) Wood boring beetles inhabiting citrus in the Lower Rio Grande Valley of Texas. Part I. Cerambycidae. *Journal of the Rio Grande Valley Horticultural Society*, 30, 45–53.
- Mannerheim, C.G. (1843) Beitrag zur Käfer-Fauna der Aleutischen Inseln, der Insel Sitkha und Neu-Californiens. *Bulletin de la Société Impériale des Naturalistes de Moscou* 16 (1), 175-314.

- Mannerheim, C.G. (1852) Zweiter Nachtrag zur Käferfauna der Nord-Amerikanischen Laender der Russischen Reiches. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 25(2), 283–387.
- Mannerheim, C. G. (1853) Dritter Nachtrag zur Käfer-Fauna der Nord-Amerikanischen Laender des Russischen Reiches. *Bulletin de la Société Impériale des Naturalistes de Moscou* 26 (3-4), 95-273.
- Marten, J. (1890) The tile-horned *Prionus Prairie Farmer*, January 1890, 60.
- Martínez, C. (2000) Escarabajos longicórnios (Coleoptera, Cerambycidae) de Colombia. *Biota Colombiana*, 1(1), 76-105.
- Mason, F.R. (1910) Collecting beetles in Mexico. *The Canadian Entomologist*, 42(1), 22–24
- Meeking J.M., Seabrook. W.D. & Paim, U. (1974) Perception of carbon dioxide by larvae of *Orthosoma brunneum* (Coleoptera: Cerambycidae) as indicated by recordings from the ventral nerve cord *The Canadian Entomologist*, 106 (3), 257-262, 3 figs.
- Melander, A.L. (1904) Destructive beetles, a note on landscape gardening. *Entomological News*, 15 (1), 19-20, pl. 5.
- Melzer, J. (1919) Os longicórneos brasileiros da subfamília "Prioninae" tomando em consideração particular as espécies do Estado de São Paulo. *Revista do Museu Paulista*, 11, 1–207.
- Moennich, H. C. (1934) H. C., 1934 Trapping for Coleoptera *Bulletin of the Brooklyn Entomological Society* 29 (3): 97-98.
- Monné, M.A. (1994) *Catalogue of the Cerambycidae (Coleoptera) of the Western Hemisphere. Part XII. Subfamilies: Parandrinae, Anoplodermatinae, Spondylinae, Aseminae and Oxypeltinae*. Sociedade Brasileira de Entomologia, São Paulo, 56 pp.
- Monné, M.A. (1995b) *Catalogue of the Cerambycidae (Coleoptera) of the Western Hemisphere. Part XXII. Subfamily Prioninae*. Sociedade Brasileira de Entomologia, São Paulo, 115 pp.
- Monné, M.A. (2002) Catalogue of the Neotropical Cerambycidae (Coleoptera) with known host plant - Part V: Subfamilies Prioninae, Parandrinae, Oxypeltinae, Anoplodermatinae, Aseminae and Lepturinae. *Publicações Avulsas do Museu Nacional*, 96, 1–70.
- Monné, M.A. (2006) Catalogue of the Cerambycidae (Coleoptera) of the Neotropical Region. Part III. Subfamilies Parandrinae, Prioninae, Anoplodermatinae, Aseminae, Spondylidinae, Lepturinae, Oxypeltinae, and addenda to the Cerambycinae and Lamiinae. *Zootaxa*, 1212, 1-244.
- Monné, M. A. (2012) Catalogue of the type-species of the genera of the Cerambycidae, Disteniidae, Oxypeltidae and Vesperidae (Coleoptera) of the Neotropical Region. *Zootaxa*, 3213, 1-183.
- Monné, M.A. & Giesbert, E.F. (1994) *Checklist of the Cerambycidae and Disteniidae (Coleoptera) of the Western Hemisphere*. Wolfsgarden Books, Burbank, 409 pp.
- Monné, M.A. & Giesbert, E.F. (1995) *Checklist of the Cerambycidae and Disteniidae (Coleoptera) of the Western Hemisphere*. Wolfsgarden Books, Burbank, 419 pp.
- Monné, M.A. & Hovore, F.T. (2006) *Checklist of the Cerambycidae, or longhorned wood-boring beetles of the Western Hemisphere*. BioQuip Publications, 393 pp
- Monné, M.L, Monné, M.A. & Wang, Q. (2017) *Chapter 1. General Morphology, Classification, and Biology of Cerambycidae*. In: Wang, Q. *Cerambycidae of the World: Biology and Pest Management*. CRC Press, Taylor & Francis, Boca Raton, 1-642 p.
- Monné, M.L. & Santos-Silva, A. (2003) Sinopse do gênero *Strongylaspis* Thomson, 1860 (Coleoptera, Cerambycidae, Prioninae, Macrotomini). *Revista Brasileira de Entomologia*, 47(1), 31–47.
- Moore, I. (1937) A list of the beetles of San Diego County, California. *Occasional Papers of the San Diego Society of Natural History*, 2, 1–109.
- Morris, F. J. A. (1908) «Some beetle-haunts» by an amateur botanist *The Canadian Entomologist*, 40 (12), 441-449.
- Morris, F.J.A. (1909) «Some beetle-haunts» by an amateur botanist *39th Annual Report of the Entomological Society of the Province of Ontario*, [1908], 56-63, figs 11-15.
- Morris, F J.A. (1916) Reports on insects of the year. Division No. 5, Port Hope District *46th Annual Report of the Entomological Society of the Province of Ontario* 1915, 17-21.

- Morris, F. J. A. (1918) Popular and practical Entomology. The Heart of a Wood-pile *The Canadian Entomologist*, 50 (2), 37-43
- Morris, R. F. (1987) New records of Georgia Cerambycidae (Coleoptera: Cerambycidae) *Journal of Entomological Science* 22 (2), 140-141.
- Morris, R.F. (2002) Distribution and biological notes for some Cerambycidae (Coleoptera) occurring in the Southeastern United States. *Insecta Mundi* 16(4), 209-213.
- Motschulsky, V. (1845) Remarques sur la collection de coléoptères russes de Victor de Motschulsky. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 18(1), 1-127.
- Moura, L.A. & Von Groll, E. (2018) Catalogue of Coleoptera type specimens housed in the collection of the Museu de Ciências Naturais, Fundação Zoobotânica do Rio Grande do Sul, Brazil *Zootaxa*, 4318 (3), 439-473.
- Mroczkowski, M. (1986) *Tetropium* Kirby, 1837 (Insecta, Coleoptera, Cerambycidae): proposed conservation by the suppression of *Isarthron* Dejean, 1835. Z.N.(S.)2534. *Bulletin of Zoological Nomenclature*, 43 (2), 188-190.
- Mullins, A.J. (1975) A note on the fecundity of *Prionus imbricornis* (Linn) (Coleoptera: Cerambycidae). *The Coleopterists' Bulletin* 29 (1), 43.
- Mulsant, E. (1839) *Histoire naturelle des Coléoptères de France*. Longicornes. Maison, Paris & Lyon, 304 pp.
- Mulsant, E. (1863) *Histoire naturelle des Coléoptères de France*. Longicornes. Maison, Paris, 540 pp.
- Mulsant, E. (1864) Tribu des Longicornes (suite). *Annales des Sciences Physiques et Naturelles, d'Agriculture et d'Industrie*, 8(3), 1-208.
- Murray, A. (1862) On the geographical relations of the Coleoptera of Old Calabar. *The Transactions of the Linnean Society of London* 23, 449-455.
- Mutchler, A.J. (1912) In Proceedings of the New York Entomological Society. Meeting of March 5, 1912. "Notes and Exhibition of Some Species of Cerambycidae in the Collection of the American Museum of Natural History. *Journal of the New York Entomological Society* 20 (3), 213-215.
- Mutchler, A.J. (1914) A new species of *Criocephalus*. *Bulletin of the American Museum of Natural History*, 33, 345-346.
- Napp, D.S. (1994) Phylogenetic relationships among the subfamilies of Cerambycidae (Coleoptera, Chrysomeloidea). *Revista Brasileira de Entomologia*, 38(2), 265-419.
- Nicolay, A.S. (1917) Buprestidae and Cerambycidae from Maine *Bulletin of the Brooklyn Entomological Society*, 12 (4), 92-95.
- Nicolay, A.S. (1919) A list of the Buprestidae and Cerambycidae taken on Long Island, N. Y. *Bulletin of the Brooklyn Entomological Society*, 14(1), 17-20; (2), 63-72.
- Nishio, Y. (1956a) Morphologie und geographische Verbreitung von *Prionus californicus*, *laticollis*, *coriarius* und *insularis*. *Kontyu*, 24, 240-246.
- Nishio, Y. (1956b) Die Gattung *Ergates* Serville, 1832, insbesondere Phylogenie und Verbreitung der Arten. *Seibutsu Kagaku*, 8(2), 68-73.
- Noguera, F.A. & Chemsak, J.A. (1996) Cerambycidae (Coleoptera), pp. 381-409. In: *Biodiversidad taxonomía, y biogeografía de artrópodos de México: Hacia una síntesis de su conocimiento*. Universidad Nacional Autónoma de México, México, D. F.
- Noguera, F.A., Zaragoza-Caballero, S., Chemsak, J.A., Rodriguez-Palafox, A., Ramirez, E., González-Soriano, E. & Ayala, A. (2002) Diversity of the family Cerambycidae (Coleoptera) of the Tropical Dry Forest of Mexico, I. Sierra de Huautla, Morelos. *Annals of the Entomological Society of America*, 95(5), 617-627.
- Noguera, F.A., Chemsak, J.A., Zaragoza-Caballero, S., Rodriguez-Palafox, A., Ramirez-García, E., González-Soriano, E & Ayala, R. (2007) A faunal study of Cerambycidae (Coleoptera) from one region with tropical dry forest in México: San Buenaventura, Jalisco. *The Pan-Pacific Entomologist* 83(4), 296-314.
- Noguera, F.A., Ortega-Huerta, M.A., Zaragoza-Caballero, S., González-Soriano, E & Ramirez-García, E. (2009) A faunal study of Cerambycidae (Coleoptera) from one region with tropical dry forest in Mexico: Sierra de San Javier, Sonora. *The Pan-Pacific Entomologist* 85(2), 70-90.

- Noguera, F.A., Ortega-Huerta, M.A., Zaragoza-Caballero, S., González-Soriano, E., & Ramírez-García, E. (2017) Species richness and abundance of Cerambycidae (Coleoptera) in Huatulco, Oaxaca, Mexico; relationships with phenological changes in the Tropical Dry Forest. *Neotropical Entomology*, (Anexo 13 p.).
- Nussler, H. (1961) Ein beständiges Vorkommen des nearktischen Bockkäfers *Parandra brunnea* Fabr. (Col. Cerambycidae) in Deutschland. *Abhandlungen und Berichte aus dem staatlichen Museum für Tierkunde in Dresden* 26 (16), 125-130.
- Olivier, A.G. (1790) *Encyclopédie méthodique, ou par ordre de matières; par une société de gens de lettres, de savans et d'artistes; précédée d'un vocabulaire universel, servant de table pour tout l'ouvrage, ornée des portraits de Mm. Diderot & d'Alembert, premiers éditeurs de l'Encyclopédie*. Plomteux, Paris; Panckoucke Libr., Liège, 5(1), 368 pp.
- Olivier, A.G. (1792) *Encyclopédie méthodique, ou par ordre de matières; par une société de gens de lettres, de savans et d'artistes; précédée d'un vocabulaire universel, servant de table pour tout l'ouvrage, ornée des portraits de Mm. Diderot & d'Alembert, premiers éditeurs de l'Encyclopédie*. Plomteux, Paris; Panckoucke Libr., Liège, 6(2), 369-704; 7(1), 368 pp.
- Olivier, A.G. (1795) *Entomologie, ou histoire naturelle des insectes, avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur figure enluminée*. Coléoptères. Desray, Paris, 4, 519 pp.
- O'Neill K.M., Fultz J.E. & Ivie M.A. (2008) Distribution of adult Cerambycidae and Buprestidae (Coleoptera) in a Subalpine Forest under shelterwood management *The Coleopterists' Bulletin* 62 (1), 27-36.
- Ortenburger, A.I. & Hatch, M.H. (1926) Notes on Coleoptera from southeastern Oklahoma with a few records from adjacent portions of Texas and Arkansas including a new species. *Transactions of the Oklahoma Academy of Sciences* 6. 142-148.
- Osten-Sacken, C.R. (1862) Descriptions of some larvae of North American Coleoptera *Proceedings of the Entomological Society of Philadelphia* 1 (5): 105-130, 1 pl.
- Ouellet, C.J. (1902) Liste des Coléoptères les plus remarquables capturés dans la province de Québec en 1899, 1900 et 1901. *Le Naturaliste Canadien*, 39 (8), 120-124.
- Pack, H.J. (1930) Notes on Utah Coleoptera. *Entomological News*, 41(7), 219-222.
- Packard, A.S. (1877) Alpheus Spring, 1877 Insects injuring shade and forest trees In 9th Annual Report of the U. S. Geological and Geographical survey of the territories embracing Colorado and parts of adjacent territories: being a report of progress of the exploration for the year 1875 by F.V. Hayden : 802-815.
- Packard, A.S. (1881) Insects injurious to forest and shade trees. *Bulletin of the United States Entomological Commission*, 7, 1-275.
- Packard, A.S. (1888) List of the spiders, myriopods and insects of Labrador. *The Canadian Entomologist*, 20(8), 141-149.
- Packard, A.S. (1889) Notes on the epipharynx, and the epipharyngeal organs of taste in mandibulate insects. *Psyche*, 5(157-159), 222-228.
- Packard, A.S. (1890) Insects injurious to forest and shade trees. *Bulletin of the United States Entomological Commission*, 7, 1-955. (enlarged edition).
- Paim, U. & Beckel W.E. (1963) The influence of oxygen and carbone dioxide on the spiracles of a wood-boring insect, *Orthosoma brunneum* (Forster) (Coleoptera: Cerambycidae) *Canadian Journal of Zoology* 41, 1149-1167, 10 figs.
- Paim, U. & Beckel, W. E.(1964) Effects of environmental gases on the mortality and survival of larvae and pupae of *Orthosoma brunneum* (Forster) (Coleoptera: Cerambycidae) *Canadian Journal of Zoology* 42, 59-69.
- Paim, U. & Beckel, W. E. (1964) The carbon dioxide related behavior of the adults of *Orthosoma brunneum* (Forster) (Coleoptera, Cerambycidae) *Canadian Journal of Zoology* 42, 295-304.
- Paim, U. & Beckel, W.E. (1964) The behavior of the larvae of *Orthosoma brunneum* (Forster) (Coleoptera, Cerambycidae), in relation to gases found in the logs inhabited by the larvae *Canadian Journal of Zoology* 42, 327-353.
- Palisot de Beauvois, A.M.J.F. (1805-1821) *Insectes recueillis en Afrique et Amérique dans les Royaumes d'Oware et de Benin, à Saint-Domingue et dans les États-Unis, pendant les années 1786-1797*. Imprimerie de Fain et Cie., Paris, 276 pp.

- Parmelee, F.T. (1941) Longhorned and flat-headed borers attacking fire-killed coniferous timbers in Michigan *Journal of Economic Entomology* 34, 377-380.
- Pascoe, F.P. (1860) Notices of new or little-known genera and species of Coleoptera. *The Journal of Entomology*, 1, 36-64; 98-132.
<https://doi.org/10.5962/bhl.title.24970>
- Pascoe, F.P. (1864-1869) Longicornia Malayana: or a descriptive catalogue of the species of the three longicorn families Lamiidae, Cerambycidae and Prionidae collected by Mr. A. R. Wallace in the Malay Archipelago. *The Transactions of the Entomological Society of London*, (3)3, 1-689.
- Payne, J.A., Lowman, H & Pate, R.R. (1975) Artificial diets for rearing the tilehorned *Prionus* *Annals of the Entomological Society of America*, 68 (4), 680-682, 2 figs.
- Payne J.A., Polles S.G., Sparks D. & Wehunt E. J. (1976) The distribution, economic importance, and chemical control of the Tilehorned *Prionus* (Coleoptera: Cerambycidae) in Georgia *Journal of the Georgia Entomological Society* 11 (1), 9-16, 2 figs.
- Peck, S.B. (2005) A checklist of the beetles of Cuba with data on distributions and bionomics. (Insecta, Coleoptera). *Arthropods of Florida and neighboring land areas*. Vol. 18: 1-214 Florida Department of Agriculture and Consumer Services, Gainesville.
- Peck, S.B. (2016) The beetles of the Lesser Antilles (Insecta, Coleoptera): diversity and distribution. *Insecta Mundi*, 460, 1-360.
- Peck, S.B., Cook, J., & Hardy, J.D. (2002) Beetle fauna of the island of Tobago, Trinidad and Tobago, West Indies. *Insecta Mundi*, 16(1-3), 9-23.
- Peck S. B. & Thomas M. C. (1998) A distributional checklist of the beetles (Coleoptera) of Florida. *Arthropods of Florida and Neighboring Land Areas*. 16: 1-180
- Penrose, R L. & Westcott, R.L. (1974) Notes on the distribution, hosts and bionomics of some Pacific Northwest Cerambycidae (Coleoptera) *The Coleopterists' Bulletin* 28 (4). 233-236.
- Perry, R.H. (1974) Notes on the longhorned beetles of Virginia. Part II (Coleoptera, Cerambycidae). *The Coleopterists' Bulletin*, 28(4), 215-217.
- Perry, R H. (1975) Notes on the long-horned beetles of Virginia, Part III (Coleoptera: Cerambycidae). *The Coleopterists' Bulletin* 29 (1), 59.
- Perry, R.H. (1977) Notes on the longhorned beetles of Virginia. Part IV (Coleoptera, Cerambycidae). *The Coleopterists' Bulletin*, 31(1), 97-99.
- Pettit, J. (1871) List of Coleoptera taken at Grimsby, Ontario. Cerambycidae. *The Canadian Entomologist*, 3 (6), 105-107.
- Pettit, R.H. (1904) Insects injurious to fruits in Michigan *Special Publications Michigan Agricultural Experiment Station* 24, 1-79, 70 figs
- Pic, M. (1916) Notes diverses, descriptions et diagnoses (Suite.) *L'Échange, Revue Linnéenne* 32 (375), 9-11.
- Pic, M. (1945) Nouvelles variétés de Coléoptères Longicornes *L'Échange, Revue Linnéenne* 61 (500), 5-7.
- Picard, F. (1929) Coléoptères. Cerambycidae *Faune de France* 20. 1-166, 71 figs. Paris, Paul Lechevalier.
- Planet, L.M. (1924) Histoire Naturelle des Longicornes de France. *Encyclopédie Entomologique*, (A) 2, 1-386.
- Plavilstshikov, N.N. (1915) Zhuki-usachi, sobranie A.I. Aleksandrovym v Manchzhurii (Col., Cerambycidae) *Entomologicheskyy Vestnik*, 2 (2), 103-110.
- Plavilstshikov, N.N. (1936) Fauna de l'URSS, Insectes Coléoptères. Cerambycidae I. *Fauna SSSR*, 21, 611 pp.
- Popenoe, E.A. (1877) A list of Kansas Coleoptera. *Transactions of the Kansas Academy of Sciences*, 5, 21-40
- Portevin, G. (1927) Tableaux dichotomiques pour la détermination des Longicornes de France. *Encyclopédie Entomologique*, (A) 2(Suppl.), 1-53.
- Procter, W. (1927) Biological survey of the Mount Desert Region. Part I. The insect fauna with reference to the flora and other biological features. *Philadelphia Wistar Institute of Anatomy and Biology* 1, 1-247.

- Procter, W. (1946) Biological survey of the Mountain Desert Region incorporated. Part VII , being a revision of parts I and IV with the addition of 1100 species. The insect fauna with references to method of capture, food plants, the flora and other biological features *Philadelphia Wistar Institute of Anatomy and Biology* 566 pp.
- Provancher, L.A. (1877) *Petite faune entomologique du Canada, précédés d'un traite élémentaire d'entomologie*. Les Coleoptères. C. Darveau, Québec, 1, 786 pp.
<https://doi.org/10.5962/bhl.title.8169>
- Provancher, L.A. (1877) Additions et corrections à la faune coléoptérologique de la Province de Québec. *Naturaliste Canadien*, 9(11), 321–338.
- Quayle, H.J. (1938) *Insects of citrus and other subtropical fruits*. Comstock Publ. Co., Ithaca, 583 pp.
- Quentin R.M & Villiers, A. (1974) Révision des Closterini de Madagascar (Col. Cerambycidae, Prioninae) *Annales de la Société Entomologique de France*, (N. S.) 10 (2,249-341, 149 figs.
- Quentin, R.M. & Villiers, A. (1975) Insectes Coléoptères Cerambycidae Parandrinae et Prioninae Faune de Madagascar 40: 1-251, 262 figs, Orstom, Paris.
- Rafinesque, C.R.(1815) *Analyse de la nature ou tableau de l'univers et des corps organisés* C. S. Rafinesque, Palermo, 1-224.
- Randall, J.W. (1838) Description of new species of coleopterous insects inhabiting the State of Maine. *Boston Journal of Natural History*, 2(1), 1–33
- Raske, A. G. (1973) *Tetropium parvulum* elevated to species rank and contrasted to *T. cinnamopterum* in morphology and host preference (Coleoptera, Cerambycidae) *The Canadian Entomologist*, 105, 745-755, 14 figs.
- Redtenbacher, L. (1845) *Die Gattungen der Deutschen Käfer-Fauna nach der analytischen Methode bearbeitet einem kurz gefassten Leitfaden, zum Studium dieses Zweiges der Entomologie*. Wien, Carl Gerold 177 pp, 2 pls.
- Redtenbacher, L. (1868) *Coleopteren In: Reise des österreichischen Fregatte Novara und die Erde in den Jahren 1857, 1858, 1859, unter der befehlen des Commodore B. von Wüllerstorff-Urbair*. Karl Gerold's Sohn, Wien, Zoologischer Theil, Zweiter Band, 249 pp.
<https://doi.org/10.5962/bhl.title.1597>
- Redtenbacher, L. (1874) *Fauna Austriaca*. Die Käfer, nach der analytischen Methode bearbeitet. Gerald, Wien, 2, 1–571.
- Reitter, E (1895) Beschreibung neuer oder wenig gekannter Coleopteren aus der Umgebung von Akbes in Syrien. *Wiener Entomologische Zeitung* 14 (3): 79-88, 3 figs.
- Reitter, E. (1912) *Fauna Germanica. Die Käfer des Deutschen Reiches*. Stuttgart, K. G. Lutz' Verlag, 4, 1–236.
- Rice, M.E. (1981) Notes on Cerambycidae from Missouri. *The Coleopterists Bulletin*, 35(4), 459–462.
- Rice, M. E., MacRae, T. & Merickel, F. (2017) The longhorned beetles (Coleoptera, Cerambycidae) of Idaho. *The Coleopterists Bulletin*, 71 (4), 667-678.
- Rice M. E., Turnbow R. H. Jr. & Hovore- F. T. (1985) Biological and distributional observations on Cerambycidae from the southwestern United States (Coleoptera). *The Coleopterists' Bulletin* 39 (1), 18-24.
- Ricksecker, L.E. (1885) [Lignivorous beetles]. *Entomologia Americana*, 1(5), 96–98
- Riley, C.V. (1870) Second annual report on the noxious, beneficial and other insects of the State of Missouri. *5th Annual Report of the Missouri State Board of Agriculture* 1869, 1-136, 99 figs.
- Riley, C.V. (1872) Miscellaneous Notes. Stridulation of *Orthosoma cylindricum*, Fabr. *The Canadian Entomologist*, 4 (7), 139-140.
- Riley, C.V. (1873) Fifth annual report on the noxious, beneficial and other insects of the State of Missouri *8th Annual Report of the Missouri State Board of Agriculture*, Jefferson City 1872 , 1-160, 75 figs.
- Riley, C.V. (1880) Food habits of the longicorn beetles or woodborers. *The American Entomologist*, 3(10), 237–239; (11), 270–271.
- Riley, C.V. (1884) Dwarfing of oaks by *Mallodon melanopus*. *Annual Report of the United States Commissioner of Agriculture for 1884*, 285-418, 10 pls.

- Riley C.V. & Howard L.O. (1891) The grape root *Prionus*. USDA Insect Life, 3 (9-10), 407.
- Rivers, J.J. (1886a) Contributions to the larval history of Pacific Coast Coleoptera. *Bulletin of the California Academy of Sciences*, 2, 64–72.
- Rivers, J.J. (1886b) *Contributions to the larval history of Pacific Coast Coleoptera*. The State Printer, Sacramento, 10 pp.
- Roff J. W. (1967) A record of *Tetropium cinnamopterum* Kirby in white spruce logs in central British Columbia Canadian Forest Service Bi-monthly Research Notes 23: 27.
- Rohwer, S.A. (1921) The North American Ichneumon-flies of the tribes Labenini, Rhyssini, Xoridini, Odontomerini, and Phytodietini. *Proceedings of the United States National Museum*, 57 (2317), 405-474, 13 figs.
- Rosales, C.J. (1966) Contribución al conocimiento de los Prioninae (Coleoptera, Cerambycidae) de Venezuela. *Revista de la Facultad de Agronomía*, 11, 1–237.
- Ross, D.A. (1968) Wood and bark-feeding Coleoptera of felled spruce in interior British Columbia. *Journal of the Entomological Society of British Columbia*, 65, 10–12.
- Ross D.A. (1967a) Wood and bark-feeding Coleoptera of felled western larch in British Columbia. *Journal of the Entomological Society of British Columbia* 64, 23-24.
- Ross D.A. (1967b) The western larch borer, *Tetropium velutinum* LeConte, in the interior British Columbia. *Journal of the Entomological Society of British Columbia* 64, 25-28
- Ross, D.A. (1968) Wood and bark-feeding Coleoptera of felled spruce in interior British Columbia *Journal of the Entomological Society of British Columbia* 65, 10-12.
- Ross D.A. & Geistlinger N.J. (1968) Protecting larch logs from *Tetropium velutinum* LeConte with Lindane emulsion. *Journal of the Entomological Society of British Columbia* 65, 14-15.
- Ross D.A. & Vanderwal, H. (1969) A spruce borer *Tetropium cinnamopterum* Kirby, in interior British Columbia. *Journal of the Entomological Society of British Columbia* 66, 10-14
- Ruette, R. (1970) A catalogue of types of Coleoptera in the Canadian National collection of insects. *Memoirs of the Entomological Society of Canada*, 72, 1–134.
- Saalas, U. (1936) Ueber das Flügelgeäder und die phylogenetische Entwicklung der Cerambyciden. *Annales Societatis Zoologicae Botanicae Fennicae Vanamo*, 4(1), 1–198.
- Sama, G. (2003) *Atlas of the Cerambycidae of Europe and the Mediterranean Area. Volume 1: Northern, Western, Central and Eastern Europe. British Isles and Continental Europe from France (excl. Corsica) to Scandinavia and Urals*. Vít Kabourek, Zlín, 2003 [2002]: 1-173, 729 figs.
- Sama, G. (2005) Description of *Neospondylis* gen. nov. from North America & Mexico (Coleoptera, Cerambycidae, Spondylidinae). *Les Cahiers Magellanes*, 43, 1–10.
- Santos-Silva, A. (2002) Notas e descrições em Parandrinae (Coleoptera, Cerambycidae, Parandrinae). *Iheringia*, Serie Zoologia, 92(2), 29–52.
- Santos-Silva, A. (2007) Revisão do gênero *Derobrachus* Audinet-Serville, 1832 (Coleoptera, Cerambycidae, Prioninae) *Arquivos de Zoologia*, 38 (1), 1-94, 362 figs.
- Santos-Silva, A. & Durán, J.R.E. (2009) Description of the female of *Strongylaspis granigera* Bates, 1884 (Coleoptera, Cerambycidae, Prioninae). *Spanish Journal of Agricultural Research*, 7(2), 349-354.
- Santos-Silva, A. & Galileo, M.H.M. (2011) Sobre alguns Cerambycidae (Coleoptera) das coleções do Instituto Nacional de Pesquisas da Amazônia (INPA) e da Universidade Federal do Amazonas (UFAM): Parandrinae, Prioninae e Disteniinae. *Les Cahiers Magellanes n.s.* 3: 20-51.
- Santos-Silva, A., Heffern, D. & Matsuda, K. (2010) Revision of Hawaiian, Australasian, Oriental, and Japanese Parandrinae (Coleoptera, Cerambycidae). *Insecta Mundi*, 0130, 1-120.
- Santos-Silva, A. & Martins, U.R. (2006) Novos táxons e notas taxonômicas em Prioninae (Coleoptera, Cerambycidae). *Iheringia, Série Zoologia*, 95(4), 393-403.
- Santos-Silva, A. & Martins, U.R. (2010) Subfamília Parandrinae, pp. 5-79. In: Martins, U. R. (org.) *Cerambycidae Sul-americanos (Coleoptera) Taxonomia*. Sociedade Brasileira de Entomologia, São Paulo, 11, 274 pp.
- Santos-Silva, A., Nearn, E.H. & Swift, I.P. (2016) Revision of the American species of the genus *Prionus* Geoffroy, 1762 (Coleoptera, Cerambycidae, Prioninae, Prionini). *Zootaxa*, 4134(1), 1-103.

- Santos-Silva, A., Swift, I.P. & Nearn, E.H. (2010) Division of the genus *Nothopleurus* Lacordaire, 1869 (Coleoptera, Cerambycidae, Prioninae). *Zootaxa*, 2643, 1-44.
- Santos-Silva, A. & Wappes, J.E. (2013) The genus *Aplagiognathus* Thomson, 1861 (Coleoptera, Cerambycidae, Prioninae, Macrotomini). *Insecta Mundi*, 264, 1-13.
- Saunders, H.S. (1887) Collecting at the electric light, 1886 *The Canadian Entomologist*, 19 (2), 21-29.
- Saunders, W. (1875) The cylindrical Orthosoma - *Orthosoma cylindricum*, Fabr. *The Canadian Entomologist*, 7 (2), 29, 2 figs
- Saunders, W. (1884) Insects injurious to the white pine - *Pinus strobus* 14th Annual Report of the Entomological Society of the Province of Ontario 1883, 52-59, figs 19-29.
- Savely, H. E. (1939) Ecological relations of certain animals in dead pine and oak logs *Ecological Monographs*, Durham 9, 321-385, 8 figs.
- Say, T. (1824) Descriptions of coleopterous insects in the late expedition to the Rocky Mountains, performed by order of Mr. Calhoun, Secretary of War, under the command of Major Long. *Journal of the Academy of Natural Sciences of Philadelphia*, 3, 238-282; 298-331; 403-462.
- Say, T. (1827) Descriptions of new species of Coleopterous Insects inhabiting the United States. *Journal of the Academy of Natural Sciences of Philadelphia* 5 (2) [1826], 237-284.
- Say, T. (1835) Descriptions of New North American coleopterous insects, and observations on some already described. *Boston Journal of Natural History*, 1(2), 151-203.
- Schaeffer, C.F.A. (1901) [*Derobrachus forreri* and *D. geminatus*]. *Journal of the New York Entomological Society*, 9, 199.
- Schaeffer, C.F.A. (1902) Notes on *Derobrachus* and *Acanthocinus*. *Entomological News*, 13(8), 235-237.
- Schaeffer, C.F.A. (1905) Some additional new genera and species of Coleoptera found within the limit of the United States. *Science Bulletin of the Museum of the Brooklyn Institute of Arts and Sciences*, 1(7), 141-179.
- Schaeffer, C.F.A. (1908) List of the longicorn Coleoptera collected on the Museum expeditions to Brownsville, Texas and the Huachuca Mts., Arizona, with descriptions of new genera and species and notes on known species. *Science Bulletin of the Museum of the Brooklyn Institute of Arts and Sciences*, 1(12), 325-352.
- Schaeffer, C.F.A. (1909) Three Cuban Coleoptera new to the fauna of the United States. *Journal of the New York Entomological Society*, 17(4), 148-150.
- Schaeffer, C.F.A. (1929) The North American species of *Parandra* (Col., Cerambycidae). *Bulletin of the Brooklyn Entomological Society*, 24, 39-40.
- Schaeffer C. F. & Leng C.W. (1912) In Proceedings of the New York Entomological Society. Meeting of March 5, 1912.
- Schaufuss, C. (1916) In: *Calwer's Käferbuch einfuhrung in die Kenntnis der Käfer Europas*. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 2, 709-1390.
- Schaupp, F.G. (1878) The best collecting ground (North Branch, Sullivan Co., N. Y.) *Bulletin of the Brooklyn Entomological Society* 1 (3): 20.
- Schaupp, F. G. (1879) Inhabitants of a beech tree. *Bulletin of the Brooklyn Entomological Society* 2 (3), 24.
- Schiefer, T.L. (1998) A preliminary List of the Cerambycidae and Disteniidae (Coleoptera) of Mississippi *Transactions of the American Entomological Society*, 124 (2), 113-131.
- Schjodte, J.C. (1865) On the classification of Cerambyces, with particular regard to the Danish fauna. *The Annals and Magazine of Natural History*, (3)15, 182-209.
- Schoenherr, C. J. (1817) *Synonymia Insectorum, oder: Versuch einer Synonymia aller bisher bekannten Insecten; nach Fabricii Systema Eleutheratorum &c. geordnet. Mit Berichtigungen und Anmerkungen, wie auch Beschreibungen neuer Arten und illuminirten Kupfern. Erster Band. Eleutherata oder Käfer*. vol. 3; Lewerentzischen Buchdruckerey, Skara, 1(3), xi, 1-506.
- Schoening, E.H. & Tilden, J.W. (1959) *Anoplodera laetifica* (LeConte) and *Ergates spiculatus* LeConte from knobcone pine. *The Pan-Pacific Entomologist*, 35, 167-168.

- Schuh, J. & Mote, D.C. (1948) Insect pests of nursery and ornamental trees and shrubs in Oregon. *Bulletin of the Oregon Agricultural Experiment Station*, 449, 1–163.
- Schwarz, E.A. (1896) [Food-habits of *Atimia confusa*] *Proceedings of the Entomological Society of Washington* 4, 44.
- Schwarz, E.A. (1901) [Communication] *Proceedings of the Entomological Society of Washington* 4 (1), 44.
- Schwarz, E.A. (1917) In: Holland, E. List of the Coleoptera collected on the Isle of Pines by Gustav A. Link, Sr., 1912-1913. *Annals of the Carnegie Museum*, 11, 333–345.
- Schwerdtfeger, F. (1955) Cerambycidae (Col.) an Pinus-Arten in Guatemala *Zeitschrift für angewandte Entomologie*, 37 (3), 315-322, 8 figs.
- Semenov-Tian-Shanskij, A. (1899) Specierum rossicarum generis *Polyarthron* Serv., 1832 synopsis. *Horae Societatis Entomologicae Rossicae*, 34, 249–259.
- Semenov-Tian-Shanskij, A. (1908) Analecta coleopterologica. XIV. *Revue Russe d'Entomologie*, 7 [1907], 258-265.
- Sharp, D. (1905) The genus *Criocephalus*. *The Transactions of the Entomological Society of London*, 1905, 145–164.
- Shenefelt, R.D. & Benjamin, D.M. (1955) Insects of Wisconsin forests *Circular of the Wisconsin University Agriculture College Ext. Service*, 500. 1-110, 28 figs.
- Sheppard, R. W. (1935) Coleopterous collections from Japanese beetle traps in southern Ontario *65th Annual Report of the Entomological Society of the Province of Ontario* 1934, 93-97.
- Sheppard, R.W. (1936) Insects collections from Niagara's colored search-lights *66th Annual Report of the Entomological Society of the Province of Ontario* 1935, 69-75.
- Sherman, F.Jr. (1946) Notes on Cerambycidae of South Carolina (Coleoptera). *Entomological News*, 57(5), 125–127.
- Sikes, D.S. & Webster, R.P. Reginald P. (2005) Bioinventory of Rhode Island Coleoptera: 45 New Records *The Coleopterists' Bulletin* 59 (3). 311-327.
- Skiles, D.D. (1978a) Taxonomy and description of two prionine Cerambycidae from southern Arizona: a new species of *Stenodontes* and new status for *Neomallodon arizonicus* (Coleoptera). *Proceedings of the Entomological Society of Washington*, 80(3), 407–423.
- Skiles, D.D. (1978b) Notes on the larval habits of *Asemum caseyi* Linsley and *A. nitidum* LeConte (Coleoptera: Cerambycidae). *The Pan-Pacific Entomologist*, 54(1), 14.
- Skinner, H. (1901) [*Stenodontes mandibularis*]. *Entomological News*, 12, 93.
- Skinner, H. (1905) Descriptions of new Coleoptera from Arizona with notes on some other species. *Entomological News*, 16(9), 289–292.
- Slosson, A.T. (1894) List of insects taken in alpine region of Mt. Washington. *Entomological News*, 5 (1), 1-6.
- Slosson, A.T. (1895) Additional list of insects taken in alpine region of Mt. Washington. *Entomological News*, 6 (1), 4-7.
- Smith G.A. & Hurley J.E., (2005) First Records in Atlantic Canada of *Spondylis upiformis* Mannerheim and *Xylotrechus sagittatus sagittatus* (Germar) (Coleoptera: Cerambycidae). *The Coleopterists' Bulletin* 59 (4), 488.
- Smith, J.B. (1888) [Host of *Notorhina aspera*]. *Proceedings of the Entomological Society of Washington*, 1(2), 90–91.
- Smith, J.B. (1900) Insects of New Jersey. A list of the species occurring in New Jersey, with notes on those of economic importance. *27th Annual Report of the Board State of Agriculture*, 1899, 1–755.
- <https://doi.org/10.5962/bhl.title.39987>
- Smith, J.B. (1910) Annual report of the insects of New Jersey. *Annual Report New Jersey State Museum*, 1910, 1–888.
- Snow, F.H. (1877) List of Coleoptera collected in Colorado in June, July and August, 1876 by the Kansas University Scientific Expedition. *Transactions of the Kansas Academy of Sciences*, 5, 15–20.
- Snow, F.H. (1878) List of Coleoptera collected near Dome Rock, Platte Canyon, Colorado by the Kansas University Scientific Expedition for 1878. *Transactions of the Kansas Academy of Sciences*, 6, 75–77.

- Snow, F.H. (1881) List of Coleoptera collected in Santa Fé Canyon, N. M. by the Kansas University Scientific Expedition for 1880. *Transactions of the Kansas Academy of Sciences*, 7, 67–71.
- Snow, F.H. (1883) List of Lepidoptera and Coleoptera collected in New Mexico by the Kansas University Scientific Expeditions of 1881 and 1882. *Transactions of the Kansas Academy of Sciences*, 8, 35–45.
- Snow, F.H. (1906a) Some results of the University of Kansas Entomological Expeditions to Galveston and Brownsville, Texas, in 1904 and 1905. *Transactions of the Kansas Academy of Sciences*, 20(1), 136–154.
- Snow, F.H. (1906b) Some results of the University of Kansas Entomological Expeditions to Arizona in 1904 and 1905. *Transactions of the Kansas Academy of Sciences*, 20(1), 155–181.
- Snyder T.E. (1910) Damage to chestnut telephone and telegraph poles by wood-boring insects. *Proceedings of the Entomological Society of Washington* 15, 131-133.
- Snyder T. E. (1913) The ovipositor of *Parandra brunnea* Fab. *USDA Bureau of Entomology Bulletin* 94 (1) 1-12.
- Solomon, J.D. Newsome, L. & Darwin, W.N. (1972) Carpenterworm moths (*Prionoxystus robiniae*: Lep., Cossidae) and cerambycid (Col.) hardwood borers caught in light-traps. *Journal of the Georgia Entomological Society*, 7(1), 76-79.
- Solomon J. D., Doolittle R. E. & Spilman J. T. (1976) Cerambycid beetles captured in sticky-traps in Mississippi. *The Coleopterists' Bulletin* 30 (3) , 289-290.
- Spencer, G.J. (1949) The egg potential of *Ergates spiculatus* LeC. (Coleoptera, Cerambycidae). *Proceedings of the Entomological Society of British Columbia*, 45, 20.
- Spencer, G.J. & Buckell, E.R. (1957) On the acridiophagous Sarcophagidae of British Columbia, with records of all others taken in the Province. *Proceedings of the Entomological Society of British Columbia*, 54, 29–36.
- Spieth, H.T. (1950) The David Rockefeller Mexican Expedition of the American Museum of Natural History. *American Museum Novitates*, 1454, 1–67.
- Spomer, S.M. (2014) The Longhorn Beetles (Coleoptera: Cerambycidae) of Nebraska, USA: new State records, a checklist of known species, and distribution maps *The Coleopterists' Bulletin* 68 (2), 297-315
- Stein J. D. & Tagestad A.D. (1976) The Long-Horned Wood-Boring Beetles of North Dakota (Coleoptera: Cerambycidae) USDA Forest Service. Rocky Mountain Forest and Range Experiment Station Research Paper RM -171 : 1-58.
- Stephens, J.F. (1831) *Illustrations of British Entomology or, a synopsis of indigenous insects; containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food, and economy, as far as possible*. Mandibulata. Baldwin & Craddock, London, 4, 413 pp.
- Steury B.W. & MacRae, T. C. (2014) The Longhorned Beetles (Insecta: Coleoptera: Cerambycidae) of the George Washington Memorial Parkway Virginia Natural History Society. *Banisteria* 44: 7-12.
- Strauch, A. 1861. *Catalogue Systématique de tous les Coléoptères décrits dans les Annales de la Société Entomologique de France depuis 1832 jusqu'à 1859*. H. W. Schmidt, Libraire-Éditeur, Halle, i-iv + 1-160.
- Svácha, P. & Danilevsky, M.L. (1987) Cerambycid larvae of Europe and Soviet Union (Coleoptera, Cerambycoidea). Part I. *Acta Universitatis Carolinae (Biologica)*, 30, 1–176.
- Svácha, P. & Danilevsky, M.L. (1989) Cerambycid larvae of Europe and Soviet Union (Coleoptera, Cerambycoidea). Part III. *Acta Universitatis Carolinae (Biologica)* 32, 1–205.
- Svácha P. & Lawrence J.F., (2014) Arthropoda: Insecta: Coleoptera. Volume 3: Morphology and Systematics (Phytophaga). 2.4 Cerambycidae Latreille, 1802. In Walter de Gruyter, *Handbook of Zoology 3*: i-xii + 1-676, 465 figs Edited by Richard A.B. Leschen & Rolf G. Beutel
- Swan, L.A. & Papp, C.S. (1972) *The common insects of North America*. Harper & Row, New York, 750 pp.

- Swenk, M. H. (1922) Prionus grubs (*Prionus fissicornis* Hald.) *USDA Bureau of Entomology and Plant Quarantine, Insect Pest Survey Bulletin* 2, 200.
- Swift, I.P., Santos-Silva, A., & Nearn, E.H. (2010). The genus *Trichocnemis* LeConte, 1851 (Coleoptera, Cerambycidae, Prioninae). *ZooKeys*, 61, 33-46.
- Tanner, V.M. (1927) A preliminary study of the genitalia of female Coleoptera. *Transactions of the American Entomological Society*, 53, 5-50.
- Tanner, V.M. (1928) The Coleoptera of Zion National Park, Utah. *Annals of the Entomological Society of America*, 21(2), 269-281.
- Tanner, V.M. & Tanner, W.W. (1974) Additional records of Coleoptera collected at the Nevada test site Mercury, Nevada. *The Great Basin Naturalist*, 34(3), 218-220.
- Terrón, R.A. (1992) Fauna de coleópteros Cerambycidae de la Reserva de la Biosfera "La Michilia", Durango, Mexico. *Folia Entomologica Mexicana*, 81, 285-314.
- Thomas, M.C. (1977) New host records and behavior observations on Florida Cerambycidae. *The Coleopterologists' Bulletin*, 31(1), 83-86.
- Thomas, S.L. (1951) *Derobrachus geminatus* on grape roots (Coleoptera: Cerambycidae). *The Pan-Pacific Entomologist*, 27(1), 35.
- Thomson, G.C. (1859) *Skandinaviens Coleoptera, synoptiskt bearbetade*. Lund, 1, 290 pp.
- Thomson, J. (1860) *Essai d'une classification de la famille des cérambycides et matériaux pour servir a une monographie de cette famille*. Bouchard-Huzard, Paris, xvi +128 p.
- Thomson, J. (1861a) *Essai d'une classification de la famille des cérambycides et matériaux pour servir a une monographie de cette famille*. Bouchard-Huzard, Paris, 129-396.
- Thomson, J. (1861b) Monographie de la famille des parandrides. *Musée Scientifique*, 2, 73-87.
- Thomson, J. (1864) Systema cerambycidarum ou exposé de tous les genres compris dans la famille des cérambycides et familles limitrophes. *Mémoires de la Société Royale des Sciences de Liège*, 19, 1-352.
- Thomson, J. (1865) Diagnoses d'espèces nouvelles qui seront décrites dans l'appendix du systema cerambycidarum. *Mémoires de la Société Royale des Sciences de Liège*, 19, 353-578.
- Thomson, J. (1867a) D'une classification nouvelle de la famille des cérambycides (Insectes, coléoptères). *Physis Recueil d'Histoire Naturelle*, 1(1), 1-10.
- Thomson, J. (1867b) Révision des parandrides (Insectes coléoptères). *Physis Recueil d'Histoire Naturelle*, 1(2), 106-118.
- Thomson, J. (1867c) Révision du groupe des mallodonites (Insectes coleoptères, prionites, cérambycides). *Physis Recueil d'Histoire Naturelle*, 1(2), 85-106.
- Thomson, J. (1878a) *Typi cerambycidarum Musei Thomsoniani*. E. Deyrolle, Paris, 21 pp.
- Toledo, V.H., Noguera, F.A., Chemsak, J.A., Hovore, F.T. & Giesbert, E.F. (2002) The Cerambycid Fauna of the tropical dry forest of "El Aguacero", Chiapas, Mexico (Coleoptera, Cerambycidae). *The Coleopterists' Bulletin*, 56(4), 515-532.
- Touroult, J. (2007) Contribution à la connaissance des longicornes de Sainte Lucie et de Martinique (Coleoptera, Cerambycidae). *Les Cahiers Magellanes*, 70, 1-14.
- Touroult, J. (2012) Longicornes des Petites Antilles: mise à jour des connaissances. *ACOREP France, Coléoptères des Petites Antilles*, 1, 70-85.
- Townes, H.K. (1944) A Catalogue and Reclassification of the Nearctic Ichneumonidae (Hymenoptera) *Memoirs of the American Entomological Society* 11 (1), 1-800.
- Townes, H. & Townes, M. (1960) Ichneumon-flies of America, north of Mexico. 2. Subfamilies Ephialtinae, Xoridinae, Acaenitinae *Bulletin of the United States National Museum*, 216 (2), i-vii + 1-676, 378 figs.
- Townsend, C.H.T. (1892) Biologic notes on New Mexico insects. *Insect Life*, 5(1), 37-40.
- Townsend, C.H.T. (1895) On the Coleoptera of New Mexico and Arizona, including biologic and other notes. *The Canadian Entomologist*, 27(2), 39-51.
- Townsend, C.H.T. (1903) Contribution to a knowledge of the coleopterous fauna of the lower Rio Grande Valley in Texas and Tamaulipas, with biological notes and special reference to geographic distribution. *Transactions of the Texas Academy of Sciences*, 5, 49-101.
- Treat, M.L. (1874) The enemies of the oak *American Agriculture* 33, 344, 5 figs.
- Tucker, E.S. (1906) Determinations of some Texas Coleoptera with records. *Entomological News*, 17, 10-14.

- Tucker, E.S. (1907) Concerning some insects collected and bred from dead and dying elm. *Transactions of the Kansas Academy of Science*, 21 (1), 158-162.
- Turienzo, P. (2007) Nuevo registro y primera planta hospedadora conocida para *Arhopalus rusticus rusticus* (Linnaeus, 1758) (Coleoptera, Cerambycidae) en Argentina. *Boletín de Sanidad Vegetal*, Plagas, 33, 321-323.
- Turnbow, R.H., Cave, R.D. & Thomas, M.C. (2003) A list of the Cerambycidae of Honduras, with additions of previously unrecorded species. *Ceiba* 44(1), 1-43.
- Turnbow R.H. & Franklin R.T. (1980) An annotated checklist of the Cerambycidae of Georgia (Coleoptera). *Journal of the Georgia Entomological Society* 15 (3), 337-349.
- Turnbow, R.H. & Wappes, J.E. (1978) Notes on Texas Cerambycidae (Coleoptera). *The Coleopterists' Bulletin*, 32(4), 367-372.
- Tyson, W.H. (1966) Notes on reared Cerambycidae. *The Pan-Pacific Entomologist*, 42(3), 201-207.
- Tyson, W.H. (1967) California *Ergates* and *Tragosoma*, with keys to the adult and immature forms (Coleoptera, Cerambycidae). *The Pan-Pacific Entomologist*, 43(2), 122-126.
- Tyson, W.H. (1970) The Cerambycidae of the Panamint Mountains, California (Coleoptera). *The Pan-Pacific Entomologist*, 46(4), 296-299.
- Ulke, H. (1903) A list of the beetles of the District of Columbia. *Proceedings of the United States National Museum*, 25(1275), 1-57.
- Van Dyke, E.C. (1934) A root-boring *Derobrachus*. *The Pan-Pacific Entomologist*, 10(2), 58.
- Van Dyke, E.C. (1937) Notes and descriptions of North American Buprestidae and Cerambycidae (Coleoptera). *Bulletin of the Brooklyn Entomological Society*, 32(3), 105-116.
- Villiers, A. (1980a) Une collection de coléoptères Cerambycidae de la Dominique appartenant à l'U. S. National Museum (Washington). *Annales de la Société Entomologique de France*, (n.s.) 16(1), 129-131.
- Villiers, A. (1980b) Coléoptères Cerambycidae des Antilles françaises. I. Parandrinae, Prioninae, Lepturinae. *Annales de la Société Entomologique de France*, (n.s.) 16(1), 133-157.
- Vitali, F. & Rezbanyai-Reser, L. (2003) Beiträge zur Insektenfauna von Jamaika, Westindien (Karabik) 5. Bockkäfer - 1 (Coleoptera, Cerambycidae). *Les Cahiers Magellanes*, 26, 1-16.
- Vlasák, J. (2014) New Larval Host Records for North American Cerambycidae (Coleoptera) *The Coleopterists' Bulletin* 68 (2), 316-320, 1 fig.
- Vlasák, J. & Vlasakova K. (2002) Records of Cerambycidae (Coleoptera) in Massachusetts with notes on larval hosts *The Coleopterists' Bulletin* 56 (2): 203-219.
- Vlasák, J. & Vlasakova K. (2021) New larval host plants and ecological observations on North American Cerambycidae (Coleoptera). *Insecta Mundi*. 0901, 1-23
- Vogt, G.B. (1949) Notes on Cerambycidae from the Lower Rio Grande Valley, Texas (Coleoptera). *The Pan-Pacific Entomologist*, 25(3), 137-144; (4), 175-184.
- Walsh, B.D. (1866) Answers to correspondents *Practical Entomology* 1, 89-90.
- Walsh B.D. & Riley C.V. (1868a) A new grape root-borer *The American Entomologist* 1 (1): 19.
- Walsh, B.D. & Riley C.V. (1868b) Grape vine borer *The American Entomologist* 1 (2): 40.
- Ware, R. E. (1929) Some notes on collecting Cerambycidae. *Proceedings of the Iowa Academy of Sciences* 36, 367-369.
- Warren, J.C. (1899) [Coleoptera from Tioga County, Pennsylvania] *Entomological News*, 10 (10), 295-296.
- Waters, D. L. & Hyche, L.L. (1984) Notes on Cerambycidae (Coleoptera) collected on dead or stressed deciduous trees in East Central Alabama *The Coleopterists' Bulletin* 38 (3), 283-287.
- Webster, F.M. (1879) Entomological Notes. *Bulletin of the Brooklyn Entomological Society* 2 (2), 20.
- Webster, F.M. (1892) Insects affecting the hackberry and raspberry. *Bulletin of the Ohio Agricultural Experiment Station* 45 (8), 151-217
- Webster, F.M. (2016) Checklist of the Coleoptera of New Brunswick, Canada. *ZooKeys*, 573-512
- Webster, R.P., McCorquodale D.B. & Majka C.G. (2009) New records of Cerambycidae (Coleoptera) for New Brunswick, Nova Scotia, and Prince Edward Island, Canada. *ZooKeys* 22, 285-308.

- Webster, R.P., Sweeney, J.D., De Merchant I., Silk P.J. & Mayo P. (2012) New Coleoptera records from New Brunswick, Canada: Cerambycidae *ZooKeys* 179, 309-319.
- Webster R.P., Alderson, C.A., Webster, V. L., Hughes C, C. & Sweeney, J.D., 2016 Further contributions to the longhorn beetle (Coleoptera, Cerambycidae) fauna of New Brunswick and Nova Scotia, Canada. *ZooKeys* 552 109-122.
- Weed, C. M. (1884) Grape root borers *Prairie Farmer*, December 1884, 13.
- Wendt, H. (1984) Die Cerambyciden-Typen (Coleoptera, Phytophaga) des Zoologischen Museums Berlin. Teil I. Unterfamilie Prioninae. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 2, 327–342.
- Westcott R. L., LaBonté J. R., Parsons G. L. & Johnson. Paul P.J. (2006) New records and other notes for Oregon Coleoptera Zootaxa, 1142, 1-33..
- Westwood J.O. (1838) Synopsis of the genera of British insects [pp. 1–48]. In: An introduction to the modern classification of insects, founded on the natural habits and corresponding organization of the different families. Vol. II. Longman, Orme, Brown, Green & Longmans, London. [7 May (pp. 1–16); 9 Aug (17–32); 5 Nov 1838 (33–48).
- White, A. (1853) *Catalogue of the coleopterous insects in the collection of the British Museum*. Longicornia 1. British Museum, London, 7,174 pp.
<https://doi.org/10.5962/bhl.title.60043>
- White, A. (1855) *Catalogue of the coleopterous insects in the collection of the British Museum*. Longicornia 2. British Museum, London, 8, 175–412.
- White R.A., Paim, U. & Seabrook, W.D. (1974) Maxillary and labial sites of carbon dioxide-sensitive receptors of larval *Orthosoma brunneum* (Forster) (Coleoptera, Cerambycidae). *Journal of Comparative Physiology* 88 (3), 235-246.
- Wickham, H.F. (1893) Field notes from Texas and Louisiana. *The Canadian Entomologist*, 25, 139–143.
- Wickham, H.F. (1897a) Recollections of old collecting grounds. I. The Coeur d'Alene district. *Entomological News*, 8(7), 165–168.
- Wickham, H.F. (1897b) The Coleoptera of Canada. The Cerambycidae of Ontario and Quebec. *The Canadian Entomologist*, 29(4), 81–88; (5), 105–111; (6), 148–153; (7), 169–173; (8), 187–193; (9), 201–208.
- Wickham, H.F. (1897c) A list of Coleoptera from the southern shore of Lake Superior, with remarks on geographical distribution. *Proceedings of the Davenport Academy of Natural Sciences* 6, 125-169.
- Wickham H.F. (1899) Recollections of old collecting grounds. IX. The Alpine Districts about Leadville. *Entomological News*, 10, 196-199.
- Wickham, H.F. (1909a) A list of the Coleoptera of Iowa. *Bulletin of the Laboratory of Natural History*, State University of Iowa, 6(2), 1–40.
- Wickham, H.F. (1909b) A list of the Van Duzee collection of Florida beetles. *Bulletin of the Buffalo Society of Natural History*, 9(3), 399–405.
- Wolcott A.B. & Montgomery B.E. (1933) An ecological study of the coleopterous fauna of a tamarack swamp. *American Midland Naturalist* 14 (2), 113-169.
- Yanega, D. (1996) Field Guide to Northeastern Longhorned Beetles (Coleoptera: Cerambycidae) *Illinois Natural History Survey Manual* 6: 1-174
- Zagajkevich. I. K. (1991) Taksonomiya i ekologiya usachey [Taxonomy and Ecology of Longicorn Beetles] Académie des Sciences de l'Ukraine & Conseil Scientifique de Kiev , 1-180, 20 figs.
- Zajciw, D. (1970) Notas sobre o gênero *Strongylaspis* Thoms., 1860, com descrição de uma espécie nova do Brasil (Coleoptera, Cerambycidae, Prioninae). *Atas da Sociedade de Biologia*, 14(1–2), 1–3.
- Zayas, F.de. (1957) Revisión de los longicórnios prionidos de Cuba (Coleópteros, Cerambycidae). *Memorias de la Sociedad Cubana de Historia Natural*, 23, 149–181.
- Zayas, F.de. (1975) *Revisión de la familia Cerambycidae* (Coleoptera, Phytophagoidea). Academia de Ciencias de Cuba, Habana, 443 pp.
- Zimsen, E. (1964) *The type material of I. C. Fabricius*. Muuksgaard, Copenhagen, 656 pp.

